Q1.

The growth of daisy plants on a lawn is affected by biotic factors and by abiotic factors.


(a) The table below shows six factors.

Tick (\checkmark) one box in each row to show whether the factor is biotic or abiotic.

Factor	Biotic	Abiotic
Nitrates in the soil		
Rabbits eating the plants		
Shading by a building		
Soil pH		
Temperature		
Trampling by people		

(3)

The figure below shows a plan of a garden.



A student estimates the number of daisy plants growing on the lawn.

The student places a quadrat at 10 different positions on the lawn.

The quadrat measures 50 cm \times 50 cm.

The student counts the number of daisy plants in each quadrat.

(b) How should the student decide where to place the quadrat?


AQA Biology GCSE - Adaptation, Interdependence and Competition

_	
Γ	he mean number of daisy plants in each quadrat is 6.
C	Calculate the number of daisy plants on the lawn.
	Give your answer to 3 significant figures.
_	
_	
_	
_	
	lumber of daisy plants on the lawn =
	Ising the mean from this investigation to calculate the number of daisy lants on the lawn may not be accurate.
(Give two reasons why.
1	

(2) (Total 13 marks)

Q2.


The image below shows what the extinct Siberian rhinoceros (*Elasmotherium sibiricum*) might have looked like.



(a)	What is the genus of the Siberian rhinoceros?		
	Tick (✓) one box.		
	Elasmotherium		
	Elasmotherium sibiricum		
	sibiricum		
			(1)
The 'three-domain system' of classification places all living organisms in one of three domains.			
(b)	Which domain was the Siberian	rhinoceros in?	
	Tick (√) one box.		
	Archaea		

Eukaryota	
Prokaryota	
Mho dovolonos	I the 'three domain evetem' of election?
•	the 'three-domain system' of classification?
Tick (√) one bo Carl Woese	DX.
Charles Darwi	n
Gregor Mende	
	Siberian rhinoceros is estimated to have been 150 cm long
Suggest one a	dvantage of this adaptation to the Siberian rhinoceros.
The only parts o	of the Siberian rhinoceros that have been found are s.
	on why only the bones of the body of the Siberian
hinoceros bec	ame rossiis.
	cientists can estimate when the Siberian rhinoceros was
Suggest how sc alive.	eientists can estimate when the Siberian rhinoceros was

The below diagram shows when the Siberian rhinoceros existed and when modern humans existed.



How many million years ago did the Siberian rhinoceros become extinct? million years ago
Determine the time in years when both the Siberian rhinoceros and modern humans existed together.
Use the diagram above and your answer to Question (g).
Time = years
Suggest two factors that may have caused the extinction of the Siberian
rhinoceros.

(Total 12 marks)


Q3.					
Li	Living organisms are classified into the following groups:				
•	• Kingdom				
•	Phylum				
•	Class				
•	Order				
•	Family				
•	Genus				
•	Species				
(a) Which scientist first sugg	gested this type of classification system?			
	Tick one box.				
	Alfred Russel Wallace				
	Carl Linnaeus				
	Charles Darwin				
	Gregor Mendel				
			(1)		

The stone plant, *Lithops bromfieldi*, is adapted to live in very dry deserts.

Figure 1 shows several stone plants.

(1)

Figure 1



(b) Give the genus to which the stone plant belongs.

(c)

The stone plant has many adaptations that help it to survive in the desert.

Draw **one** line from each adaptation to how the adaptation helps the stone plant to survive.

How the adaptation helps Adaptation survival Can trap a lot of light Absorb water from deep in Plants look like stones the ground Leaves with thick, waxy Help cross-pollination cuticles Are not easy to see and Many long, branching roots so are not eaten Thick, fleshy leaves Reduce water loss Store water

(Total 9 marks)

(4)

The jerboa is a small desert animal.

Figure 2 shows a jerboa.

Figure 2



The jerboa is adapted for survival in the desert.

The jerboa spends the daytime in its underground burrow.

The jerboa only leaves its burrow to look for food during the night.

Describe how the	nese adaptations help the jerboa to survive in the desert.
What type of ac	laptations are described in Question (d)?
Γick one box.	
Behavioural	
Functional	
Structural	


(2)

Q4.

Figure 1 shows a type of camel called a dromedary (Camelus dromedarius).

The dromedary lives in hot, dry deserts.

Figure 1



(a) One adaptation of the dromedary is 'temperature tolerance'.

This means that the animal's body temperature can rise by up to 6 °C before it starts to sweat.

Explain how temperature tolerance can help the dromedary to survive in the desert.

(b) Three more adaptations of the dromedary are given in **Figure 1**.

Give a reason why each adaptation helps the animal survive in the desert.

Fat store _____

Produces little urine and very dry faeces _____

Hard mouth	
	(3)


There are several species of the camel family alive today.

Scientists think these species evolved from a common ancestor that lived in North America about 45 million years ago (Mya).

Figure 2 shows:

- where four modern species of the camel family live today
- how the ancestors of these camels migrated from North America.

Figure 2



(c)	Which two of the four modern species of camel do scientists believe to be
	most closely related to each other?

Give the reason for your answer.

	 	and	 	
Reason	 		 	

(1)

(d)	Describe the type of evidence used for developing the theory of camel migration shown in Figure 2 .			
		(2)		
(e)	Explain how several different species of camel could have evolved from a common ancestor over 45 million years.			
		(6)		
	(Total 14 n	narke)		