
(2)

Q1.

People eat fish caught in the North Sea.

Figure 1 shows a food chain.

Figure 1



	Algae	Small animals	Herring	Humar
				Not to scale
(a)	The algae make g	lucose by photos	synthesis.	
	Which two substa	inces do the alga	ae need for photosyn	thesis?
	Tick (✓) two boxe	S.		
	Carbon dioxide			
	Nitrogen			
	Oxygen			
	Starch			
	Water			
(b)	What is the source	e of energy for pl	notosynthesis?	

Tick (✓) one box.

Light	
Mineral ions	3





Figure 2 shows the biomass of adult herring in the North Sea between 1950 and 1990.

Figure 2



(d) Too many herring were caught in the 1960s.

Calculate the percentage decrease in the biomass of adult herring between 1960 and 1970.

Use the equation:

percentage decrease =
$$\frac{\text{(biomass in 1960 - biomass in 1970)}}{\text{biomass in 1960}} \times 100$$

Give your answer to the nearest whole number.

(Total 12 marks)


1977, laws were introduced to help conserve herring. Describe the change in biomass of adult herring from 1977 to 1990.			
Use data from Figure 2 in your answer.			
One of the laws was to control mesh size of fishing nets.			
Figure 3 shows a fishing net with a legal mesh size.			
Figure 3			
Mesh size 8-year-old herring 4-year-old herring			
Herring can live for up to 12 years.			
Herring start to reproduce when they are 3 to 4 years old.			
Explain how the control of mesh size of fishing nets has helped to conserve stocks of herring.			

Q2.

A food web contains several food chains.

(d)

Figure 1 shows a food web.




Describe how the algae get energy.	
Name one primary consumer in Figure 1 .	
Name one producer in Figure 1 .	

The different food chains in **Figure 1** have different numbers of organisms.

(1)

(2)

Complete **Figure 2** to show a food chain in **Figure 1** with **five** organisms, including the human.



(e) Figure 1 shows that mackerel eat krill and squid.

The biomass of mackerel is much less than the combined biomass of krill and squid.


One reason for this is that the mackerel cannot digest all parts of the krill and squid.

Give two other reasons.

1	
2	_

Figure 3 shows how the biomass of adult herring in the North Sea has changed between 1950 and 2010.

Figure 3



(f) Calculate the percentage decrease in the biomass of herring between 1960 and 1977.

Give your ans	swer to the	nearest w	noie numi	Jei.	

Percentage decrease = _

(4)

(g) Too many herring were caught by fishermen between 1960 and 1977.

Herring can live for up to 12 years and begin to reproduce when 3 to 4 years old.

Laws have been introduced to help conserve herring:

- 1977 to 1981 herring fishing was banned in the North Sea
- 1984 to present day control of mesh size of fishing nets
- 1997 to present day fishing quotas were introduced
- 1998 to present day herring fishing was banned in breeding grounds during the breeding season.

Figure 4 shows how a minimum mesh size helps to conserve herring.





Figure 3 is repeated below.

Figure 3




Evaluate the effect of these laws on the conservation of herring stocks.
Use data from Figure 3 and information from Figure 4 in your answer.

 (0)
(6)
(Total 17 marks)

Q3.

A food for pet dogs contains meat from chickens.

The below diagram shows the food chain.



(a) What is the trophic level of the dog?


Tick (\checkmark) one box.

1 2 3

(1)

(b) Draw **one** line from each organism to the description of the organism's position in the food chain.

	Organism	Description	
		Herbivore	
	Chicken		
		Producer	
	Dog		
		Secondary consumer	
	Wheat		
		Tertiary consumer	
(c)	Name the process wheat pla	ants use to make glucose.	(3)
(d)	Some of the chicken biomas What is one reason why?	ss does not become part of the dog's biomass.	(1)
	Tick (✓) one box.		
	Some of the chicken is use	d for the dog to grow	
	The dog produces waste in	faeces	
	The wheat is eaten by the	dog	(1)
A ne	w dog food has been develop	ped.	(-,
The	new dog food is made from ir	nsects.	
The	insects in the dog food factor	y are fed on vegetables.	
(e)	Which pyramid of biomass r this food chain?	represents the vegetables, insects and dogs in	
	Tick (✓) one box.		



Q4.

A new dog food has been developed that does **not** contain meat from cows, sheep or chickens.

The new dog food contains insects.

The insects in the dog food factory are fed on waste vegetables.

Sketch the pyramid of biomass for the food chain that produces food for dogs from insects. Label the pyramid. (2) Describe two reasons why the biomass of the insects eaten by dogs does not all become biomass of the dogs. (2) Explain how making dog food from insects could improve human food (c) security in the future.

nymph

 -	 	
(4)		
(Total 8 marks)		

Q5.

Figure 1 shows a food chain in a pond.

Algae Daphnia Hydra Dragonfly

Figure 1

(a) Which term describes the Daphnia in this food chain?

Tick (✓) one box.

Apex predator

Primary consumer

Producer

Secondary consumer

(1)

(b) Draw a pyramid of biomass for the food chain.

Label each trophic level.

(2)

(c)	Give one reason why the total biomass of the Daphnia in the pond is different from the total biomass of the algae.			
Stud	ents investi	gated the size o	of the population of Daphnia	a in the pond.
This	is the metho	od used.		
1. C	ollect 1 dm ³	of pond water t	from near the edge of the po	ond.
2. P	our the wate	r through a fine	e net.	
3. C	ount the nun	nber of Daphni	a caught in the net.	
4. R	epeat steps	1–3 four more	times.	
The	table below	shows the resu	ults.	
		Sample number	Number of Daphnia in 1 dm³ water	
		1	5	
		3	21	
		4	16	
		5	28	
(d)	Calculate t	he mean numb	per of Daphnia in 1 m³ of po	nd water.
	$1 \text{ m}^3 = 100$	0 dm³		
	-			

(e) The pond was a rectangular shape, measuring:

Mean number of Daphnia in 1 m³ of pond water = _____

- length = 2.5 metres
- width = 1.5 metres

AQA Biology GCSE - Trophic Levels in an Ecosystem

• depth = 0.5 metres.	
Calculate the estimated number of Daphnia in the pond.	
Use your answer from part (d).	
Give your answer in standard form.	
Number of Daphnia in the pond =	•
	(4


Rainfall can cause fertiliser to be washed from farmland into a pond.

The students investigated the effect of fertiliser on the population of Daphnia in water from the pond.

- The students put 20 Daphnia in each of five different concentrations of fertiliser.
- The students counted the total number of Daphnia in each concentration of fertiliser after 2 weeks.

Figure 2 shows the results.

Figure 2




(f) A concentration of 5.0 mg/dm³ of fertiliser caused a large increase in the population of Daphnia.

Explain	why
---------	-----

(2)

(g) Figure 1 is repeated below.

Figure 1




The population of **Hydra** will decrease when 20 mg/dm³ of fertiliser is added to the pond.

Explain why.


(2)		
(2)		
(Total 14 marks)		

Q6.

The diagram below shows a food chain in a river.



(a) Draw **one** line from each scientific term to the correct organism in the food chain.



(b) The table below shows the biomass of the organisms at each stage in the food chain.

Organism	Biomass in arbitrary units
Algae	840
Invertebrate animals	200
Small fish	40
Large fish	10

Calculate the percentage of the biomass of the invertebrate animals that is transferred to the large fish.

percentage = -	biomass of la biomass of inverted	rge fish ×100	
, t	piomass of inverte	brate animals	
	P	ercentage =	
		-	
A large amount of biom	ass is lost from the	e food chain.	
Complete the sentence	es.		
Choose answers from	the box.		
coordination	digestion		
	uigestion	excretion	
filtration When the small fish ea broken down during	ingestion t the invertebrate a	respiration animals, not all of this	material is
filtration When the small fish ea	ingestion t the invertebrate a m the gut may ente	respiration animals, not all of this respiration animals, not all of this response to the ser the body cells of the response to the response	small fish
filtration When the small fish ear broken down during Materials absorbed from these materials are browster by The carbon dioxide and	ingestion It the invertebrate a In the gut may enter In the down into can Industry the control of the contr	respiration animals, not all of this respiration animals, not all of this response to the respiration t	small fish
filtration When the small fish ear broken down during Materials absorbed from these materials are browater by The carbon dioxide and removed from the small fish by	ingestion It the invertebrate a m the gut may ente oken down into ca d other waste mate	respiration animals, not all of this respiration animals, not all of this response to the respiration t	small fish
filtration When the small fish early broken down during Materials absorbed from these materials are browster by The carbon dioxide and removed	ingestion It the invertebrate a If the gut may enter oken down into can If the small fish.	respiration animals, not all of this respiration animals, not all of this response to the respective to the response to the response to the response to the re	small fish

(4)

Q7.

Figure 1 shows:

- a food chain for organisms in a river
- the biomass of the organisms at each trophic level.

Figure 1

Algae Invertebrate Small fish Large fish animals

Biomass in g/m²: 840 200 40 10

(a) Draw a pyramid of biomass for the food chain in **Figure 1** on **Figure 2**.

You should:

- use a suitable scale
- label the x-axis
- label each trophic level.

Figure 2

(b) Calculate the percentage of the biomass lost between the algae and the large fish.

Give your answer to 2 significant figures.

AQA Biology GCSE - Trophic Levels in an Ecosystem

Percentage loss =			
Give one way that biomass is lost between trophic levels. A large amount of untreated sewage entered the river. Many fish died. Untreated sewage contains organic matter and bacteria. Explain why many fish died.			
Give one way that biomass is lost between trophic levels. A large amount of untreated sewage entered the river. Many fish died. Untreated sewage contains organic matter and bacteria. Explain why many fish died.			
A large amount of untreated sewage entered the river. Many fish died. Untreated sewage contains organic matter and bacteria. Explain why many fish died.		Percentage loss =	
A large amount of untreated sewage entered the river. Many fish died. Untreated sewage contains organic matter and bacteria. Explain why many fish died.	Give one way that bio	omass is lost between trophic levels.	
Untreated sewage contains organic matter and bacteria. Explain why many fish died.			
Untreated sewage contains organic matter and bacteria. Explain why many fish died.			
Untreated sewage contains organic matter and bacteria. Explain why many fish died.	A large amount of uni	treated sewage entered the river. Many fish died	
Explain why many fish died.			
		sh died.	
		sh died.	— —
		sh died.	

(3)

(Total 13 marks	

Q8.

Cows are reared for meat production.

The cows can be reared indoors in heated barns, or outdoors in grassy fields.

The table shows energy inputs and energy outputs for both methods of rearing cows.

	kJ / m² / year		
	Energy	y input	Energy output
	Food	Fossil fuels	Meat production
Indoors	10 000	6 000	40
Outdoors	5 950	50	Х

(a) The percentage efficiency for rearing cows **outdoors** is 0.03%

Calculate the energy output value X.

Use the equation:

(b) The percentage efficiency for rearing cows **outdoors** is 0.03%

AQA Biology GCSE - Trophic Levels in an Ecosystem

Give two ways in which the energy is wasted. 1. 2. Suggest two reasons why it is more efficient to rear cows indoors than to
A large amount of energy is wasted in both methods of rearing cows. Give two ways in which the energy is wasted. 1. 2. Suggest two reasons why it is more efficient to rear cows indoors than to
Answer = time A large amount of energy is wasted in both methods of rearing cows. Give two ways in which the energy is wasted. 1. 2. Suggest two reasons why it is more efficient to rear cows indoors than to rear cows outdoors.
A large amount of energy is wasted in both methods of rearing cows. Give two ways in which the energy is wasted. 1. 2. Suggest two reasons why it is more efficient to rear cows indoors than to
A large amount of energy is wasted in both methods of rearing cows. Give two ways in which the energy is wasted. 1. 2. Suggest two reasons why it is more efficient to rear cows indoors than to
A large amount of energy is wasted in both methods of rearing cows. Give two ways in which the energy is wasted. 1. 2. Suggest two reasons why it is more efficient to rear cows indoors than to
A large amount of energy is wasted in both methods of rearing cows. Give two ways in which the energy is wasted. 1. 2. Suggest two reasons why it is more efficient to rear cows indoors than to
A large amount of energy is wasted in both methods of rearing cows. Give two ways in which the energy is wasted. 1. 2. Suggest two reasons why it is more efficient to rear cows indoors than to
A large amount of energy is wasted in both methods of rearing cows. Give two ways in which the energy is wasted. 1. 2. Suggest two reasons why it is more efficient to rear cows indoors than to
Give two ways in which the energy is wasted. 1. 2. Suggest two reasons why it is more efficient to rear cows indoors than to
Give two ways in which the energy is wasted. 1. 2. Suggest two reasons why it is more efficient to rear cows indoors than to
1. 2. Suggest two reasons why it is more efficient to rear cows indoors than to
2. Suggest two reasons why it is more efficient to rear cows indoors than to
2. Suggest two reasons why it is more efficient to rear cows indoors than to
Suggest two reasons why it is more efficient to rear cows indoors than to
Suggest two reasons why it is more efficient to rear cows indoors than to
Suggest two reasons why it is more efficient to rear cows indoors than to
real cows outdoors.
1.
2.

Q9.

The diagram below shows a food chain in a garden.

	Lettuce → Snail → Shrew
e©c a)	destillat/iStock/Thinkstock; Snail ©Valengilda/iStock/Thinkstock; Shrew © GlobalT/iStock/Thinkstock; Shrew © GlobalT/iStock/Thinkstock/Thin
b)	Name one carnivore shown in the diagram above.
c)	A disease kills most of the shrews in the garden.
	Suggest why the number of snails in the garden may then increase.
d)	What is the name given to all the snails in the garden shown in the diagram above?
d)	
d)	above?
d)	above? Tick one box.
d)	above? Tick one box. Community
(d)	above? Tick one box. Community Ecosystem
(d)	above? Tick one box. Community Ecosystem Population

Shrew Snail Lettuce	Shrew Snail Lettuce	Shrew Snail Lettuce
A	В	c
Some snails ate som	ne lettuces.	
The lettuces contain	ed 11 000 kJ of energy.	
Only 10% of this end	ergy was transferred to the s	snails.
Calculate the energy	y transferred to the snails fro	om the lettuces.
	Energy =	kJ
Give one reason whe to the snails.	ny only 10% of the energy in	the lettuces is transferred
	ny only 10% of the energy in	the lettuces is transferred
to the snails.		the lettuces is transferred
to the snails. Tick one box. The lettuces carry of		the lettuces is transferred
to the snails. Tick one box. The lettuces carry of	out photosynthesis at the roots of the lettuces	the lettuces is transferred
to the snails. Tick one box. The lettuces carry of the snails do not each of the snails.	out photosynthesis at the roots of the lettuces	the lettuces is transferred
to the snails. Tick one box. The lettuces carry of the snails do not each of the snails.	out photosynthesis at the roots of the lettuces hail can be eaten affect the food chain.	the lettuces is transferred


Q10.

Figure 1 shows how energy and biomass pass along a food chain.

(2)

(1)

Figure 1




(a)	The parsley shown in Figure 1 carries out photosynthesis.

(b) Which diagram shows the pyramid of biomass for the food chain in **Figure 1**?

Why is photosynthesis important in the food chain?

Tick (**√**) one box.




(c) **Figure 2** shows the ways a swallowtail caterpillar transfers 20 J of energy from food.

(2)

(2)


Figure 2



What percentage of the energy in the caterpillar's food is used for growth?
Percentage =

- (d) The organisms in the food chain are adapted for survival.
 - (i) Figure 3 shows a swallowtail caterpillar seen from the back.

Figure 3



to reduce the chance of being eaten by blue tits.							

Suggest how the swallowtail caterpillar shown in Figure 3 is adapted

(ii) Figure 4 shows a hawk.

Figure 4



Suggest two ways that the hawk is adapted to catch and kill blue tits.

l.	
2.	

(2) (Total 9 marks)

Blue tit: ©JensGade/iStock Parsley: © Warren_Price/iStock Caterpillar ©prettyzhizhi/iStock Hawk: © kojihirano/iStock Swallowtail caterpillar: © Anna_Po/iStock

Q11.

Students investigated a food chain in a garden.

lettuce \rightarrow snail \rightarrow thrush (bird)

The students:

- estimated the number of lettuce plants in the garden
- estimated the number of snails feeding on the lettuces
- counted two thrushes in the garden in 5 hours.

The table below shows the students' results and calculations.

Organism	Population size	Mean mass of each organism in g	Biomass of population in g	Biomass from previous organism that is lost in	Percentage of biomass lost
----------	-----------------	--	----------------------------	--	----------------------------

(a)

				g	
Lettuce	50	120.0	6000		
Snail	200	2.5	500	5500	91
Thrush	2	85.0	170	330	66

•	
	Scientists estimate that about 90% of the biomass in food is lost at each step in a food chain.
	Suggest one reason why the students' value for the percentage of biomass lost between the snails and the thrushes is only 66%.

(b) European banded snails have shells with different colours (light or dark) and with stripes or with no stripes.

Figure 1 shows two examples of European banded snails.





Figure 2 shows results from surveys in woodlands and in grasslands of the percentage of snails with light-coloured shells and the percentage of snails with no stripes.

Each point on the graph represents the results of one survey in one habitat.

Figure 2



Figur	e 2 is a scatter graph.
Why i	s a scatter graph used for this data?
	pare the general appearance of snails that live in woodlands with eneral appearance of snails that live in grasslands.

(iii) Suggest a reason for the general appearance of snails that live in woodlands.

			(1)
		(Total 7 r	marks)

AQA Biology GCSE - Trophic Levels in an Ecosystem

PhysicsAndMathsTutor.com