

CIE Chemistry A-Level Topic 18 - Carbonyl Compounds

Flashcards

This work by PMT Education is licensed under CC BY-NC-ND 4.0











How are primary alcohols oxidised to aldehydes?







How are primary alcohols oxidised to aldehydes?

If a primary alcohol is heated with acidified potassium dichromate(VI), the alcohol will be oxidised to form an aldehyde.

$$CH_3CH_2OH + [O] \rightarrow CH_3CHO + H_2O$$

The acidified potassium dichromate(VI) will change colour from orange to green.









How are secondary alcohols oxidised to ketones?









How are secondary alcohols oxidised to ketones?

If a secondary alcohol is heated under reflux with acidified potassium dichromate(VI), the alcohol will be oxidised to form a ketone:

$$CH_3CH(OH)CH_3 + [O] \rightarrow CH_3COCH_3 + H_2O$$

The acidified potassium dichromate(VI) will change colour from orange to green.









How are aldehydes and ketones reduced using cyanide ions?











How are aldehydes and ketones reduced using cyanide ions?

By using HCN in ethanol or NaCN solution:

$$CH_3CHO + HCN \rightarrow CH_3CH(OH)CN$$

$$CH_3COCH_3 + HCN \rightarrow CH_3C(OH)CNCH_3$$









How are aldehydes and ketones reduced using NaBH, or LiAlH, ions?









How are aldehydes and ketones reduced using NaBH₄ or LiAlH₄ ions?

NaBH₄ and LiAlH₄ are reducing agents and can both be represented as [H] in chemical equations:

$$CH_3CHO + 2[H] \rightarrow CH_3CH_2OH$$









What is the mechanism for the reaction between HCN and aldehydes and ketones?








What is the mechanism for the reaction between HCN and aldehydes and ketones?

Nucleophilic addition











What is the test for a carbonyl compound?











What is the test for a carbonyl compound?

Brady's reagent / 2,4-dinitrophenylhydrazine (2,4-DNPH):

- Add a few drops of the unknown sample to 5 cm³ of Brady's reagent.
- The presence of a carbonyl compound is indicated by a yellow/orange precipitate.









How can you test for an aldehyde using Fehling's solution?











How can you test for an aldehyde using Fehling's solution?

Warm the sample with Fehling's solution.

If an aldehyde is present, the blue solution will form a brick red precipitate.









How can you test for an aldehyde using Tollens' reagent?











How can you test for an aldehyde using Tollens' reagent?

Add Tollens' reagent to the sample and warm in a water bath.

The presence of a silver mirror on the test tube indicates that the sample contains an aldehyde functional group.









How do CH₃CO- compounds react with alkaline aqueous iodine?











How do CH₃CO- compounds react with alkaline aqueous iodine?

lodine is added to the aldehyde or ketone solution. Just enough NaOH is added until the iodine is decolourised.

A pale yellow precipitate of tri-iodomethane, CH₃I, is formed.





