



A-level
CHEMISTRY
7405/3

Paper 3

Mark scheme

June 2023

Version: 0.3 Post-Standardisation



2 3 6 A 7 4 0 5 / 3 / M S

Mark schemes are prepared by the Lead Assessment Writer and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all associates participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every associate understands and applies it in the same correct way. As preparation for standardisation each associate analyses a number of students' scripts. Alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, associates encounter unusual answers which have not been raised they are required to refer these to the Lead Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this mark scheme are available from aqa.org.uk

Copyright information

AQA retains the copyright on all its publications. However, registered schools/colleges for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to schools/colleges to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Copyright © 2023 AQA and its licensors. All rights reserved.

AS and A-Level Chemistry

Mark Scheme Instructions for Examiners

1. General

The mark scheme for each question shows:

- the marks available for each part of the question
- the total marks available for the question
- the typical answer or answers which are expected
- extra information to help the examiner make his or her judgement and help to delineate what is acceptable or not worthy of credit or, in discursive answers, to give an overview of the area in which a mark or marks may be awarded.

The extra information in the 'Comments' column is aligned to the appropriate answer in the left-hand part of the mark scheme and should only be applied to that item in the mark scheme.

You should mark according to the contents of the mark scheme. If you are in any doubt about applying the mark scheme to a particular response, consult your Team Leader.

At the beginning of a part of a question a reminder may be given, for example: where consequential marking needs to be considered in a calculation; or the answer may be on the diagram or at a different place on the script.

In general the right-hand side of the mark scheme is there to provide those extra details which might confuse the main part of the mark scheme yet may be helpful in ensuring that marking is straightforward and consistent.

The use of M1, M2, M3 etc in the right-hand column refers to the marking points in the order in which they appear in the mark scheme. So, M1 refers to the first marking point, M2 the second marking point etc.

2. Emboldening

- 2.1** In a list of acceptable answers where more than one mark is available 'any **two** from' is used, with the number of marks emboldened. Each of the following bullet points is a potential mark.
- 2.2** A bold **and** is used to indicate that both parts of the answer are required to award the mark.
- 2.3** Alternative answers acceptable for a mark are indicated by the use of **OR**. Different terms in the mark scheme are shown by a / ; eg allow smooth / free movement.

3. Marking points

3.1 Marking of lists

This applies to questions requiring a set number of responses, but for which students have provided extra responses. The general 'List' principle to be followed in such a situation is that 'right + wrong = wrong'.

Each error / contradiction negates each correct response. So, if the number of error / contradictions equals or exceeds the number of marks available for the question, no marks can be awarded.

However, responses considered to be neutral (often prefaced by 'Ignore' in the mark scheme) are not penalised.

For example, in a question requiring 2 answers for 2 marks:

Correct answers	Incorrect answers (i.e. incorrect rather than neutral)	Mark (2)	Comment
1	0	1	
1	1	1	They have not exceeded the maximum number of responses so there is no penalty.
1	2	0	They have exceeded the maximum number of responses so the extra incorrect response cancels the correct one.
2	0	2	
2	1	1	
2	2	0	
3	0	2	The maximum mark is 2
3	1	1	The incorrect response cancels out one of the two correct responses that gained credit.
3	2	0	Two incorrect responses cancel out the two marks gained.
3	3	0	

3.2 Marking procedure for calculations

Full marks should be awarded for a correct numerical answer, without any working shown, unless the question states 'Show your working' or 'justify your answer'. In this case, the mark scheme will clearly indicate what is required to gain full credit.

If an answer to a calculation is incorrect and working is shown, process mark(s) can usually be gained by correct substitution / working and this is shown in the 'Comments' column or by each stage of a longer calculation.

3.3 Errors carried forward, consequential marking and arithmetic errors

Allowances for errors carried forward are most likely to be restricted to calculation questions and should be shown by the abbreviation ECF or consequential in the marking scheme.

An arithmetic error should be penalised for one mark only unless otherwise amplified in the marking scheme. Arithmetic errors may arise from a slip in a calculation or from an incorrect transfer of a numerical value from data given in a question.

3.4 Equations

In questions requiring students to write equations, state symbols are generally ignored unless otherwise stated in the 'Comments' column.

Examiners should also credit correct equations using multiples and fractions unless otherwise stated in the 'Comments' column.

3.5 Oxidation states

In general, the sign for an oxidation state will be assumed to be positive unless specifically shown to be negative.

3.6 Interpretation of 'it'

Answers using the word 'it' should be given credit only if it is clear that the 'it' refers to the correct subject.

3.7 Phonetic spelling

The phonetic spelling of correct scientific terminology should be credited **unless** there is a possible confusion with another technical term or if the question requires correct IUPAC nomenclature.

3.8 Brackets

(.....) are used to indicate information which is not essential for the mark to be awarded but is included to help the examiner identify the sense of the answer required.

3.9 Ignore / Insufficient / Do not allow

Ignore or insufficient is used when the information given is irrelevant to the question or not enough to gain the marking point. Any further correct amplification could gain the marking point.

Do **not** allow means that this is a wrong answer which, even if the correct answer is given, will still mean that the mark is not awarded.

3.10 Marking crossed out work

Crossed out work that **has not been** replaced should be marked as if it were not crossed out, if possible. Where crossed out work **has been** replaced, the replacement work and not the crossed out work should be marked.

3.11 Reagents

The command word "Identify", allows the student to choose to use **either** the name or the formula of a reagent in their answer. In some circumstances, the list principle may apply when both the name and the formula are used. Specific details will be given in mark schemes.

The guiding principle is that a reagent is a chemical which can be taken out of a bottle or container. Failure to identify complete reagents **will be penalised**, but follow-on marks (eg for a subsequent equation or observation) can be scored from an incorrect attempt (possibly an incomplete reagent) at the correct reagent. Specific details will be given in mark schemes.

For example, **no credit** would be given for

- the cyanide ion or CN^- when the reagent should be potassium cyanide or KCN;
- the hydroxide ion or OH^- when the reagent should be sodium hydroxide or NaOH;

- the $\text{Ag}(\text{NH}_3)_2^+$ ion when the reagent should be Tollens' reagent (or ammoniacal silver nitrate). In this example, no credit is given for the ion, but credit could be given for a correct observation following on from the use of the ion. Specific details will be given in mark schemes.

In the event that a student provides, for example, **both** KCN and cyanide ion, it would be usual to ignore the reference to the cyanide ion (because this is not contradictory) and credit the KCN. Specific details will be given in mark schemes.


3.12 Organic structures

Where students are asked to draw organic structures, unless a specific type is required in the question and stated in the mark scheme, these may be given as displayed, structural or skeletal formulas or a combination of all three as long as the result is unambiguous.

In general

- Displayed formulae must show all of the bonds and all of the atoms in the molecule, but need not show correct bond angles.
- Skeletal formulae must show carbon atoms by an angle or suitable intersection in the skeleton chain. Functional groups must be shown and it is essential that all atoms other than C atoms are shown in these (except H atoms in the functional groups of aldehydes, secondary amines and N-substituted amides which do not need to be shown).
- Structures must not be ambiguous, eg 1-bromopropane should be shown as $\text{CH}_3\text{CH}_2\text{CH}_2\text{Br}$ and not as the molecular formula $\text{C}_3\text{H}_7\text{Br}$ which could also represent the isomeric 2-bromopropane.
- Bonds should be drawn correctly between the relevant atoms. This principle applies in all cases where the attached functional group contains a carbon atom, eg nitrile, carboxylic acid, aldehyde and acid chloride. The carbon-carbon bond should be clearly shown. Wrongly bonded atoms will be penalised **on every occasion**. (see the examples below)
- The same principle should also be applied to the structure of alcohols. For example, if students show the alcohol functional group as C – HO, they should be penalised **on every occasion**.
- Latitude should be given to the representation of C – C bonds in alkyl groups, given that CH_3- is considered to be interchangeable with $\text{H}_3\text{C}-$ even though the latter would be preferred.
- Similar latitude should be given to the representation of amines where NH_2- C will be allowed, although $\text{H}_2\text{N}-$ C would be preferred.
- Poor presentation of vertical C – CH_3 bonds or vertical C – NH_2 bonds should **not** be penalised. For other functional groups, such as – OH and – CN, the limit of tolerance is the half-way position between the vertical bond and the relevant atoms in the attached group.

By way of illustration, the following would apply.



- Representation of CH_2 by C-H_2 will be penalised
- Some examples are given here of **structures** for specific compounds that should **not** gain credit (but, exceptions may be made in the context of balancing equations)

CH_3COH for ethanal

$\text{CH}_3\text{CH}_2\text{HO}$ for ethanol

OHCH_2CH_3 for ethanol

$\text{C}_2\text{H}_6\text{O}$ for ethanol

CH_2CH_2 for ethene

$\text{CH}_2\cdot\text{CH}_2$ for ethene

$\text{CH}_2:\text{CH}_2$ for ethene

- Each of the following **should gain credit** as alternatives to correct representations of the structures.

$\text{CH}_2 = \text{CH}_2$ for ethene, $\text{H}_2\text{C}=\text{CH}_2$

$\text{CH}_3\text{CHOHCH}_3$ for propan-2-ol, $\text{CH}_3\text{CH}(\text{OH})\text{CH}_3$

- In most cases, the use of “sticks” to represent C – H bonds in a structure should **not** be penalised. The exceptions to this when “sticks” will be penalised include
 - structures in mechanisms where the C – H bond is essential (e.g. elimination reactions in halogenoalkanes and alcohols)
 - when a displayed formula is required
 - when a skeletal structure is required or has been drawn by the candidate

3.13 Organic names

As a general principle, non-IUPAC names or incorrect spelling or incomplete names should **not** gain credit. Some illustrations are given here.


Unnecessary but not wrong numbers will **not** be penalised such as the number ‘2’ in 2-methylpropane or the number ‘1’ in 2-chlorobutan-1-oic acid.

but-2-ol	should be butan-2-ol
2-hydroxybutane	should be butan-2-ol
butane-2-ol	should be butan-2-ol
2-butanol	should be butan-2-ol
ethan-1,2-diol	should be ethane-1,2-diol
2-methpropan-2-ol	should be 2-methylpropan-2-ol
2-methylbutan-3-ol	should be 3-methylbutan-2-ol
3-methylpentan	should be 3-methylpentane
3-mythylpentane	should be 3-methylpentane
3-methypentane	should be 3-methylpentane
propanitrile	should be propanenitrile
aminethane	should be ethylamine (although aminoethane can gain credit)
2-methyl-3-bromobutane	should be 2-bromo-3-methylbutane
3-bromo-2-methylbutane	should be 2-bromo-3-methylbutane
3-methyl-2-bromobutane	should be 2-bromo-3-methylbutane
2-methylbut-3-ene	should be 3-methylbut-1-ene
difluorodichloromethane	should be dichlorodifluoromethane


3.14 Organic reaction mechanisms

Curly arrows should originate either from a lone pair of electrons or from a bond.

The following representations should not gain credit and will be penalised each time within a clip.



For example, the following would score zero marks



When the curly arrow is showing the formation of a bond to an atom, the arrow can go directly to the relevant atom, alongside the relevant atom or **more than half-way** towards the relevant atom.

In free-radical substitution

- the absence of a radical dot should be penalised **once only** within a clip.
- the use of half-headed arrows is not required, but the use of double-headed arrows or the incorrect use of half-headed arrows in free-radical mechanisms should be penalised **once only** within a clip

The correct use of skeletal formulae in mechanisms is acceptable, but where a C-H bond breaks, both the bond and the H must be drawn to gain credit.

3.15 Extended responses

For questions marked using a 'Levels of Response' mark scheme:

Level of response mark schemes are broken down into three levels, each of which has a descriptor. Each descriptor contains two statements. The first statement is the Chemistry content statement and the second statement is the communication statement.

Determining a level

Start at the lowest level of the mark scheme and use it as a ladder to see whether the answer meets the Chemistry content descriptor for that level. The descriptor for the level indicates the qualities that might be seen in the student's answer for that level. If it meets the lowest level, then go to the next one and decide if it meets this level, and so on, until you have a match between the level descriptor and the answer.

When assigning a level you should look at the overall quality of the answer and not look to pick holes in small and specific parts of the answer where the student has not performed quite as well as the rest. If the answer covers different aspects of different levels of the mark scheme you should use a best fit approach for defining the level.

Once the level has been decided, the mark within the level is determined by the communication statement:

- If the answer completely matches the communication descriptor, award the higher mark within the level.
- If the answer does not completely match the communication descriptor, award the lower mark within the level.

The exemplar materials used during standardisation will help you to determine the appropriate level. There will be an exemplar in the standardising materials which will correspond with each level of the mark scheme and for each mark within each level. This answer will have been awarded a mark by the Lead Examiner. You can compare the student's answer with the exemplar to determine if it is the same standard, better or worse than the example. You can then use this to allocate a mark for the answer based on the Lead Examiner's mark on the exemplar.

You may well need to read back through the answer as you apply the mark scheme to clarify points and assure yourself that the level and the mark are appropriate.

Indicative content in the mark scheme is provided as a guide for examiners. It is not intended to be exhaustive and you must credit other chemically valid points. Students may not have to cover all of the points mentioned in the indicative content to reach the highest level of the mark scheme. The mark scheme will state how much chemical content is required for the highest level.

An answer which contains nothing of relevance to the question must be awarded no marks.

For other extended response answers:

Where a mark scheme includes linkage words (such as 'therefore', 'so', 'because' etc), these are optional. However, a student's marks for the question may be limited if they do not demonstrate the ability to construct and develop a sustained line of reasoning which is coherent, relevant, substantiated and logically structured. In particular answers in the form of bullet pointed lists may not be awarded full marks if there is no indication of logical flow between each point or if points are in an illogical order. The mark schemes for some questions state that the maximum mark available for an extended response answer is limited if the answer is not coherent, relevant, substantiated and logically structured. During the standardisation process, the Lead Examiner will provide marked exemplar material to demonstrate answers which have not met these criteria. You should use these exemplars as a comparison when marking student answer.

MARK SCHEME – A-LEVEL CHEMISTRY – 7405/3 – JUNE 2023

Question	Answers	Additional comments/Guidelines	Mark
1.1	catalyst	ALLOW reduces E_a IGNORE speeds up reaction IGNORE provides alternative path IGNORE proton donor IGNORE dehydrating agent	1
Question	Answers	Additional comments/Guidelines	Mark
1.2	electric heater/heat mantle or (hot) water bath	IGNORE not with a Bunsen/naked flame / gently ALLOW hot water ALLOW heating/hot plate/ sand bath / oil bath ALLOW reference to flame/Bunsen if in context of heating a water bath NOT any indication of direct heat from Bunsen	1

MARK SCHEME – A-LEVEL CHEMISTRY – 7405/3 – JUNE 2023

Question	Answers	Additional comments/Guidelines	Mark
1.3	<p>M1 there is a bung/stopper (in the end of the condenser)</p> <p>M2 idea of pressure build up</p> <p>M3 water goes the wrong way through the condenser</p> <p>M4 water does not fill the condenser / condenser is not cool enough</p>	<p>owtte</p> <p>M2 stopper could be forced out</p> <p>IGNORE glass shatters / explodes</p> <p>water in at top / out at bottom</p> <p>IGNORE condenser is wrong way round</p> <p>ALLOW less condensing / vapour/gas/reactants/products will not condense / not as effective at cooling/condensing vapour/reactants/products escapes</p> <p>IGNORE uneven cooling</p> <p>NOT mixture in flask not cooled</p> <p>ALLOW M1/M3 neck of flask not sealed owtte</p> <p>M2/M4 vapour/reactants/products can escape</p> <p>IGNORE references to clamps</p>	4

MARK SCHEME – A-LEVEL CHEMISTRY – 7405/3 – JUNE 2023

Question	Answers	Additional comments/Guidelines	Mark
1.4	<p>M1 to neutralise/react with/remove the <u>acid</u></p> <p>M2 carbon dioxide / gas is produced</p>	<p>ALLOW carboxylic/ethanoic and/or sulfuric</p> <p>IGNORE react with/neutralise the distillate/mixture</p> <p>IGNORE to act as a base</p> <p>NOT if incorrect acid named</p> <p>ALLOW effervescence / bubbles / fizzes</p> <p>IGNORE water vapour produced</p> <p>NOT if incorrect gas named</p> <p>IGNORE pressure build up (as in Q)</p>	2

Question	Answers	Additional comments/Guidelines	Mark
1.5	<p>M1 ethyl ethanoate/it is immiscible with / insoluble in water</p> <p>M2 ethyl ethanoate/it is less dense / has <u>lower</u> density (than water)</p>	<p>ALLOW water/solution and ethyl ethanoate/it do not mix,</p> <p>OR aqueous and organic layers do not mix</p> <p>ALLOW ethyl ethanoate/it is hydrophobic</p> <p>IGNORE references to polarity / intermolecular forces</p> <p>IGNORE different/low density</p> <p>NOT lighter</p> <p>ALLOW answers for either mark from either part</p>	2

MARK SCHEME – A-LEVEL CHEMISTRY – 7405/3 – JUNE 2023

Question	Answers	Additional comments/Guidelines	Mark
1.6	to remove/absorb water / as a drying agent	ALLOW reacts with water ALLOW to dry the product/it IGNORE dehydrates NOT reference to crystals forming NOT to dry the reactants NOT to remove soluble impurities	1


MARK SCHEME – A-LEVEL CHEMISTRY – 7405/3 – JUNE 2023


Question	Answers	Additional comments/Guidelines	Mark
1.7	<p>M1 mass of ethanol = 10×0.790 (= 7.90 g)</p> <p>M2 amount of ethanol = $\frac{7.90}{46.0}$ (= 0.172 mol) AND amount of ethanoic acid = $\frac{5.25}{60.0}$ (= 0.0875 mol)</p> <p>M3 (limiting reagent is) ethanoic acid</p> <p>M4 (max amount of ethyl ethanoate = 0.0875 mol) max mass of ethyl ethanoate = 88.0×0.0875 (= 7.70 g)</p> <p>M5 % yield = $\frac{5.47}{M4} \times 100 = 71.0\%$ (70.6 to 71.1 to min 2sf)</p>	<p>Allow ECF at each stage</p> <p>M1 scores from 0.172 mol of ethanol</p> <p>M2 need to see numbers or sums for both substances</p> <p>M2 10/46 can only be ECF if 10 is identified as a mass</p> <p>M3 ECF from M2 if both amounts clearly shown and $n_{\text{ethanol}} < n_{\text{ethanoic acid}}$</p> <p>M4 independent of M3</p> <p>Alternative M4 & 5</p> <p>M4 Amount of ethyl ethanoate formed $= \frac{5.47}{88.0}$ (=0.0622)</p> <p>M5 % yield = $\frac{M4}{0.0875} \times 100 = 71.0\%$</p> <p>Correct answer scores M4 and M5 but mark M1/2/3 separately</p> <p>M5 must show an attempt at mass or moles of ester formed divided by mass or moles of ester expected</p>	5

MARK SCHEME – A-LEVEL CHEMISTRY – 7405/3 – JUNE 2023



Question	Answers	Additional comments/Guidelines	Mark
1.8	reaction is an equilibrium/reversible	ALLOW losses during distillation/isolation/purification/transfer / incomplete distillation / side reactions / by-products ALLOW incomplete reaction ALLOW impurities/contamination/water present / not dry IGNORE water is also produced (during the reaction)	1





Question	Answers	Additional comments/Guidelines	Mark
2.1	Displayed formula of pentan-1-ol $ \begin{array}{cccccc} & \text{H} & \text{H} & \text{H} & \text{H} & \text{H} \\ & & & & & \\ \text{H} & -\text{C} & -\text{C} & -\text{C} & -\text{C} & -\text{C}-\text{O}-\text{H} \\ & & & & & \\ & \text{H} & \text{H} & \text{H} & \text{H} & \text{H} \end{array} $	NOT pentan-3-ol NOT -OH	1


Question	Answers	Additional comments/Guidelines	Mark
2.2	Skeletal formula of 3-methylbutan-2-ol or 2-methylbutan-2-ol 	IGNORE numbers on C atoms IGNORE 'dots' at junctions IGNORE other non-skeletal structures IGNORE skeletal structure of pentan-2-ol NOT other incorrect skeletal structures NOT O-H NOT if bond clearly to H of OH	1

Question	Answers	Additional comments/Guidelines	Mark
2.3	one of these compounds 	Any structural representation of correct compound	1

MARK SCHEME – A-LEVEL CHEMISTRY – 7405/3 – JUNE 2023

Question	Answers	Additional comments/Guidelines	Mark
2.4	$ \begin{array}{c} \text{CH}_3 \\ \\ \text{CH}_3 - \text{C} - \text{CH}_2 - \text{OH} \\ \\ \text{CH}_3 \end{array} $	Any structural representation of correct compound	1
2.5	 <p>M1 loss of H₂O: arrow from C–O bond to O M2 loss of H⁺: arrow from correct C–H bond to correct C–C bond M3 elimination</p>	<p>M1/M2 list principle for additional arrows on any structure M1 NOT if arrow to + M3 IGNORE acid-catalysed / dehydration NOT nucleophilic / addition / electrophilic</p>	3
2.6		<p>Any structural representation of correct compound</p> <p>If skeletal CH₂ not needed</p> <p>Allow rings in place of C₆H₅</p>	1

Question	Answers	Additional comments/Guidelines	Mark
3.1	<p data-bbox="286 240 748 651">  </p> <p data-bbox="286 743 495 770">Alternative form</p> <p data-bbox="286 794 748 1209">  </p> <p data-bbox="286 1233 927 1318"> M1 Phe structure drawn with correct peptide link M2 amide group shown on end </p>	<p data-bbox="1193 240 1756 373"> M1 if Phe drawn with COOH or CONH₂ M2 ALLOW if no Phe drawn i.e. if NH₂ only attached directly to C=O on diagram </p> <p data-bbox="1193 397 1570 724">  </p> <p data-bbox="1193 748 1675 775">Scores M2 for ending in amide group</p> <p data-bbox="1193 799 1653 1158">  </p> <p data-bbox="1193 1182 1809 1315"> Scores M1 for Phe group M1 H needed on N of peptide link drawn unless C-CH-CH₂ drawn skeletal </p>	2

Question	Answers	Additional comments/Guidelines	Mark
3.2		<p>ALLOW zwitterion</p> <p>ALLOW -NH_2^+ and/or COO^-</p> <p>ALLOW with C shown in COOH group</p> <p>ALLOW without H on N</p> <p>ALLOW N–H</p> <p>NOT N–</p>	1

Question	Answers	Additional comments/Guidelines	Mark
3.3	<p>M1 (aqueous) HCl/hydrochloric acid</p> <p>M2 reflux/heat</p>	<p>Name or formula of any strong acid or alkali</p> <p>ALLOW warm / hot / high temperature for heat</p> <p>NOT $T > 200^\circ\text{C}$</p> <p>IGNORE conc as condition with acid/alkali</p> <p>IGNORE pressure</p> <p>Alternative</p> <p>M1 protease/(poly)peptidase/peptase/named protease</p> <p>IGNORE enzyme</p> <p>M2 warm</p> <p>NOT hot / high temperature / $T > 50^\circ\text{C}$</p>	2

MARK SCHEME – A-LEVEL CHEMISTRY – 7405/3 – JUNE 2023

Question	Answers	Additional comments/Guidelines	Mark
3.4	<p>M1 lid/cover (on beaker)</p> <p>Then any 2 from these 3</p> <ul style="list-style-type: none"> prevents escape of vapour (from beaker) / evaporation of solvent (from beaker) so atmosphere in beaker is saturated with solvent vapour owtte to reduce evaporation from the plate 	<p>ALLOW (for bullet point 3) so solvent can rise up plate</p> <p>ALLOW (for bullet point 3) to avoid plate drying out</p>	3

Question	Answers	Additional comments/Guidelines	Mark
3.5	Difference in the balance between solubility in solvent/mobile phase and attraction to/retention on stationary phase	<p>ALLOW difference between (relative) affinity/attraction for solvent and stationary phase</p> <p>ALLOW absorption/adsorption for retention on stationary phase</p>	1

Question	Answers	Additional comments/Guidelines	Mark
3.6	<p>M1 ninhydrin</p> <p>M2 amino acids are colourless / to make the amino acids visible</p>	<p>ALLOW iodine</p> <p>IGNORE UV</p> <p>IGNORE stated final colour e.g. “turns the amino acids purple” is not enough on its own</p> <p>IGNORE clear</p>	2

Question	Answers	Additional comments/Guidelines	Mark
3.7	0.54	ALLOW 0.53 – 0.55 (to min two sig figs)	1

Question	Answers	Additional comments/Guidelines	Mark								
4	<p>This question is marked using Levels of Response. Refer to the Mark Scheme Instructions for Examiners for guidance.</p> <table border="1" data-bbox="286 347 1164 1276"> <tr> <td data-bbox="286 347 448 563">Level 3 5–6 marks</td> <td data-bbox="448 347 1164 563"> <p>All stages are covered and the explanation of each stage is correct and virtually complete (ie two from stages 2 and 3 and three from stage 1).</p> <p>Answer communicates the whole explanation, including equations, coherently and shows a logical progression through all three stages.</p> </td> </tr> <tr> <td data-bbox="286 563 448 906">Level 2 3–4 marks</td> <td data-bbox="448 563 1164 906"> <p>All stages are covered (NB ‘covered’ means min two from stage 1) but the explanation of each stage may be incomplete or may contain inaccuracies.</p> <p>OR</p> <p>two stages covered and the explanations are generally correct and virtually complete (i.e. two from stages 2 and/or 3 and/or three from stage 1).</p> <p>Answer is coherent and shows some progression through all three stages. Some steps in each stage may be incomplete.</p> </td> </tr> <tr> <td data-bbox="286 906 448 1185">Level 1 1–2 marks</td> <td data-bbox="448 906 1164 1185"> <p>Two stages are covered (NB ‘covered’ means min two from stage 1) but the explanation of each stage may be incomplete or may contain inaccuracies.</p> <p>OR</p> <p>only one stage is covered but the explanation is generally correct and virtually complete (i.e. two from stages 2 and/or 3 and/or three from stage 1).</p> <p>Answer shows some progression between two stages.</p> </td> </tr> <tr> <td data-bbox="286 1185 448 1276">0 mark</td> <td data-bbox="448 1185 1164 1276">Insufficient correct chemistry to gain a mark.</td> </tr> </table>	Level 3 5–6 marks	<p>All stages are covered and the explanation of each stage is correct and virtually complete (ie two from stages 2 and 3 and three from stage 1).</p> <p>Answer communicates the whole explanation, including equations, coherently and shows a logical progression through all three stages.</p>	Level 2 3–4 marks	<p>All stages are covered (NB ‘covered’ means min two from stage 1) but the explanation of each stage may be incomplete or may contain inaccuracies.</p> <p>OR</p> <p>two stages covered and the explanations are generally correct and virtually complete (i.e. two from stages 2 and/or 3 and/or three from stage 1).</p> <p>Answer is coherent and shows some progression through all three stages. Some steps in each stage may be incomplete.</p>	Level 1 1–2 marks	<p>Two stages are covered (NB ‘covered’ means min two from stage 1) but the explanation of each stage may be incomplete or may contain inaccuracies.</p> <p>OR</p> <p>only one stage is covered but the explanation is generally correct and virtually complete (i.e. two from stages 2 and/or 3 and/or three from stage 1).</p> <p>Answer shows some progression between two stages.</p>	0 mark	Insufficient correct chemistry to gain a mark.	<p>Indicative Chemistry content</p> <p>Stage 1 IDs (Allow ppts if shown as products in equations or on the table) (Allow names – with oxidation states for Fe compounds)</p> <p>(1a) BOTH red-brown ppts from L = $\text{Fe}(\text{H}_2\text{O})_3(\text{OH})_3 / \text{Fe}(\text{OH})_3$</p> <p>(1b) green ppt from M = $\text{Fe}(\text{H}_2\text{O})_4(\text{OH})_2 / \text{Fe}(\text{OH})_2$ for NH_3 AND M = FeCO_3 with Na_2CO_3</p> <p>(1c) white ppt with L = AgCl AND white ppt with M = BaSO_4</p> <p>(1d) L = FeCl_3 AND M = FeSO_4</p> <p>Stage 2 Reactions with sodium carbonate (ALLOW if any 3+ compared with any 2+)</p> <p>(2a) carbon dioxide is gas produced from L (could come from equation)</p> <p>(2b) $\text{Fe}^{3+} / \text{Fe}(\text{H}_2\text{O})_6^{3+}$ (in L) is more acidic than $\text{Fe}^{2+} / \text{Fe}(\text{H}_2\text{O})_6^{2+}$ (in M) ora</p> <p>(2c) Fe^{3+} is smaller / has higher charge / greater charge density of Fe^{3+} / is more polarising (makes ion a better proton donor)</p>	6
Level 3 5–6 marks	<p>All stages are covered and the explanation of each stage is correct and virtually complete (ie two from stages 2 and 3 and three from stage 1).</p> <p>Answer communicates the whole explanation, including equations, coherently and shows a logical progression through all three stages.</p>										
Level 2 3–4 marks	<p>All stages are covered (NB ‘covered’ means min two from stage 1) but the explanation of each stage may be incomplete or may contain inaccuracies.</p> <p>OR</p> <p>two stages covered and the explanations are generally correct and virtually complete (i.e. two from stages 2 and/or 3 and/or three from stage 1).</p> <p>Answer is coherent and shows some progression through all three stages. Some steps in each stage may be incomplete.</p>										
Level 1 1–2 marks	<p>Two stages are covered (NB ‘covered’ means min two from stage 1) but the explanation of each stage may be incomplete or may contain inaccuracies.</p> <p>OR</p> <p>only one stage is covered but the explanation is generally correct and virtually complete (i.e. two from stages 2 and/or 3 and/or three from stage 1).</p> <p>Answer shows some progression between two stages.</p>										
0 mark	Insufficient correct chemistry to gain a mark.										

		<p>Stage 3 Equations (ECF from incorrect 3+ ion in L and/or 2+ ion in M)</p> <p>(3a) $\text{Fe}(\text{H}_2\text{O})_6^{3+} + 3\text{NH}_3 \rightarrow \text{Fe}(\text{H}_2\text{O})_3(\text{OH})_3 + 3\text{NH}_4^+$ AND $2\text{Fe}(\text{H}_2\text{O})_6^{3+} + 3\text{CO}_3^{2-} \rightarrow 2\text{Fe}(\text{H}_2\text{O})_3(\text{OH})_3 + 3\text{CO}_2 + 3\text{H}_2\text{O}$</p> <p>(3b) $\text{Fe}(\text{H}_2\text{O})_6^{2+} + 2\text{NH}_3 \rightarrow \text{Fe}(\text{H}_2\text{O})_4(\text{OH})_2 + 2\text{NH}_4^+$ AND $\text{Fe}(\text{H}_2\text{O})_6^{2+} + \text{CO}_3^{2-} \rightarrow \text{FeCO}_3 + 6\text{H}_2\text{O}$ OR $\text{Fe}^{2+} + \text{CO}_3^{2-} \rightarrow \text{FeCO}_3$</p> <p>(3c) $\text{Ag}^+ + \text{Cl}^- \rightarrow \text{AgCl}$ AND $\text{Ba}^{2+} + \text{SO}_4^{2-} \rightarrow \text{BaSO}_4$</p>	
--	--	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--

MARK SCHEME – A-LEVEL CHEMISTRY – 7405/3 – JUNE 2023

Question	Answers	Additional comments/Guidelines	Mark
5.1	<p>M1 energy transferred = 2.4×100 (= 240 kJ / 240000 J)</p> <p>M2 amount of water vaporised = $\frac{103}{18}$ (= 5.72 mol)</p> <p>M3 $\Delta H_{\text{vap}} = \frac{\text{M1 in kJ}}{\text{M2}} = (+)41.9 / 42.0$ (kJ mol⁻¹)</p>	<p>M1 IGNORE sign</p> <p>ALLOW ecf in M3 (if $q = mc\Delta T$ used for incorrect M1 then a value in kJ could score ecf in M3)</p> <p>ALLOW 41.9 to 42.11 to two or more sig figs</p> <p>M3 NOT if negative</p> <p>Correct answer = 3 marks</p>	3
5.2	<p>M1 O AND N more electronegative than C and/or H (so all have polar bonds)</p> <p>M2 CH₃CH₂OH and CH₃OCH₃ both v-shaped/non-linear/bent AND CH₃CH₂NH₂ (trigonal) pyramidal</p> <p>M3 shapes are not symmetrical (so molecules are polar)</p> <p>M4 O more electronegative than N (so ethylamine is least polar)</p>	<p>ALLOW 'different electronegativities' PLUS diagrams labelled $\delta+$ and $\delta-$</p> <p>ALLOW angular for v-shaped in M2</p> <p>Diagrams from M2 do not require lone pairs</p> <p>ALLOW M3 if diagrams in M2 show asymmetry</p> <p>Correct diagrams of the three shapes gives M2 and M3</p>	4

MARK SCHEME – A-LEVEL CHEMISTRY – 7405/3 – JUNE 2023




Question	Answers	Additional comments/Guidelines	Mark
5.3	<p>M1 hydrogen bonding in $\text{CH}_3\text{CH}_2\text{OH}$ and $\text{CH}_3\text{CH}_2\text{NH}_2$ AND (permanent) dipole-dipole forces in CH_3OCH_3</p> <p>M2 hydrogen bonding stronger (than other (intermolecular) forces)</p> <p>M3 hydrogen bonding stronger in $\text{CH}_3\text{CH}_2\text{OH}$ than in $\text{CH}_3\text{CH}_2\text{NH}_2$</p>	<p>IGNORE van der Waals' / temporary/induced dipole-dipole forces</p> <p>M1 NOT any reference to breaking covalent bonds</p> <p>M3 ALLOW reference to O being more/most electronegative (than N) OR ethanol has greater dipole moment / more polar than ethylamine</p> <p>If none of M1, M2 or M3 have been awarded: ALLOW one mark for an indication that higher boiling point = stronger intermolecular forces but NOT if reference to breaking covalent bonds</p>	3

MARK SCHEME – A-LEVEL CHEMISTRY – 7405/3 – JUNE 2023


Question	Answers	Additional comments/Guidelines	Mark
6.1	<p>M1 amount of HCl = 0.010 mol AND amount Ca(OH)_2 ($= \frac{0.6}{74.1}$) = 0.00810</p> <p>M2 0.00810 mol of Ca(OH)_2 requires 0.0162 mol HCl OR 0.01 mol of HCl requires 0.005 mol of Ca(OH)_2</p> <p>NB There must be an indication that 0.005 is the amount of Ca(OH)_2 needed and not just 0.01/2</p>	<p>Alternative:</p> <p>M1 amount of HCl = 0.010 mol amount of Ca(OH)_2 needed = 0.0050 mol</p> <p>NB There must be an indication that 0.005 is the amount of Ca(OH)_2 needed and not just 0.01/2</p> <p>M2 mass of Ca(OH)_2 needed = $0.0050 \times 74.1 = 0.3705$ g (so 0.6 g is excess)</p>	2

Question	Answers	Additional comments/Guidelines	Mark
6.2	<p>M1 0.400 g dm^{-3} means $0.400 \div 74.1$ ($= 0.00540 \text{ mol dm}^{-3}$)</p> <p>M2 $[\text{OH}^-] = \text{M1} \times 2$ ($= 0.0108 \text{ mol dm}^{-3}$)</p> <p>M3 $[\text{H}^+] = 6.80 \times 10^{-15} \div \text{M2}$ ($= 6.30 \times 10^{-13} \text{ mol dm}^{-3}$)</p> <p>M4 $\text{pH} = -\log[\text{H}^+]$</p> <p>M5 $= -\log \text{M3 with answer to 2dp}$ ($= 12.20$)</p>	<p>Correct answer to 2 dp = 5 marks ALLOW 12.21 for 5 marks</p> <p>M4 ALLOW if calculation shown containing a number that has been calculated as $[\text{H}^+]$</p>	5


MARK SCHEME – A-LEVEL CHEMISTRY – 7405/3 – JUNE 2023

Question	Marking Guidance	Mark	Comments					
07	C	1 (AO2)	<table border="1"> <tr> <td>C</td> <td>12</td> <td>13</td> <td>10</td> <td></td> </tr> </table>	C	12	13	10	
C	12	13	10					
08	B	1 (AO2)	166.0					
09	A	1 (AO1)	graphite					
10	C	1 (AO1)	$\text{Ba(s)} + \text{Cl}_2\text{(g)} \rightarrow \text{BaCl}_2\text{(s)}$					
11	D	1 (AO1)	total number of particles present					
12	C	1 (AO1)	The proportion of successful collisions increases because the catalysed reaction has a lower activation energy.					
13	D	1 (AO2)	6.87×10^{-4}					
14	B	1 (AO1)	magnesium in MgO					
15	B	1 (AO1)	Intermolecular forces are overcome.					
16	B	1 (AO1)	HCO_3^-					
17	D	1 (AO2)	adding 10 cm ³ of 1.0 mol dm ⁻³ HCl					
18	C	1 (AO2)	1.00×10^{-4}					
19	B	1 (AO2)	Sodium has the lowest melting point.					

MARK SCHEME – A-LEVEL CHEMISTRY – 7405/3 – JUNE 2023

20	A	1 (AO1)	ammonia and ammonium chloride
21	B	1 (AO1)	a colourless solution with effervescence
22	D	1 (AO1)	I ⁻
23	A	1 (AO1)	[CoCl ₄] ²⁻ is square planar.
24	C	1 (AO1)	FeSO ₄
25	C	1 (AO2)	They have no effect on the value of the equilibrium constant.
26	D	1 (AO3)	Step 5 H ₂ with a nickel catalyst
27	C	1 (AO1)	decomposition of ozone
28	A	1 (AO2)	0.133
29	B	1 (AO3)	
30	D	1 (AO1)	Ozone absorbs ultraviolet radiation.
31	A	1 (AO2)	propene
32	D	1 (AO1)	$\text{C}_6\text{H}_5\text{CH}(\text{OH})\text{CH}_3 + [\text{O}] \rightarrow \text{C}_6\text{H}_5\text{COCH}_3 + \text{H}_2\text{O}$
33	A	1 (AO1)	comparing fingerprint regions of their infrared spectra

MARK SCHEME – A-LEVEL CHEMISTRY – 7405/3 – JUNE 2023

34	A	1 (AO1)	
35	C	1 (AO1)	Enzymes work equally well on both optical isomers of a substrate.
36	D	1 (AO2)	replacement of two Cl ⁻ ligands