



Cambridge International AS & A Level

CHEMISTRY

9701/14

Paper 1 Multiple Choice

May/June 2025

1 hour 15 minutes

You must answer on the multiple choice answer sheet.

You will need: Multiple choice answer sheet
Soft clean eraser
Soft pencil (type B or HB is recommended)

INSTRUCTIONS

- There are **forty** questions on this paper. Answer **all** questions.
- For each question there are four possible answers **A, B, C** and **D**. Choose the **one** you consider correct and record your choice in soft pencil on the multiple choice answer sheet.
- Follow the instructions on the multiple choice answer sheet.
- Write in soft pencil.
- Write your name, centre number and candidate number on the multiple choice answer sheet in the spaces provided unless this has been done for you.
- Do **not** use correction fluid.
- Do **not** write on any bar codes.
- You may use a calculator.

INFORMATION


- The total mark for this paper is 40.
- Each correct answer will score one mark.
- Any rough working should be done on this question paper.
- The Periodic Table is printed in the question paper.
- Important values, constants and standards are printed in the question paper.

This document has **20** pages. Any blank pages are indicated.



2


- 1 The diagram shows the logarithm of the first 13 ionisation energies of an element.



Which statement is correct?

- A** A proton is lost for each successive ionisation energy.
- B** The element is aluminium.
- C** The element is silicon.
- D** The element has only one outer electron.
- 2 What is the empirical formula of butanoic acid?
- A** C_2H_4O **B** C_3H_6O **C** C_4H_8O **D** $C_5H_{10}O$
- 3 In which pair is the bond angle in the first species smaller than the smallest bond angle in the second species?
- A** CH_4 and SF_6
- B** CO_2 and BF_3
- C** H_2O and H_3O^+
- D** NH_4^+ and NH_3

- 4 Two glass vessels, M and N, are connected by a closed valve.



M contains helium at 20 °C at a pressure of 1×10^5 Pa. N has been evacuated, and has three times the volume of M.

The valve is opened and the temperature of the whole apparatus is raised to 100 °C.

What is the final pressure in the system?

- A 3.18×10^4 Pa
 B 4.24×10^4 Pa
 C 1.25×10^5 Pa
 D 5.09×10^5 Pa
- 5 Solid sulfur consists of molecules made up of eight atoms covalently bonded together.

The bonding in sulfur dioxide is O=S=O.

enthalpy change of combustion of S_8 , $\Delta H_c^\ominus S_8(s) = -2376 \text{ kJ mol}^{-1}$

energy required to break 1.0 mol $S_8(s)$ into gaseous atoms = 2232 kJ mol^{-1}

O=O bond enthalpy = 496 kJ mol^{-1}


Using these data, what is the value of the S=O bond enthalpy?

- A 239 kJ mol^{-1} B 257 kJ mol^{-1} C 319 kJ mol^{-1} D 536 kJ mol^{-1}
- 6 In this question, the average oxidation state of sulfur in $\underline{S}_2O_3^{2-}$ and sulfur in $\underline{S}_2O_4^{2-}$ should be used.

In which reaction does the underlined element have the largest increase in oxidation state?

- A $3\underline{Cr}O_4^{3-}(aq) + 8H^+(aq) \rightarrow 2CrO_4^{2-}(aq) + Cr^{3+}(aq) + 4H_2O(l)$
 B $2\underline{N}O_2(g) + H_2O(l) \rightarrow HNO_3(aq) + HNO_2(aq)$
 C $\underline{S}_2O_3^{2-}(aq) + 2H^+(aq) \rightarrow S(s) + SO_2(g) + H_2O(l)$
 D $2\underline{S}_2O_4^{2-}(aq) + H_2O(l) \rightarrow S_2O_3^{2-}(aq) + 2HSO_3^-(aq)$

- 7 Methanol, CH_3OH , is made industrially from carbon monoxide and hydrogen in the equilibrium reaction shown.







Which statement about this equilibrium is correct?

- A K_p for the process is $\frac{p_{\text{CH}_3\text{OH}}}{p_{\text{CO}} \times p_{\text{H}_2}}$.
- B An increase in pressure increases the equilibrium yield of methanol.
- C An increase in temperature increases the equilibrium yield of methanol.
- D The addition of an effective catalyst increases the equilibrium yield of methanol.
- 8 The distribution of molecular energies in an ideal gas can be represented in a Boltzmann distribution.

Which change in conditions leads to a **larger** value for the number of molecules that have the most probable energy?

- A keeping the temperature constant but decreasing the pressure
- B keeping the pressure constant but decreasing the temperature
- C keeping the temperature constant but increasing the pressure
- D keeping the pressure constant but increasing the temperature

- 9 Which graph represents the number of unpaired electrons in the atoms of six elements in Period 3 of the Periodic Table?



- 10 A 69.0 g sample of nitrogen dioxide is placed in a reaction vessel.

The initial pressure of the nitrogen dioxide is P . An effective catalyst is then added and the nitrogen dioxide begins to decompose into its elements.




After ten minutes, the total pressure is $1.1P$.

What is the mass of oxygen molecules in the reaction vessel after ten minutes?

- A** 4.80 g **B** 9.60 g **C** 38.4 g **D** 48.0 g
- 11 Which statement about the molecule PF_5 is correct?
- A** Every F-P-F bond angle is 90° .
- B** The central atom in the molecule does **not** have a lone pair of electrons.
- C** The molecule has an overall dipole moment.
- D** The shape of the molecule is octahedral.
- 12 Which solid compound has both ionic and covalent bonding but **not** coordinate bonding?

- A** Al_2Cl_6 **B** CH_3COONa **C** MgCl_2 **D** NH_4Cl

13 Which equation represents the standard enthalpy change of formation, $\Delta H_f^\ominus$, for ethanol?



14 When K_2MnO_4 reacts with concentrated hydrochloric acid, the products include chlorine molecules and MnCl_2 . All of the manganese atoms are reduced to MnCl_2 .

Both Mn and Cl change their oxidation numbers during the reaction. No other element is oxidised or reduced.

Using these changes in oxidation number, how many moles of chlorine will be produced when 1.0 mol of K_2MnO_4 reacts with an excess of hydrochloric acid?


A 2.0 mol

B 2.5 mol

C 3.0 mol

D 4.0 mol

15 A nitrogen–hydrogen mixture, initially in the mole ratio of 1 : 3, reaches equilibrium with ammonia when 50% of the nitrogen has reacted. The total final pressure is p .



What is the partial pressure of ammonia in the equilibrium mixture?

A $\frac{p}{6}$

B $\frac{p}{4}$

C $\frac{p}{3}$

D $\frac{p}{2}$

- 16** Propyl methanoate is hydrolysed with NaOH(aq) at 20 °C to form two products, X and Y. Product X is an alcohol.

Data from the experiment is shown.

time / s	[X] / mol dm ⁻³
0	0.000
40	0.004
80	0.007
120	0.010
180	0.015
240	0.019
300	0.022

Which row is correct?

	average rate of reaction between 240 and 300 s	product Y
A	$5.00 \times 10^{-5} \text{ mol dm}^{-3} \text{ s}^{-1}$	HCOONa
B	$5.00 \times 10^{-5} \text{ mol dm}^{-3} \text{ s}^{-1}$	HCOOH
C	$7.33 \times 10^{-5} \text{ mol dm}^{-3} \text{ s}^{-1}$	HCOONa
D	$7.33 \times 10^{-5} \text{ mol dm}^{-3} \text{ s}^{-1}$	HCOOH

17 The table shows the numbers of bond pairs and lone pairs in four different species.

Which row is correct?

	species	total number of bond pairs	total number of lone pairs
A	nitrogen molecule	3	1
B	ammonia molecule	4	1
C	ammonium ion	4	0
D	hydroxide ion	1	4

18 Which row is correct?

	property	explanation
A	AgI dissolves in aqueous ammonia more readily than AgCl does	AgI reacts with aqueous ammonia
B	HCl decomposes more readily than HI does	Cl is more electronegative than I
C	I ₂ has a higher melting point than Cl ₂	I ₂ has stronger van der Waals' forces than Cl ₂
D	I ₂ is a stronger oxidising agent than Cl ₂	an I atom loses electrons more readily than a Cl atom

19 Substance J reacts with water. A gas is given off and the pH of the solution increases. The solution is then reacted with sulfuric acid and a white precipitate forms.

What could be substance J?

- A** barium
- B** barium oxide
- C** magnesium
- D** magnesium oxide

- 20** Compound L has empirical formula NH. It decomposes on gentle heating to produce ammonia and compound M only. An aqueous solution of compound L is a good conductor of electricity.

Which row could be correct?


	identity of M	species present in L(aq)
A	N_2H_4	N^{3-} and H^+
B	N_2H_4	NH_4^+ and NH_2^-
C	HN_3	NH_3 and HN_3
D	HN_3	NH_4^+ and N_3^-

- 21** X, Y and Z are elements in Period 3 of the Periodic Table. The results of some experiments carried out with compounds of these elements are shown.

element	result of adding the oxide of the element to $\text{H}_2\text{O}(\text{l})$	result of adding the chloride of the element to $\text{H}_2\text{O}(\text{l})$	result of adding the oxide of the element to $\text{HCl}(\text{aq})$
X	no reaction	hydrolyses	forms chloride salt
Y	forms hydroxide	dissolves	forms chloride salt
Z	forms acid	hydrolyses	hydrolyses

Which statement is correct?

- A** Element X is Al and element Y is Mg.
B Element X is Si and element Y is Na.
C Element Y is Al and element Z is P.
D Element Y is Na and element Z is Al.
- 22** The flow diagram shows two successive reactions starting from element Q. Element Q is either calcium or barium.



Element Q forms a nitrate that is less thermally stable than strontium nitrate.

What is the identity of compound R?

- A** CaO **B** $\text{Ca}(\text{OH})_2$ **C** BaO **D** $\text{Ba}(\text{OH})_2$


- 23** Which anions are formed when chlorine is passed into cold aqueous potassium hydroxide?
- A** Cl^- and ClO^-
 - B** Cl^- and ClO_3^-
 - C** Cl^- and ClO_4^-
 - D** ClO^- and ClO_3^-
- 24** What increases for each successive element in Period 3 from sodium to sulfur?
- A** the highest oxidation number of the element seen in an oxide
 - B** the melting point of the elements
 - C** the number of occupied orbitals in the atom
 - D** the pH of the solutions of the chlorides in water
- 25** Which statement about an ammonium ion is correct?
- A** All of the H–N–H bond angles in the ion are 90° .
 - B** All of the H–N–H bond angles in the ion are 107° .
 - C** The ion contains a N–H dative covalent bond which is weaker than the other three N–H covalent bonds.
 - D** The ion will react with a base as it is a weak acid.
- 26** An alcohol, U, is reacted with hot acidified $\text{K}_2\text{Cr}_2\text{O}_7$ solution. The organic product of the reaction contains 58.8% C, 9.8% H and 31.4% O by mass.
- What is the identity of alcohol U?
- A** 2-methylbutan-2-ol
 - B** 3-methylbutan-2-ol
 - C** pentan-1-ol
 - D** propan-1-ol

27 Which row shows a primary, a secondary and a tertiary alcohol?

	primary	secondary	tertiary
A	$\begin{array}{c} \text{CH}_2\text{OH} \\ \\ \text{CH}_2 \\ \\ \text{CH}_3 \end{array}$	$\begin{array}{c} \text{CH}_2\text{OH} \\ \\ \text{CHOH} \\ \\ \text{CH}_3 \end{array}$	$\begin{array}{c} \text{CH}_2\text{OH} \\ \\ \text{CHOH} \\ \\ \text{CH}_2\text{OH} \end{array}$
B	$\begin{array}{c} \text{CH}_2\text{OH} \\ \\ \text{CH}_3-\text{C}-\text{H} \\ \\ \text{CH}_3 \end{array}$	$\begin{array}{c} \text{CH}_3 \\ \\ \text{CH}_3-\text{C}-\text{OH} \\ \\ \text{CH}_3 \end{array}$	$\begin{array}{c} \text{CH}_3 \\ \\ \text{CH}_3-\text{C}-\text{H} \\ \\ \text{CH}_2\text{OH} \end{array}$
C	$\begin{array}{c} \text{CH}_2\text{OH} \\ \\ \text{CH}_3-\text{C}-\text{H} \\ \\ \text{H} \end{array}$	$\begin{array}{c} \text{CH}_2\text{OH} \\ \\ \text{CH}_3-\text{C}-\text{CH}_2\text{OH} \\ \\ \text{H} \end{array}$	$\begin{array}{c} \text{CH}_2\text{OH} \\ \\ \text{CH}_3-\text{C}-\text{CH}_2\text{OH} \\ \\ \text{CH}_2\text{OH} \end{array}$
D	$\begin{array}{c} \text{H} \\ \\ \text{CH}_3-\text{C}-\text{OH} \\ \\ \text{H} \end{array}$	$\begin{array}{c} \text{CH}_3 \\ \\ \text{CH}_3-\text{C}-\text{OH} \\ \\ \text{H} \end{array}$	$\begin{array}{c} \text{CH}_3 \\ \\ \text{CH}_3-\text{C}-\text{OH} \\ \\ \text{CH}_3 \end{array}$


28 1-chloro-2-methylbutane reacts with sodium cyanide in a nucleophilic substitution reaction.

What is the most likely intermediate or transition state in this reaction?




- 29** When 0.010 mol of a hydrocarbon X reacts with 720 cm³ of hydrogen at room conditions, an alkane is formed.


What is hydrocarbon X?



- 30** An alkene reacts with hot concentrated acidified KMnO₄ to produce a single organic product as shown.




What is the structure of the alkene?



- 31 Testosterone is an optically active organic molecule.


testosterone



How many chiral centres are there in one molecule of testosterone?

- A** 5 **B** 6 **C** 7 **D** 8
- 32 Quinone is an unsaturated molecule.

quinone



Which statement about quinone is correct?

- A** Quinone is non-planar and has an overall dipole moment.
B Quinone is non-planar and does **not** have an overall dipole moment.
C Quinone is planar and has an overall dipole moment.
D Quinone is planar and does **not** have an overall dipole moment.
- 33 Which reagent would react with 1-bromopropane to give the highest yield of propene?
- A** ammonia
B aqueous potassium hydroxide
C potassium cyanide
D ethanolic sodium hydroxide
- 34 Acidified potassium dichromate(VI), $K_2Cr_2O_7$, is added to propan-1-ol and the mixture is immediately distilled. The distillate is treated with HCN in the presence of KCN.

What is the organic product?

- A** $CH_3C(CN)(OH)CH_3$
B $CH_3CH_2CH_2CO_2H$
C $CH_3CH_2CH(OH)CN$
D $CH_3CH_2CH_2CN$

- 35** Compound Q gives positive results when tested separately with alkaline $I_2(aq)$ and with Tollens' reagent.

What is compound Q?

- A** $CHOCH_2CH_2CHO$
B CH_3CH_2COCHO
C CH_3COCH_2CHO
D $CH_3COCOCH_3$

- 36** Structural isomerism and stereoisomerism should be considered when answering this question.

How many isomeric esters of methanoic acid can be made with the molecular formula $C_5H_{10}O_2$?

- A** 2 **B** 3 **C** 4 **D** 5

- 37** The juice of one lemon reacts completely with 120 cm^3 of 0.50 mol dm^{-3} sodium carbonate solution.

The formula of citric acid is $HOOCCH_2C(OH)(COOH)CH_2COOH$.

No sodium carbonate is left unreacted.

What is the amount of citric acid in one lemon assuming that it is the only acid in the sample?

- A** 0.02 mol **B** 0.04 mol **C** 0.06 mol **D** 0.09 mol

- 38** Separate samples of 1-bromopropane are used in two different reactions.


reaction 1 1-bromopropane is converted into compound X by heating it under pressure with ammonia in ethanol.

reaction 2 1-bromopropane is converted into compound Y. Compound Y undergoes hydrolysis to form butanoic acid.


Which row identifies compound X and describes the reagents used in reaction 2 to make compound Y?

	identity of compound X	reagents for reaction 2
A	$CH_3CH_2CH_2NH_2$	$HCN(aq)$
B	$CH_3CH_2CH_2NH_2$	KCN dissolved in ethanol
C	$CH_3CH_2CH_2OH$	$HCN(aq)$
D	$CH_3CH_2CH_2OH$	KCN dissolved in ethanol


39 The diagram shows part of a polymer chain.



Which monomer would form this polymer?



40 The mass spectrum of element Z is shown.



What is element Z?

- A** arsenic
- B** chlorine
- C** gallium
- D** tungsten

BLANK PAGE

BLANK PAGE

BLANK PAGE

Important values, constants and standards

molar gas constant	$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
Faraday constant	$F = 9.65 \times 10^4 \text{ C mol}^{-1}$
Avogadro constant	$L = 6.02 \times 10^{23} \text{ mol}^{-1}$
electronic charge	$e = -1.60 \times 10^{-19} \text{ C}$
molar volume of gas	$V_m = 22.4 \text{ dm}^3 \text{ mol}^{-1}$ at s.t.p. (101 kPa and 273 K) $V_m = 24.0 \text{ dm}^3 \text{ mol}^{-1}$ at room conditions
ionic product of water	$K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ (at 298 K (25 °C))
specific heat capacity of water	$c = 4.18 \text{ kJ kg}^{-1} \text{ K}^{-1}$ ($4.18 \text{ J g}^{-1} \text{ K}^{-1}$)

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

The Periodic Table of Elements

Group																					
1	2											13	14	15	16	17	18				
		Key atomic number atomic symbol name relative atomic mass												1 Hhydrogen1.0							
3 Li lithium 6.9	4 Be beryllium 9.0	21 Sc scandium 45.0	22 Ti titanium 47.9	23 V vanadium 50.9	24 Cr chromium 52.0	25 Mn manganese 54.9	26 Fe iron 55.8	27 Co cobalt 58.9	28 Ni nickel 58.7	29 Cu copper 63.5	30 Zn zinc 65.4	31 Ga gallium 69.7	32 Ge germanium 72.6	33 As arsenic 74.9	34 Se selenium 79.0	35 Br bromine 79.9	36 Kr krypton 83.8				
11 Na sodium 23.0	12 Mg magnesium 24.3	39 Y yttrium 88.9	40 Zr zirconium 91.2	41 Nb niobium 92.9	42 Mo molybdenum 95.9	43 Tc technetium	44 Ru ruthenium 101.1	45 Rh rhodium 102.9	46 Pd palladium 106.4	47 Ag silver 107.9	48 Cd cadmium 112.4	49 In indium 114.8	50 Sn tin 118.7	51 Sb antimony 121.8	52 Te tellurium 127.6	53 I iodine 126.9	54 Xe xenon 131.3				
55 Cs caesium 132.9	56 Ba barium 137.3	57–71 lanthanoids	72 Hf hafnium 178.5	73 Ta tantalum 180.9	74 W tungsten 183.8	75 Re rhenium 186.2	76 Os osmium 190.2	77 Ir iridium 192.2	78 Pt platinum 195.1	79 Au gold 197.0	80 Hg mercury 200.6	81 Tl thallium 204.4	82 Pb lead 207.2	83 Bi bismuth 209.0	84 Po polonium	85 At astatine	86 Rn radon				
87 Fr francium	88 Ra radium	89–103 actinoids	104 Rf rutherfordium	105 Db dubnium	106 Sg seaborgium	107 Bh bohrium	108 Hs hassium	109 Mt meitnerium	110 Ds darmstadtium	111 Rg roentgenium	112 Cn copernicium	113 Nh nihonium	114 Fl flerovium	115 Mc moscovium	116 Lv livermorium	117 Ts tennessine	118 Og oganesson				
lanthanoids		57 La lanthanum 138.9	58 Ce cerium 140.1	59 Pr praseodymium 140.9	60 Nd neodymium 144.2	61 Pm promethium	62 Sm samarium 150.4	63 Eu europium 152.0	64 Gd gadolinium 157.3	65 Tb terbium 158.9	66 Dy dysprosium 162.5	67 Ho holmium 164.9	68 Er erbium 167.3	69 Tm thulium 168.9	70 Yb ytterbium 173.1	71 Lu lutetium 175.0					
actinoids		89 Ac actinium	90 Th thorium 232.0	91 Pa protactinium 231.0	92 U uranium 238.0	93 Np neptunium	94 Pu plutonium	95 Am americium	96 Cm curium	97 Bk berkelium	98 Cf californium	99 Es einsteinium	100 Fm fermium	101 Md mendelevium	102 No nobelium	103 Lr lawrencium					

lanthanoids

actinoids