

**UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS**  
GCE Advanced Subsidiary Level and GCE Advanced Level

**MARK SCHEME for the May/June 2012 question paper**  
**for the guidance of teachers**

**9701 CHEMISTRY**

**9701/23**

Paper 2 (AS Structured Questions), maximum raw mark 60

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

- Cambridge will not enter into discussions or correspondence in connection with these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2012 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.



|        |                                |          |       |
|--------|--------------------------------|----------|-------|
| Page 2 | Mark Scheme: Teachers' version | Syllabus | Paper |
|        | GCE AS/A LEVEL – May/June 2012 | 9701     | 23    |

- 1 (a) (i) from Na to Cl
- nuclear charge increases (1)  
 electrons are in the same shell/have the same shielding (1)  
 nuclear attraction increases (1)
- (ii) argon does not form any bonds/compounds **or**  
 argon exists as single atoms/is monatomic (1) [4]
- (b) (i)
- | radius of cation/nm |                  |                  | radius of anion/nm |                 |                 |
|---------------------|------------------|------------------|--------------------|-----------------|-----------------|
| Na <sup>+</sup>     | Mg <sup>2+</sup> | Al <sup>3+</sup> | P <sup>3-</sup>    | S <sup>2-</sup> | Cl <sup>-</sup> |
| 0.095               | 0.065            | 0.050            | 0.212              | 0.184           | 0.181           |
- (1)
- (ii) cations contain fewer electrons than the corresponding atoms **or**  
 cations contain fewer electrons than they do protons (1)  
 nucleus has a greater attraction (1)
- (iii) anions contain more electrons than the corresponding atoms **or**  
 anions contain more electrons than they do protons (1)  
 nucleus has a smaller attraction (1) [5]
- (c) (i) Na<sub>2</sub>O + H<sub>2</sub>O → 2NaOH (1)  
 SO<sub>2</sub> + H<sub>2</sub>O → H<sub>2</sub>SO<sub>3</sub> (1)
- (ii) for Na<sub>2</sub>O 10 to 14 (1)  
 for SO<sub>2</sub> 1 to 4 (1)
- (iii) NaOH + H<sub>2</sub>SO<sub>3</sub> → NaHSO<sub>3</sub> + H<sub>2</sub>O **or**  
 2NaOH + H<sub>2</sub>SO<sub>3</sub> → Na<sub>2</sub>SO<sub>3</sub> + 2H<sub>2</sub>O (1) [5]

**[Total: 14]**

| Page 3 | Mark Scheme: Teachers' version | Syllabus | Paper |
|--------|--------------------------------|----------|-------|
|        | GCE AS/A LEVEL – May/June 2012 | 9701     | 23    |

- 2 (a) (i)  $\text{Na}_2\text{CO}_3 + 2\text{HCl} \rightarrow 2\text{NaCl} + \text{H}_2\text{O} + \text{CO}_2$  (1)
- (ii)  $n(\text{HCl}) = \frac{35.8}{1000} \times 0.100 = 3.58 \times 10^{-3}$  (1)
- (iii)  $n(\text{Na}_2\text{CO}_3) = \frac{35.8}{2} \times 10^{-3} = 1.79 \times 10^{-3}$  mol in  $25.0 \text{ cm}^3$  (1)
- (iv)  $n(\text{Na}_2\text{CO}_3) = 1.79 \times 10^{-3} \times 10 = 1.79 \times 10^{-2}$  mol in  $250 \text{ cm}^3$  (1)
- (v) mass of  $\text{Na}_2\text{CO}_3 = 1.79 \times 10^{-2} \times 106 = 1.90\text{g}$   
 $M_r$  of  $\text{Na}_2\text{CO}_3 = 106$  (1)  
 mass of  $\text{Na}_2\text{CO}_3 = 1.90 \text{ g}$  (1) [6]
- (b)  $n(\text{H}_2\text{O})$  in  $5.13 \text{ g}$  of washing soda =  $\frac{5.13 - 1.90}{18} = 1.79 \times 10^{-1}$  mol (1)  
 $n(\text{Na}_2\text{CO}_3)$  in  $5.13 \text{ g}$  of washing soda =  $1.79 \times 10^{-2}$  mol  
 $n(\text{H}_2\text{O}) : n(\text{Na}_2\text{CO}_3) = 10 : 1$  (1)
- or  
 $1.90 \text{ g Na}_2\text{CO}_3$  are combined with  $3.23 \text{ g H}_2\text{O}$   
 $106 \text{ g Na}_2\text{CO}_3$  are combined with  $\frac{3.23 \times 106}{1.90} = 180.2 \text{ g H}_2\text{O}$  (1)  
 this is  $10 \text{ mol}$  of  $\text{H}_2\text{O}$  (1)
- or  
 $1.79 \times 10^{-2} \text{ mol Na}_2\text{CO}_3 \cdot x\text{H}_2\text{O} \equiv 5.13 \text{ g}$  of washing soda  
 $1 \text{ mol Na}_2\text{CO}_3 \cdot x\text{H}_2\text{O} \equiv \frac{5.13}{1.79 \times 10^{-2}} = 286.6 \text{ g}$  (1)  
 $\text{Na}_2\text{CO}_3 = 106$  and  $\text{H}_2\text{O} = 18$  hence  $x = 10$  (1) [2]

[Total: 8]

| Page 4 | Mark Scheme: Teachers' version | Syllabus | Paper |
|--------|--------------------------------|----------|-------|
|        | GCE AS/A LEVEL – May/June 2012 | 9701     | 23    |

- 3 (a)  $\text{CH}_3\text{OCH}_3(\text{l}) + 3\text{O}_2(\text{g}) \rightarrow 2\text{CO}_2(\text{g}) + 3\text{H}_2\text{O}(\text{l})$  (1)  
 the enthalpy change/heat change/heat evolved when  
 one mole of  $\text{CH}_3\text{OCH}_3$ /a compound (1)  
 is completely burned **or**  
 burned in an excess of air/oxygen (1) [3]


- (b) 
$$2\text{CH}_3\text{OH}(\text{l}) \rightarrow \text{CH}_3\text{OCH}_3(\text{g}) + \text{H}_2\text{O}(\text{l})$$

$$\Delta H_f^\ominus/\text{kJ mol}^{-1} \quad 2(-239) \quad -184 \quad -286$$

$$\Delta H_{\text{reaction}}^\ominus = -184 + (-286) - 2(-239)$$

$$= +8 \text{ kJ mol}^{-1}$$
 (1)  
 (1)  
 correct sign (1) [3]

- (c) (i)



**both** correct (1)


- (ii) structural isomerism **or** functional group isomerism (1) [2]

- (d) (i) hydrogen bonds (1)

- (ii) lone pair on O atom of  $\text{C}_2\text{H}_5\text{OH}$  (1)

correct dipole  $\text{O}^{\delta-}-\text{H}^{\delta+}$  on bond in one molecule of ethanol (1)

hydrogen bond shown between lone pair of an O atom and a hydrogen atom,  
 i.e.



(1) [4]


**[Total: 12]**

| Page 5 | Mark Scheme: Teachers' version | Syllabus | Paper |
|--------|--------------------------------|----------|-------|
|        | GCE AS/A LEVEL – May/June 2012 | 9701     | 23    |

- 4 (a) high temperature and high pressure (1)  
 high temperature and catalyst (1) [2]


- (b)  $C_{12}H_{26} \rightarrow C_4H_8 + C_8H_{18}$  or (1)  
 $C_{12}H_{26} \rightarrow 2C_4H_8 + C_4H_{10}$  (1) [1]

(c)



(4 × 1) [4]

(d) (i)




(1)

- (ii) compound B (1)  
 compound C (1) [3]

|        |                                |          |       |
|--------|--------------------------------|----------|-------|
| Page 6 | Mark Scheme: Teachers' version | Syllabus | Paper |
|        | GCE AS/A LEVEL – May/June 2012 | 9701     | 23    |

(e)

allow any orientation of CH<sub>3</sub>- groups

(1) [1]

(f) (i) CH<sub>2</sub>=CH—CH=CH<sub>2</sub>allow CH<sub>3</sub>CHOHCH=CH<sub>2</sub> and CH<sub>3</sub>C≡CCH<sub>3</sub>

(1)

(ii) CH<sub>2</sub>BrCHBrCHBrCH<sub>2</sub>Brallow CH<sub>3</sub>CBr<sub>2</sub>CBr<sub>2</sub>CH<sub>3</sub> from CH<sub>3</sub>CHOHCH=CH<sub>2</sub>allow CH<sub>3</sub>CHOHCHBrCH<sub>2</sub>Br from CH<sub>3</sub>C≡CCH<sub>3</sub>

(1)

(iii) electrophilic addition

**both** words required

(1) [3]

**[Total: 14]**

|        |                                |          |       |
|--------|--------------------------------|----------|-------|
| Page 7 | Mark Scheme: Teachers' version | Syllabus | Paper |
|        | GCE AS/A LEVEL – May/June 2012 | 9701     | 23    |

- 5 (a) (i) CO<sub>2</sub>/carbon dioxide (1)
- (ii) carboxylic acid **or** –CO<sub>2</sub>H **or** –COOH (1) [2]
- (b) (i) dehydration **or** elimination (1)
- (ii) H contains >C=C< bond (1)  
 H contains –CO<sub>2</sub>H group (1)  
 H is CH<sub>2</sub>=CHCO<sub>2</sub>H (1) [4]
- (c)  $n(\text{F}) = \frac{0.600}{90} = 6.67 \times 10^{-3} \text{ mol}$  (1)
- F contains one –OH group and one –CO<sub>2</sub>H group (1)  
 hence one mole of F produces one mole of H<sub>2</sub> with Na (1)  
 $n(\text{H}_2) = 6.67 \times 10^{-3} \text{ mol}$  (1)  
 vol. of H<sub>2</sub> =  $6.67 \times 10^{-3} \times 24000 \text{ cm}^3$   
 =  $160 \text{ cm}^3$  at room temperature and pressure (1) [4]

(d) (i)

|   |   |
|---|---|
| HOCH <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> H | CH <sub>3</sub> CH(OH)CO <sub>2</sub> H |
| <b>J</b>  | <b>K</b>                                |

one isomer correct

(1)

(ii)

|  |                                     |
|--|-------------------------------------|
| HO <sub>2</sub> CCH <sub>2</sub> CO <sub>2</sub> H | CH <sub>3</sub> COCO <sub>2</sub> H |
| <b>product from J</b>                              | <b>product from K</b>               |

**one** oxidation product correct

(1) [2]

**[Total: 12]**