

CAMBRIDGE INTERNATIONAL EXAMINATIONS
GCE Advanced Subsidiary Level and GCE Advanced Level

MARK SCHEME for the October/November 2012 series

9701 CHEMISTRY

9701/22

Paper 2 (AS Structured Questions), maximum raw mark 60

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2012 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.

Page 2	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – October/November 2012	9701	22

- 1 (a) ZnCO_3 Zn(OH)_2 ZnO
not Zn or other compounds of Zn (any 2) [2]
- (b) (i) to ensure all of the water of crystallisation had been driven off **or**
to be at constant mass (1)
- (ii) mass of $\text{ZnSO}_4 = 76.34 - 74.25 = 2.09 \text{ g}$ (1)
- $M_r \text{ ZnSO}_4 = 65.4 + 32.1 + (4 \times 16.0) = 161.5$
- allow use of $\text{Zn} = 65$ and/or $\text{S} = 32$ to give values between 161 and 161.5 (1)
- $n(\text{ZnSO}_4) = \frac{2.09}{161.5} = 0.01294 = 1.29 \times 10^{-2}$
- $\text{ZnSO}_4 = 161$ gives 1.30×10^{-2} (1)
- (iii) mass of H_2O driven off = $77.97 - 76.34 = 1.63 \text{ g}$ (1)
- $n(\text{H}_2\text{O}) = \frac{1.63}{18} = 0.0905 = 9.1 \times 10^{-2}$ (1)
- (iv) $1.29 \times 10^{-2} \text{ mol ZnSO}_4$ are combined with $9.1 \times 10^{-2} \text{ mol H}_2\text{O}$
- 1 mol ZnSO_4 is combined with $\frac{9.1 \times 10^{-2}}{1.29 \times 10^{-2}}$
- = $7.054 \approx 7 \text{ mol H}_2\text{O}$
- answer must be expressed as a whole number
allow ecf on candidate's answers to (b)(ii) and (b)(iii) (1) [7]
- (c) (i) $n(\text{Zn}) = n(\text{CH}_3\text{CO}_2)_2\text{Zn} \cdot 2\text{H}_2\text{O}$ (1)
- $n(\text{Zn}) = \frac{0.015}{65.4} = 2.290 \times 10^{-4}$
- = 2.29×10^{-4} (1)
- mass of crystals = $2.29 \times 10^{-4} \times 219.4 = 0.0502655 \text{ g}$
= $0.05 \text{ g} = 50 \text{ mg}$ (1)
- (ii) concentration of $(\text{CH}_3\text{CO}_2)_2\text{Zn} \cdot 2\text{H}_2\text{O} = \frac{2.29 \times 10^{-4}}{0.005} = 0.0458$
= $4.58 \times 10^{-2} \text{ mol dm}^{-3}$ (1)
- allow correct answers if $\text{Zn} = 65$ is used [4]

[Total: 13]

Page 3	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – October/November 2012	9701	22

- 2 (a) (i) thermal stability decreases down Group VII (1)
- (ii) from Cl to I, atomic size increases **or**
the bonding pair is further from the nucleus of X **or**
H—X bond becomes longer **or**
smaller orbital overlap occurs (1)
hence H—X bond strength decreases down Group VII (1) [3]
- (b) $K_c = \frac{[\text{HI}]^2}{[\text{H}_2] \times [\text{I}_2]}$ (1)
- no units – must be clearly stated (1) [2]
- (c) (i) no change (1)
 K_c has no units **or**
same no. of molecules / moles each side of equilibrium (1)
- (ii) equilibrium moves to RHS (1)
 K_c increases with decreasing temperature **or**
forward reaction is exothermic **or**
reverse reaction is endothermic (1) [4]
- (d)
- | | | | | | | |
|----------------------------------|------------------------|---|------------------------|---|----------------|-----|
| | H ₂ (g) | + | I ₂ (g) | = | 2HI(g) | |
| initial moles | 0.02 | | 0.02 | | 0 | |
| equil. moles | (0.02 – y) | | (0.02 – y) | | 2y | (1) |
| equil. conc/mol dm ⁻³ | $\frac{(0.02 - y)}{1}$ | | $\frac{(0.02 - y)}{1}$ | | $\frac{2y}{1}$ | |
- $K_c = \frac{\text{HI}^2}{[\text{H}_2] \times [\text{I}_2]} = \frac{(2y)^2}{(0.02 - y)^2} = 59$ (1)
- $\frac{2y}{(0.02 - y)} = \sqrt{59} = 7.7$
- $2y = (7.7 \times 0.02) - 7.7y$
- $9.7y = 0.154$
- gives $y = \frac{0.154}{9.7} = 0.0159 = 0.016$ (1)
- at equilibrium**
- $n(\text{HI}) = 2 \times 0.016 = 0.032$ **and**
 $n(\text{H}_2) = n(\text{I}_2) = (0.02 - 0.016) = 0.004$ (1)
- allow ecf where possible [4]

[Total: 13]

Page 4	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – October/November 2012	9701	22

- 3 (a) (i) $\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightleftharpoons 2\text{NH}_3(\text{g})$ **or**
 $\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightarrow 2\text{NH}_3(\text{g})$
state symbols required (1)
- (ii) **pressure** between 60 and 250 atm **or**
between 60×10^5 Pa and 250×10^5 Pa (1)
- temperature** between 300 and 550 °C (1)
- catalyst** iron / iron oxide (1)
- (iii) manufacture of HNO_3 / as a cleaning agent / refrigerant / fertiliser / manufacture of fertilisers / explosives / to remove SO_2 from combustion products of hydrocarbon fuels (1) [5]
- (b) (i) NH_4Cl and $\text{Ca}(\text{OH})_2$
both formulae required (1)
- (ii) $2\text{NH}_4\text{Cl} + \text{Ca}(\text{OH})_2 \rightarrow \text{CaCl}_2 + 2\text{NH}_3 + 2\text{H}_2\text{O}$ **or**
 $\text{NH}_4^+ + \text{OH}^- \rightarrow \text{NH}_3 + \text{H}_2\text{O}$
correct products (1)
correctly balanced equation (1)
- (iii) CaO (1)
it is not an acid / it is basic / it does not react with NH_3 **or**
both P_2O_5 / P_4O_{10} **and** H_2SO_4 are acidic / react with NH_3 (1) [5]
- (c)
- $$\begin{array}{c} \text{H} \\ | \\ \text{H}-\text{N}: \\ | \\ \text{H} \end{array} + \text{H}^+ \longrightarrow \left[\begin{array}{c} \text{H} \\ | \\ \text{H}-\text{N} \rightarrow \text{H} \\ | \\ \text{H} \end{array} \right]^+$$
- correct displayed eqn.,
with positive charge clearly shown (1)
lone pair on NH_3 (1)
co-ordinate / dative bond clearly shown (1) [3]

[Total: 13]

Page 5	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – October/November 2012	9701	22

4 (a) (i)

reaction	organic compound	reagent	structural formulae of organic products
A	$(\text{CH}_3)_3\text{COH}$	$\text{Cr}_2\text{O}_7^{2-}/\text{H}^+$ heat under reflux	no reaction
B	$\text{CH}_3\text{CH}_2\text{CHO}$	Fehling's reagent warm	$\text{CH}_3\text{CH}_2\text{CO}_2\text{H}$ or $\text{CH}_3\text{CH}_2\text{CO}_2^-$
C	$\text{HCO}_2\text{CH}(\text{CH}_3)_2$	$\text{NaOH}(\text{aq})$ warm	HCO_2Na or HCO_2^- $(\text{CH}_3)_2\text{CHOH}$
D	$\text{CH}_2=\text{CHCHO}$	NaBH_4	$\text{CH}_2=\text{CHCH}_2\text{OH}$
E	$(\text{CH}_3)_3\text{COH}$	NaBH_4	no reaction
F	$\text{CH}_3\text{CH}_2\text{COCH}_3$	$\text{MnO}_4^-/\text{H}^+$ heat under reflux	no reaction

each correct answer gets (1)

(7 × 1)


(ii)

reaction	colour at the beginning of the reaction	colour at the end of the reaction
B	blue	brick red

each correct answer gets 1

(1 + 1 + 1) [10]

(b) (i)



(1)

(ii) red or orange


(1) [2]

[Total: 12]

Page 6	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – October/November 2012	9701	22

- 5 (a) (i) carboxylic acid **or** alcohol present **or** carboxylic acid **and** alcohol present **not** acid **or** carboxyl **or** hydroxyl (1)
- (ii) carboxylic acid **not** present **or** only alcohol present (1)
- (iii) alkene **or** $>C=C<$ present (1) [3]


(b) (i)



each correct structure gets (1)


(4 × 1)

(ii) pair 1 geometrical **or** *cis-trans* **or** *E/Z* isomerism (1)




pair 2 optical isomerism – accept chiral compounds (1) [6]

#1



#2



[Total: 9]