



Cambridge International AS & A Level

CHEMISTRY**9701/23**

Paper 2 AS Structured Questions

October/November 2020

MARK SCHEME

Maximum Mark: 60

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2020 series for most Cambridge IGCSE™, Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

This document consists of **11** printed pages.

PUBLISHED**Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

PUBLISHED**GENERIC MARKING PRINCIPLE 5:**

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

Science-Specific Marking Principles

1	Examiners should consider the context and scientific use of any keywords when awarding marks. Although keywords may be present, marks should not be awarded if the keywords are used incorrectly.
2	The examiner should not choose between contradictory statements given in the same question part, and credit should not be awarded for any correct statement that is contradicted within the same question part. Wrong science that is irrelevant to the question should be ignored.
3	Although spellings do not have to be correct, spellings of syllabus terms must allow for clear and unambiguous separation from other syllabus terms with which they may be confused (e.g. ethane / ethene, glucagon / glycogen, refraction / reflection).
4	The error carried forward (ecf) principle should be applied, where appropriate. If an incorrect answer is subsequently used in a scientifically correct way, the candidate should be awarded these subsequent marking points. Further guidance will be included in the mark scheme where necessary and any exceptions to this general principle will be noted.
5	<p><u>'List rule' guidance</u></p> <p>For questions that require <i>n</i> responses (e.g. State two reasons ...):</p> <ul style="list-style-type: none"> • The response should be read as continuous prose, even when numbered answer spaces are provided. • Any response marked <i>ignore</i> in the mark scheme should not count towards <i>n</i>. • Incorrect responses should not be awarded credit but will still count towards <i>n</i>. • Read the entire response to check for any responses that contradict those that would otherwise be credited. Credit should not be awarded for any responses that are contradicted within the rest of the response. Where two responses contradict one another, this should be treated as a single incorrect response. • Non-contradictory responses after the first <i>n</i> responses may be ignored even if they include incorrect science.

6 Calculation specific guidance

Correct answers to calculations should be given full credit even if there is no working or incorrect working, **unless** the question states 'show your working'.

For questions in which the number of significant figures required is not stated, credit should be awarded for correct answers when rounded by the examiner to the number of significant figures given in the mark scheme. This may not apply to measured values.

For answers given in standard form (e.g. $a \times 10^n$) in which the convention of restricting the value of the coefficient (a) to a value between 1 and 10 is not followed, credit may still be awarded if the answer can be converted to the answer given in the mark scheme.

Unless a separate mark is given for a unit, a missing or incorrect unit will normally mean that the final calculation mark is not awarded. Exceptions to this general principle will be noted in the mark scheme.

7 Guidance for chemical equations

Multiples / fractions of coefficients used in chemical equations are acceptable unless stated otherwise in the mark scheme.

State symbols given in an equation should be ignored unless asked for in the question or stated otherwise in the mark scheme.

Question	Answer	Marks
1(a)	$\text{Mg(g)} \rightarrow \text{Mg}^{\text{+}}(\text{g}) + \text{e}^{-}$	1
1(b)	M1: distance between nucleus and outer e⁻ increases OR outer electron removed from higher energy shell	3
	M2: increased shielding	
	M3: decreased nuclear attraction	
1(c)	M1: greater nuclear attraction	2
	M2: (2nd / 2s) electron being removed from smaller (ion)	


PUBLISHED


Question	Answer	Marks			
2(a)(i)	$P_4 + 5O_2 \rightarrow P_4O_{10}$	1			
2(a)(ii)	any two from: <ul style="list-style-type: none"> reacts vigorously solid disappears / colourless solution forms hydrolysis exothermic acid(ic) (solution) steamy / misty fumes 	2			
2(a)(iii)	Simple and covalent OR molecular and covalent	1			
2(b)(i)	M1: proton / H^+ donor	2			
	M2: partially dissociates (in solution)				
2(b)(ii)	$SO_2 + H_2O \rightarrow H_2SO_3$	1			
2(b)(iii)	(homogeneous) catalyst	1			
2(c)(i)	thermal decomposition	1			
2(c)(ii)	M1: $\Delta H_r = -1434 - (-635 + -297)$	2			
	M2: = -502 (kJ mol ⁻¹)				
2(d)(i)		1			
2(d)(ii)	<table border="1" style="display: inline-table; vertical-align: middle;"> <tr> <td style="padding: 2px 10px;">0</td> <td style="padding: 2px 10px;">(+4</td> <td style="padding: 2px 10px;">-1</td> </tr> </table>	0	(+4	-1	1
0	(+4	-1			

PUBLISHED

Question	Answer	Marks
2(d)(iii)	(at 1000 K and 100 kPa) M1: (yield) decreases	4
	M2: reaction is exothermic AND equilibrium moves left	
	(at 500 K and 500 kPa) M3: (yield) increases	
	M4: fewer moles (of gas) on right-hand side AND equilibrium moves right	
2(e)(i)	M1: ionic	2
	M2: ions only able / free to move / free to conduct (when liquid / molten)	
2(e)(ii)	M1: covalent	2
	M2: hydrolysed (by water)	

PUBLISHED


Question	Answer	Marks
3(a)(i)	<p>M1: correct representation of Al_2Cl_6, dot and cross or line diagram</p>  <p>M2: TWO correct co-ordinate bonds identified</p>	2
3(a)(ii)	120	1
3(a)(iii)	Li^+ is $1s^2$ H^- is $1s^2$	1
3(a)(iv)	(Lattice of) cations / positive ions surrounded by delocalised electrons'	1
3(b)	$Al(OH)_3$ / aluminium hydroxide	1
3(c)(i)	<p>M1: potassium dichromate[(VI)]</p> <p>M2: acid(ified) AND (heat under) reflux</p>	2
3(c)(ii)	<p>(M1: correct identity of R and statement re: reaction 3 ONLY ketone reduced)</p> <p>R (is 2-hydroxybutanoic acid) AND as (only) $C=O$ / ketone reduced</p> <hr/> <p>(M2: correct explanation re: strength of reducing agents) $NaBH_4$ cannot reduce the $COOH$ / carboxylic acid OR $LiAlH_4$ can reduce the $COOH$ / carboxylic acid</p>	2

Question	Answer	Marks
3(c)(iii)	 <p>M1: Presence of :CN (if bonding shown, must be unambiguous triple bond) M2: curly arrow from :CN lone pair to carbonyl carbon M3: correct dipole AND curly arrow from double bond to oxygen M4: correct intermediate drawn</p>	4
3(c)(iv)	$\text{C}_2\text{H}_5\text{CH}(\text{OH})\text{CN} + \text{HCl} + 2\text{H}_2\text{O} \rightarrow \text{C}_2\text{H}_5\text{CH}(\text{OH})\text{COOH} + \text{NH}_4\text{Cl}$	1
3(c)(v)	<p>Any two of three absorption references:</p> <ul style="list-style-type: none"> • absorption 2200–2250 (cm⁻¹) shows presence of C≡N • lack of absorption at 1680–1730 (cm⁻¹) shows lack of C=O • lack of absorption at 2500–3000 (cm⁻¹) shows lack of RCO₂-H / O-H in RCO₂H 	2

PUBLISHED

Question	Answer	Marks
4(a)(i)	M1: (Volatility) decreases (down the group)	2
	M2: more electrons so greater intermolecular forces / intermolecular attractions OR more electrons so greater VdW between molecules	
4(a)(ii)	(HI has the) lowest bond enthalpy	1
4(a)(iii)	M1: HF has permanent dipole(-dipole forces) AND HI has ((only)) instantaneous dipole / induced dipole (forces) / permanent dipole(-dipole forces)	2
	M2: IMF's in HI are weaker (than IMF's in HF)	
4(a)(iv)	$3\text{I}_2 + 6\text{NaOH} \rightarrow 5\text{NaI} + \text{NaIO}_3 + 3\text{H}_2\text{O}$	1

PUBLISHED

Question	Answer	Marks
4(b)(i)	HI(g) / PI ₃ / P and I ₂	1
4(b)(ii)	Electrophilic addition	1
4(c)(i)	2(-)iodo(-)2(-)methylbutane	1
4(c)(ii)	Nucleophilic substitution / S _N	1
4(c)(iii)		1
4(c)(iv)	(L has) two identical / two methyl groups attached to one end / one carbon of the C=C / double bond	1
4(c)(v)	ethanoic acid / CH ₃ COOH	1
4(c)(vi)	CH ₃ COCH ₃ + 3I ₂ + 4OH ⁻ → (1)CH ₃ COO ⁻ + 3H ₂ O + 3I ⁻ + CHI ₃ M1: correctly balanced M2: CHI ₃ product	2
4(c)(vii)	yellow ppt / yellow solid	1