

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS General Certificate of Education Advanced Subsidiary Level and Advanced Level

CANDIDATE NAME					
CENTRE NUMBER		CANDIDA ^T NUMBER	re [



CHEMISTRY 9701/22

Paper 2 Structured Questions AS Core

October/November 2011 1 hour 15 minutes

Candidates answer on the Question Paper.

Additional Materials: **Data Booklet**

READ THESE INSTRUCTIONS FIRST

Write your name, Centre number and candidate number on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams, graphs, or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE ON ANY BARCODES.

Answer all questions.

You may lose marks if you do not show your working or if you do not use appropriate units.

A Data Booklet is provided.

The number of marks is given in brackets [] at the end of each question or part question.

At the end of the examination, fasten all your work securely together.

For Exam	iner's Use
1	
2	
3	
4	
5	
Total	

This document consists of 11 printed pages and 1 blank page.



Answer all the questions in the space provided.

For Examiner's Use

1 Compound A is an organic compound which contains carbon, hydrogen and oxygen.

When 0.240 g of the vapour of **A** is slowly passed over a large quantity of heated copper(II) oxide, CuO, the organic compound **A** is completely oxidised to carbon dioxide and water. Copper is the only other product of the reaction.

The products are collected and it is found that $0.352\,\mathrm{g}$ of CO_2 and $0.144\,\mathrm{g}$ of $\mathrm{H}_2\mathrm{O}$ are formed.

- (a) In this section, give your answers to three decimal places.
 - (i) Calculate the mass of carbon present in 0.352 g of CO₂.

Use this value to calculate the amount, in moles, of carbon atoms present in 0.240 g of **A**.

(ii) Calculate the mass of hydrogen present in $0.144\,\mathrm{g}$ of $\mathrm{H}_2\mathrm{O}$.

Use this value to calculate the amount, in moles, of hydrogen atoms present in $0.240\,\mathrm{g}$ of \mathbf{A} .

(iii) Use your answers to calculate the mass of oxygen present in 0.240 g of A.

Use this value to calculate the amount, in moles, of oxygen atoms present in 0.240 g of **A**.


[6]

(b)	Use your answers to (a) to calculate the empirical formula of A.	For Examiner's Use
	[1]	
(c)	When a 0.148 g sample of $\bf A$ was vapourised at 60°C, the vapour occupied a volume of 67.7 cm ³ at a pressure of 101 kPa.	
	(i) Use the general gas equation $pV = nRT$ to calculate M_r of A .	
	$M_{r} = \dots$	
	(ii) Hence calculate the molecular formula of A.	
	[3]	
(d)		
	Suggest two structural formulae for A .	
	[2]	
(e)	Compound A contains only carbon, hydrogen and oxygen.	
	Explain how the information on the opposite page about the reaction of ${\bf A}$ with CuO confirms this statement.	
	[1]	
	[Total: 13]	

2 The Periodic Table we currently use is derived directly from that proposed in 1869 by Mendeleev who had noticed patterns in the physical and chemical properties of the elements he had studied.

For Examiner's Use

The diagram below shows the first ionisation energies of the first 18 elements of the Periodic Table.



(a)	Give	e the equation, including state symbols, for the first ionisation energy of sulfur.
		[2]
(b)		lain why there is a general increase in first ionisation energies across the Period sodium to argon.
		[3]
(c)	(i)	Explain why the first ionisation energy of magnesium is greater than that of aluminium.
	(ii)	Explain why the first ionisation energy of phosphorus is greater than that of sulfur.
		[4]

The table below refers to the elements of the third Period sodium to sulfur and is incomplete.

For Examiner's Use

element	Na	Mg	Al	Si	Р	S
conductivity			high			
melting point			high			

- (d) (i) Complete the 'conductivity' row by using only the words 'high', 'moderate' or 'low'.
 - (ii) Complete the 'melting point' row by using **only** the words 'high' or 'low'. [5

When Mendeleev published his first Periodic Table, he left gaps for elements that had yet to be discovered. He also predicted some of the physical and chemical properties of these undiscovered elements.

For one element, **E**, he correctly predicted the following properties.

melting point of the element high melting point of the oxide high

boiling point of the chloride low

The element **E** was in the fourth Period and was one of the elements from gallium, proton number 31, to bromine, proton number 35.

(e)	By considering the properties of the third Period elements aluminium to chlorine, suggest the identity of the fourth Period element E .
	[1]
	[Total: 15]

3		me chemical reactions, such as the thermal decomposition of potassium nearbonate, KHCO ₃ , the enthalpy change of reaction cannot be measured directly.	For Examiner's Use
		cases, the use of Hess' Law enables the enthalpy change of reaction to be calculated enthalpy changes of other reactions.	
	(a) Stat	te Hess' Law.	
		[2]	
		to determine the enthalpy change for the thermal decomposition of potassium ncarbonate, two separate experiments were carried out.	
	experim	ent 1	
	tempera When 0.	3 of 2.00 mol dm $^{-3}$ hydrochloric acid (an excess) was placed in a conical flask and the ture recorded as 21.0 °C0200 mol of potassium carbonate, $\rm K_2CO_3$, was added to the acid and the mixture with a thermometer, the maximum temperature recorded was 26.2 °C.	
	(b) (i)	Construct a balanced equation for this reaction.	
	(ii)	Calculate the quantity of heat produced in experiment 1 , stating your units. Use relevant data from the <i>Data Booklet</i> and assume that all solutions have the same specific heat capacity as water.	
	(iii)	Use your answer to (ii) to calculate the enthalpy change per mole of $\rm K_2CO_3$. Give your answer in kJ $\rm mol^{-1}$ and include a sign in your answer.	
	(iv)	Explain why the hydrochloric acid must be in an excess.	
		[4]	

					_
ex	മ	ru	me	nt	~
-	\sim			,,,,	_

For Examiner's Use

The experiment was repeated with 0.0200 mol of potassium hydrogen carbonate, ${\rm KHCO_3}.$ All other conditions were the same.

In the second experiment, the temperature fell from 21.0 °C to 17.3 °C.

- (c) (i) Construct a balanced equation for this reaction.
 - (ii) Calculate the quantity of heat absorbed in experiment 2.
 - (iii) Use your answer to (ii) to calculate the enthalpy change per mole of KHCO₃. Give your answer in kJ mol⁻¹ and include a sign in your answer.

[3]

(d) When ${\rm KHCO_3}$ is heated, it decomposes into ${\rm K_2CO_3}$, ${\rm CO_2}$ and ${\rm H_2O}$.

$$2\mathsf{KHCO_3} \mathop{\longrightarrow} \mathsf{K_2CO_3} + \mathsf{CO_2} + \mathsf{H_2O}$$

Use Hess' Law and your answers to **(b)(iii)** and **(c)(iii)** to calculate the enthalpy change for this reaction.

Give your answer in kJ mol⁻¹ and include a sign in your answer.

[2]


[Total: 11]

- **4** But-1-ene, CH₃CH₂CH=CH₂, is an important compound in the petrochemical industry.
 - (a) Some reactions of but-1-ene are given below.

Use

For Examiner's

In each empty box, draw the structural formula of the organic compound formed.



(b) Compound **T** reacts with compound **U**.

Draw the **displayed** formula of the organic product of this reaction.

For Examiner's Use

[2]

[Total: 7]

5	Astronomers using modern telescopes of various types have found many molecules in the dust clouds in space. Many of these molecules are those of organic compounds and astronomers constantly look for evidence that amino acids such as aminoethanoic acid, $\rm H_2NCH_2CO_2H$, are present.
	One molecule that has been found in the dust clouds is hydroxyethanal, $\mathrm{HOCH}_2\mathrm{CHO}.$
	(a) Hydroxyethanal contains two functional groups.

es in For and Examiner's acid, Use

One	one molecule that has been found in the dust clouds is hydroxyethanal, HOCH ₂ CHO.				
(a)	Hyd	Iroxyethanal contains t	wo functional groups.		
	(i)	Name, as fully as hydroxyethanal.	s you can, each of the functional groups present in		
		1			
		2			
	(ii)	react with the other fu	proup, identify a reagent that will react with this group and not unctional group present. The what would be observed when this reaction is carried out.		
		functional group 1	reagent		
			observation		
		functional group 2	reagent		
			observation[7]		
(b)		e the skeletal formula cted separately with the	e of the organic compounds formed when hydroxyethanal is e following.		
	(i)	NaBH ₄			

(ii) $\operatorname{Cr_2O_7^{2-}/H^+}$ under reflux conditions

[2]

In a	a school	or colle	ege la	aboratory,	it is	possible	to	convert	а	sample	of	hydroxy	ethana	۱£
into	aminoe	thanoic	acid	in a three	-step	process								

For Examiner's Use

$$\mathsf{HOCH_2CHO} \xrightarrow{\mathsf{step 1}} \mathbf{X} \xrightarrow{\mathsf{step 2}} \mathbf{Y} \xrightarrow{\mathsf{step 3}} \mathsf{H_2NCH_2CO_2H}$$

By considering the possible reactions of the functional groups present in hydroxyethanal, you are to deduce a possible route for this conversion.

(c) (i) In the boxes below, draw the structural formulae of your suggested intermediates X and Y.



(ii) State the reagents for each of the three steps you have chosen.

step	1
step	2
step	3

[5]

[Total: 14]

12

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.