

# Cambridge International AS & A Level

### Cambridge International Examinations

Cambridge International Advanced Subsidiary and Advanced Level

CANDIDATE NAME				
CENTRE NUMBER		CANDIDATE NUMBER		

CHEMISTRY 9701/22

Paper 2 AS Level Structured Questions

October/November 2016

1 hour 15 minutes

Candidates answer on the Question Paper.

Additional Materials: Data Booklet

#### **READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units.

A Data Booklet is provided.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.





Answer **all** the questions in the spaces provided.


	exce		ample of a Group 2 metal, <b>M</b> , was added to 40.0 cm <sup>3</sup> of 1.00 moldm <sup>-3</sup> hydrochloric acid s).
equa	atior	า 1	$\mathbf{M}(s) + 2HCl(aq) \rightarrow \mathbf{M}Cl_2(aq) + H_2(g)$
(a)	Cald	cul	late the amount, in moles, of hydrochloric acid present in 40.0 cm <sup>3</sup> of 1.00 mol dm <sup>-3</sup> HC <i>l</i> .
			amount = mol [1]
	Whe		the reaction had finished, the resulting solution was made up to 100 cm³ in a volumetric
	sod	liun	ocm <sup>3</sup> sample of the solution from the volumetric flask required 15.0 cm <sup>3</sup> of 0.050 mol dm <sup>-3</sup> carbonate solution, Na <sub>2</sub> CO <sub>3</sub> , for complete neutralisation of the remaining chloric acid.
	(i)	W	/rite the equation for the complete reaction of sodium carbonate with hydrochloric acid.
			[1]
1	(ii)		alculate the amount, in moles, of sodium carbonate needed to react with the ydrochloric acid in the 10.0 cm <sup>3</sup> sample from the volumetric flask.
			amount = mol [1]
(	iii)	Ca	alculate the amount, in moles, of hydrochloric acid in the 10.0 cm <sup>3</sup> sample.
			amount = mol [1]
(	iv)		alculate the total amount, in moles, of hydrochloric acid remaining after the reaction nown in equation 1.
			amount = mol [1]

(v)	Use your answers to <b>(a)</b> and <b>(b)(iv)</b> to calculate the amount, in moles, of hydrochloric acid that reacted with the 0.50 g sample of <b>M</b> .
(vi)	$amount = \ mol \ [1]$ Use your answer to <b>(v)</b> and equation 1 to calculate the amount, in moles, of <b>M</b> in the 0.50 g sample.
(vii)	${\rm amount} = {\rm mol} \ [1]$ Calculate the relative atomic mass, $A_{\rm r}$ , of ${\bf M}$ and identify ${\bf M}$ .
	A <sub>r</sub> of <b>M</b> =identity of <b>M</b> =[2]

**2** Dinitrogen tetraoxide, N<sub>2</sub>O<sub>4</sub>, and nitrogen dioxide, NO<sub>2</sub>, exist in dynamic equilibrium with each other.

$$N_2O_4(g) \iff 2NO_2(g)$$
  $\Delta H = +54 \text{ kJ mol}^{-1}$ 

The energy profile for this reaction is shown.



- (a) Add labelled arrows to the energy profile to indicate
  - the enthalpy change of the reaction,  $\Delta H$ ,
  - the activation energy of the forward reaction, E<sub>a</sub>.

[2]

- (b) 0.0500 mol of N<sub>2</sub>O<sub>4</sub> was placed in a sealed vessel of volume 1.00 dm<sup>3</sup>, at a temperature of 50 °C and a pressure of 1.68 × 10<sup>5</sup> Pa. The mass of the resulting equilibrium mixture was 4.606 g.
  - (i) Calculate the average molecular mass,  $M_r$ , of the resulting equilibrium mixture. Give your answer to **three** significant figures.

$$M_{\rm r} = \dots [2]$$

(ii) The number of moles of  $N_2O_4$  that dissociated can be represented by n.

State, in terms of n, the amount, in moles, of  $NO_2$  in the equilibrium mixture.

moles of 
$$NO_2 = \dots$$
 [1]

© UCLES 2016

The number of moles of N <sub>2</sub> O <sub>4</sub> remaining at equilibrium is (0.05)	5 – <i>n</i> ).
---	-----------------

(iii)	State, in terms of $n$ , the total amount, in moles, of gas in the equilibrium mixture.

[1]

(iv) State, in terms of n, the mole fraction of  $NO_2$  in the equilibrium mixture.

[1]

In this equilibrium mixture, the mole fraction of  $NO_2$  is 0.400.

(v) Use your answers to (ii) and (iv) to calculate the amount in moles of each gas in the equilibrium mixture. Give your answers to **three** significant figures.

amount of  $N_2O_4 = \dots$  mol

amount of NO<sub>2</sub> = ..... mol

[2]

(vi) Write the expression for the equilibrium constant,  $\mathcal{K}_{\!_{p}}$ , for this equilibrium.

 $K_p =$ 

[1]


(vii) Use the total pressure of the mixture,  $1.68 \times 10^5$  Pa, to calculate the value of the equilibrium constant,  $K_p$ , and give its units.

 $K_p = \dots$ 

units = .....[3]


[Total: 13]

- 3 The Periodic Table is arranged such that the properties of the elements show a number of trends.
  - (a) A plot of the first ionisation energies for the first 18 elements is shown.



(i)	Explain why the values show a general increase from atomic number 11 to 18.
	[2]
(ii)	Explain the decreases in first ionisation energies between
	• atomic numbers 12 and 13,
	atomic numbers 15 and 16.
	[4]
(iii)	Suggest an explanation for the trend in the first ionisation energies of the elements with atomic numbers 2, 10 and 18.
	[2]

(b) A plot of the melting points of the elements across the third period is shown.



(i)	Explain the increase in melting point from atomic number 11 to 12.
	[2]
(ii)	Suggest a reason why the increase from atomic number 12 to 13 is much smaller than the increase from atomic number 11 to 12.
	[1]
(iii)	State and explain the pattern of the melting points from atomic number 15 to 18.
	[3]
iv)	Explain why the element with atomic number 14 has a melting point so much higher than the rest of the elements in the third period.
	[1]

[Total: 15]

4	In each section of this question the structural formula of an organic compound is shown. For each compound answer the questions about it.					
	(a)	CH	3CH <sub>2</sub> CHBrCH <sub>3</sub>			
		(i)	Name this compound.			
			[	[1]		
		(ii)	This compound shows stereoisomerism.			
			Draw the <b>two</b> stereoisomers in the conventional way.			
				<b>ာ</b> 1		
		, <u>.</u>		[2]		
	(	(iii) —	Give the structures of <b>three</b> other structural isomers of C₄H <sub>9</sub> Br.			
	(b)	(C <sub>2</sub> I	H <sub>5</sub> ) <sub>3</sub> CBr	[3]		
		(i)	Name this compound.			
			[	1]		
		(ii)	(C₂H₅)₃CBr reacts with aqueous OH⁻.			
			Complete the mechanism for this reaction including all necessary curly arrows, charge partial charges and lone pairs.	s,		
	CH <sub>3</sub>	ÇH <sub>2</sub>				
CH <sub>3</sub>	CH <sub>2</sub>	CH   	-Br <del>→</del>			
	O1 13	ا ات 12		3]		
	(	(iii)	What type of mechanism occurs in (ii)?			
	·			11		
				•		

(c)	CH	3CH2CH2CHBrCH3						
	(i)	Give the reagents and conditions necessary for the conversion of this compound into mixture of alkenes.						
				[2]				
	(ii)	Give the name of	the mechanism for the conversion in (i	<b>)</b> . [1]				
	(iii)	Draw the skeletal	formulae of the three alkenes produce					

[3]

[Total: 17]

5

In eac	In each section of this question choose the answer or answers from the options listed.								
<b>(a)</b> Si	x particles are	listed.							
		Н∙	H⁺	Cl•	Cl-	•CH <sub>3</sub>	⁺CH <sub>3</sub>		
(i)	Identify <b>two</b> pof UV light.	oarticles p	roduce	ed during	the rea	ction of m	nethane and chloi	rine in the prese	ence
									. [1]
(ii)	Identify the t	<b>wo</b> particl	es pro	duced by	y the he	terolytic	fission of a bond	in chlorometha	ıne.
									. [1]
<b>(b)</b> Se	even reaction ty	/pes are li	sted.						
	а	ddition	subs	titution	oxida	ation	elimination		
		hydro	lysis	conde	ensatior	n redu	uction		
(i)	Name the typ	oe of reac	tion inv	olved w	hen Tol	lens' rea	gent is used to id	entify an aldeh	yde.
									. [1]
(ii)	Name the typ	oe of reac	tion inv	olved ir	the tes	t for a ca	rbonyl group usir	ng 2,4-DNPH.	
									. [1]
(iii)	Name the typ	oe of reac	tion inv	olved ir	the rea	action of	a ketone with Nal	BH₄.	
									. [1]
(iv)	Name the typ	oe of reac	tion inv	olved ir	the rea	action of a	an aldehyde with	HCN.	
									[1]
								[Tota	al: 6]

11

# **BLANK PAGE**

12

## **BLANK PAGE**

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.