UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS

GCE Advanced Subsidiary Level and GCE Advanced Level

MARK SCHEME for the October/November 2009 question paper for the guidance of teachers

9701 CHEMISTRY

9701/31

Paper 31 (Advanced Practical Skills), maximum raw mark 40

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

• CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the October/November 2009 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.



Page 2	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A/AS LEVEL – October/November 2009	9701	31

Que	estion	Sections	Indicative material	Mark	
1	(a)	PDO layout	Two balance readings and mass of FA 1 clearly recorded for each experiment. (Data for 2 nd experiment could be on page 4) Examiner to check subtraction for each experiment – no penalty in this section but see section (e)	1	[1]
foll be (i) (iv)	e experowing awardo) ii) & (ix)		If the candidate has only performed one experiment the following points only can be awarded: (ii), (iii), (vi), (vii) and (x). (i) Single table recording observations for both experiments. Times at ½ minute intervals. (ii) Appropriate headings and units Allow times in minutes (min) or seconds (iii) All temps recorded to nearest 0.5 °C (Must be more than one at .5 as well as .0) (iv) Some temps recorded before mixing and some after mixing for each expt. or Candidate records initial temperature and at least three temperatures after mixing for each experiment (v) First temperature after mixing is clearly taken 1 minute after adding the zinc powder (Examiner judgement re temperatures recorded before mixing / temperatures only recorded after mixing) and cooling for at least 5 minutes after recorded maximum temperature.	1 1 1 1	
		For Superviso	or - calculate mean maximum ∆T to nearest 0.5 °C; calculate mean of time taken (to nearest ½ min) to reac temperature after mixing.	h max	
		MMO Quality	 (vi) & (vii) 1st expt. Compare ΔT with Supervisor. award (vi) and (vii) if within 2 °C award (vii) only if >2 °C and ≤5°C (viii) & (ix) 2nd expt. Compare ΔT with Supervisor. award (viii) and (ix) if within 2 °C award (ix) only if >2 °C and ≤5°C (x) (1st expt) & (xi) (2nd expt). Compare time after mixing at which max temp is obtained with same time for Supervisor, for each expt. If Supervisor ≤3 min; 1 mark for Δ time ≤1 min. 	2 1 1	
			If Supervisor >3 min; 1 mark for Δ time \leq 1½ min.		[11]

Page 3	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A/AS LEVEL – October/November 2009	9701	31

Question	Sections	Indicative material	Mark	
(c)	PDO Layout	Plots temperature on <i>y</i> -axis and time on <i>x</i> -axis and has at least one temperature and one time label (<i>ignore absent or incorrect units</i>) Scales used are linear and easy for the examiner to use, (3 or 4 min. per large square are acceptable) Scales should enable the temperature when zinc is added	1	
A completely horizontal line, drawn at the initial temperature can be accepted as equivalent to plotting of initial temperature.		and <u>all</u> points after the addition of zinc to be plotted. Points should be within a minimum of 5 large squares on temperature axis If the candidate has recorded temperatures and times before zinc is added: Correctly plots on <u>each</u> graph: the last temperature/time, from results <u>before</u> zinc is added <u>or</u> the temperature and	1	
drawn, the marks may and one fur initial and memperature plotted and	be awarded ther mark if the naximum es are correctly	maximum temperature (associated time not required If the candidate has only recorded the initial temperature of the solution: Correctly plots on each graph: the temperature when zinc is added and maximum temperature (associated time not required Draws a cooling curve or straight line and projects the curve / line back to the time of mixing	1	[4]
(d)	ACE Interpretation	For experiment 1: Correctly reads the temperature rise from the graph to within 1 °C of the value obtained from the graph by the examiner. If the value is incorrect for experiment 1, check value for experiment 2. Award mark if either value is correct.	1	[1]
(e)–(h)	PDO Display	Shows working in all sections attempted – minimum of three sections required. Significant figures in final answers. 2 or 3 sf in 1(e), 2 to 4 sf in 1(g), 3 sf only in 1(h) minimum of three sections required.	1	[2]
(e)	ACE Interpretation	Correctly calculates 2.0 x 10 ⁻² mol of CuSO ₄ , and (mass zinc/65.4) for each experiment. Answers correctly rounded for the number of significant figures displayed. Do not award this mark if there is an error in subtraction or there are missing balance readings in section (a).	1	[1]

Page 4	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A/AS LEVEL – October/November 2009	9701	31

	Question	Sections	Indicative material	Mark	
	(f)	ACE Conclusions	To gain this mark the candidate must refer to: (i) the 1:1 mole ratio from the equation and (ii) the relative moles of Cu ²⁺ and Zn(s) used, as calculated in (e) If candidate states that "more moles of zinc were present" and this fits the calculated values in (e) – accept as the relative statement.	1	[1]
	(g)	ACE Interpretation	Shows (25 × 4.3 × candidate mean ∆T) _with appropriate unit, J or kJ, on final answer.	1	
maxim use he		as given a ture in (g) , allow o allow use of ΔT	(Allow use of 4.2 or 4.18 without penalty) Award this mark for the correct expression and unit OR where the expression is not shown, a correct evaluation of that expression and unit		[1]
	(h)	ACE Interpretation	No mark is awarded in this section if there is no division by (moles of zinc) or by (moles of Cu²+). Calculates	1	[2]
	(i)	ACE Interpretation	Candidate identifies one source of error in the experiment. This must be related to: Apparatus used or method described – no human error allowed. Heat loss is most likely error to be seen Accept reference to the graduation (precision) of the thermometer.	1	[1]
	(j)	ACE Improvement	Answer must follow on from (i) Suggests a way in which method could be improved e.g. Use of a lid or increased insulation to minimise heat loss.	1	[1]
	Qn 1	Total			[26]

Page 5	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A/AS LEVEL – October/November 2009	9701	31

Que	estion	Sections	Indicative material	Mark	
FA	3 is Mr	nSO ₄ (s); FA 4	is PbCO ₃ (s); FA 5 is CuCO ₃ (s)		
2	(a)	MMO Collection MMO Decisions	As FA 5 is heated, observes: green or blue solid turning black, or green/blue (solid) turning to a black solid or residue Tests gas given off with: limewater, in any of a glowing (not burning) splint, these tests red litmus paper	1	[2]
	(b)	MMO Collection	Observes each of the following: colourless solution with FA 3 or colourless solution with FA 4, and blue or green solution with FA 5 and observation of a gas evolved with FA 4 or with FA 5. [Second mark from (a) may be awarded here if not already given in (a)]	1	[1]
	(c)	ACE Conclusion	Identifies carbonate in FA 4 <u>or</u> FA 5 and refers to carbon dioxide; providing there has been evidence in the tests: a positive test with limewater, or effervescence with dilute acid or Identifies carbonate in FA 4 <u>or</u> FA 5 and refers to specific test for carbon dioxide and its result in the conclusion.	1	[1]

Page 6	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A/AS LEVEL – October/November 2009	9701	31

Question	Sections	Indicative material	Mark	
(d)	PDO Recording	(i) All observations in a single table. Both reagents are required There must be no repetition of "headings".	1	
	MMO Collection	(ii) Reports addition of reagents to excess whenever a precipitate is formed on first addition of the reagent. (Minimum of 2 ppt)	1	
		(iii) white / off-white / buff / (light or pale) brown precipitate with solution from FA 3 .	1	
		Precipitate insoluble in excess with both reagents and turning brown (light or pale brown precipitate darkening) recorded for at least one of the reagents (iv) Give one mark for both observations. FA 4 – white precipitate – both reagents. soluble in excess NaOH; insoluble in excess NH ₃ (aq). and	1	
	ACE Conclusions	FA 5 – blue precipitate – both reagents. insoluble in excess NaOH; soluble in excess NH ₃ (aq) or colour goes to dark/deep blue. Mark conclusions consequentially to observations. (v) Expected cations: FA 3 (Mn ²⁺) and FA 5 (Cu ²⁺) Minimum observations required:	1	
		Mn ²⁺ — off-white (buff, pale or light brown) ppt with each reagent but NOT from white ppt alone. Allow from white ppt turning brown.		
		Cu ²⁺ – blue ppt insoluble in excess NaOH or dark blue colour with aqueous NH ₃ (vi) Identifies Pb ²⁺ and Al ³⁺ as possible cations. (a single consequential ion is acceptable)	1	[6]

Page 7	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A/AS LEVEL – October/November 2009	9701	31

Question	Sections	Indicative material	Mark	
(e)	MMO Decisions MMO Collection	If no pair of ions is given in (d), no mark can be awarded in this section Mark consequentially Selects appropriate reagent to distinguish between any pair of cations identified in (d). For Pb ²⁺ /Al ³⁺ accept HCl, H ₂ SO ₄ Kl or chromate/dichromate The candidate should name a reagent, e.g. potassium dichromate. If Cr ₂ O ₇ ²⁻ , or dichromate is given as the reagent the (aq) state symbol must also be given or reference made to an aqueous solution of the ions. If selected reagent is suitable; mark consequentially for chosen reagent and Pb ²⁺ . For Pb ²⁺ /Al ³⁺ FA 4 gives white precipitate with HCl and with H ₂ SO ₄ and	1	
		yellow precipitate with chromate/dichromate or iodide. Ignore any conclusion.		[2]
(f)	MMO Collection	Observes as only reaction: FA 3 gives white precipitate with Ba(NO ₃) ₂ which is insoluble in dilute nitric acid, but Ignore any white ppt or cloudiness with FA 3 and silver nitrate, and ignore cation precipitates on adding NH ₃ (aq)	1	
	ACE Conclusions	Accept a dash in the boxes for reaction of FA 3, FA 4 and FA 5 with barium nitrate and with silver chloride as evidence of "no reaction" with that reagent A conclusion that fits observations for (i) barium chloride with all solutions or (ii) silver nitrate with all solutions	1	[2]
Qn 2	Total	<u>I</u>		[14]