



Cambridge International AS & A Level

CANDIDATE NAME		
CENTRE NUMBER	CANDIDATE NUMBER	

CHEMISTRY

9701/35

Paper 3 Advanced Practical Skills 1

May/June 2020

2 hours

You must answer on the question paper.

You will need: The materials and apparatus listed in the confidential instructions

INSTRUCTIONS

- Answer all questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do not write on any bar codes.
- You may use a calculator.
- You should show all your working, use appropriate units and use an appropriate number of significant figures.
- Give details of the practical session and laboratory, where appropriate, in the boxes provided.

INFORMATION

- The total mark for this paper is 40.
- The number of marks for each question or part question is shown in brackets [].
- The Periodic Table is printed in the question paper.
- Notes for use in qualitative analysis are provided in the question paper.

Ses	ssion
Labo	oratory

For Examiner's Use	
1	
2	
3	
Total	

This document has 12 pages. Blank pages are indicated.

Quantitative Analysis

Read through the whole method before starting any practical work. Where appropriate, prepare a table for your results in the space provided.

Show your working and appropriate significant figures in the final answer to **each** step of your calculations.

1 Ethanedioic acid forms salts with Group 1 metals. In this experiment you will identify the Group 1 metal ion, Z⁺, present in an ethanedioate salt, (COO)₂Z₂. You will titrate a solution of the salt with acidified aqueous potassium manganate(VII). The equation for the reaction between manganate(VII) ions and ethanedioate ions in acidic solution is shown.

$$2MnO_4^{-}(aq) + 16H^{+}(aq) + 5(COO)_2^{2-}(aq) \rightarrow 2Mn^{2+}(aq) + 8H_2O(I) + 10CO_2(g)$$

FA 1 is 0.0200 mol dm⁻³ potassium manganate(VII), KMnO₄.

FA 2 is a solution containing 8.06 g dm⁻³ of an ethanedioate salt, (COO)₂Z₂.

FA 3 is dilute sulfuric acid, H₂SO₄.

(a) Method

- Fill the burette with **FA 1**.
- Pipette 25.0 cm³ of FA 2 into a conical flask.
- Use the measuring cylinder to transfer 25 cm³ of **FA 3** into the same conical flask.
- Place the conical flask on the tripod and gauze and heat the conical flask until the temperature of the solution is approximately 70 °C.
- Carefully remove the hot conical flask and place it on the white tile under the burette.
- During titrations, add FA 1, slowly at first, until a permanent pale pink colour is formed. (The pink colour on initial addition may take several seconds to disappear.) If the reaction mixture turns brown, reheat it to approximately 70°C. If the brown colour disappears, continue with the titration. If the brown colour remains, discard the contents of the flask and begin a new titration.
- Perform a rough titration (the end-point is a permanent pale pink colour) and record your burette readings in the space below.

The rough titre iscm³.

- Carry out as many accurate titrations as you think necessary to obtain consistent results.
- Make sure any recorded results show the precision of your practical work.
- Record all of your burette readings and the volume of FA 1 added in each accurate titration.

I	
II	
III	
IV	
V	
VI	
VII	

[7]

(b)	From your accurate titration results, obtain a suitable value for the volume of FA 1 to be used in your calculations. Show clearly how you obtained this value.		
		25.0 cm ³ of FA 2 required cm ³ of FA 1 . [1]	
(c)	Cal	lculations	
	(i)	Give your answers to (c)(ii) , (c)(iii) , (c)(iv) and (c)(v) to the appropriate number of significant figures.	
	(ii)	Calculate the number of moles of manganate (VII) ions in the volume of ${\bf FA~1}$ calculated in ${\bf (b)}.$	
		moles of $MnO_4^- = mol [1]$	
((iii)	Use the equation on page 2 to calculate the number of moles of ethanedioate ions in $25.0\mathrm{cm^3}$ of FA 2 .	
		moles of $(COO)_2^{2-} = \dots mol [1]$	
((iv)	Calculate the relative formula mass, M_r , of the ethanedioate salt, $(COO)_2Z_2$.	
		<i>M</i> _r of ethanedioate salt = [1]	
	(v)	Calculate the relative atomic mass, A_r , of the Group 1 metal, Z , in the ethanedioate salt. Show your working.	
		A_{r} of $Z = \dots$	
		Hence identify Z.	
		Z is	
		[2]	
		[Total: 14]	

2 In this experiment you will determine the value of x in the formula for hydrated manganese(II) sulfate, MnSO₄•xH₂O, where x is an integer. You will do this by measuring the mass lost when a sample of hydrated manganese(II) sulfate is heated.

$$MnSO_4 \bullet xH_2O(s) \rightarrow MnSO_4(s) + xH_2O(g)$$

FA 4 is hydrated manganese(II) sulfate, MnSO₄•xH₂O.

(a) Method

- Weigh the crucible with a lid and record the mass.
- Add all the FA 4 to the crucible.
- Reweigh the crucible with the lid and FA 4. Record the mass. Describe the appearance of FA 4.

appearance of FA 4

- Place the crucible in the pipe-clay triangle on top of the tripod.
- Heat the crucible **gently** with the lid on for approximately 1 minute.
- Remove the lid and then heat more strongly for a further 4 minutes.
- Replace the lid and allow the crucible to cool.
- While the crucible is cooling you may wish to begin work on Question 3.
- Once the crucible has cooled, reweigh the crucible with the lid and contents. Record the mass.
- Calculate and record the mass of FA 4 added to the crucible, the mass of the residue and the mass of water lost.
- Describe the appearance of the residue.

appearance of the residue	
---------------------------	--

I	
II	
III	
IV	
V	
VI	

[6]

(b) Calculations

	(i)	Calculate the number of moles of manganese(II) sulfate present in the residue. You may assume all the water of crystallisation has been removed.
		moles of $MnSO_4 = \dots mol$ [1]
((ii)	Calculate the number of moles of water lost.
		moles of water lost = mol [1]
(i	iii)	Calculate the value of x in MnSO ₄ •xH ₂ O.
		x =[1]
(c)	It is	possible that FA 4 did not lose all of the water of crystallisation in your experiment.
	(i)	Explain how you could modify the experiment to ensure all water has been removed.
		[1]
((ii)	Explain why your calculated value of x might not change if a small amount of water of crystallisation remained in the residue.
		[1]
		[Total: 11]

[3]

Qualitative Analysis

Where reagents are selected for use in a test, the **name** or **correct formula** of the element or compound must be given.

At each stage of any test you are to record details of the following:

- colour changes seen
- the formation of any precipitate and its solubility in an excess of the reagent added
- the formation of any gas and its identification by a suitable test.

You should indicate clearly at what stage in a test a change occurs.

If any solution is warmed, a **boiling tube** must be used.

Rinse and reuse test-tubes and boiling tubes where possible.

No additional tests for ions present should be attempted.

- 3 Half fill the 250 cm³ beaker with water and place it on a tripod and gauze. Heat until the water begins to boil then switch off your Bunsen burner. This is the hot water bath for part **(b)**.
 - (a) FA 5 is a solution of a salt which contains one cation and at least one anion, all of which are listed in the Qualitative Analysis Notes. Sulfur is not present in FA 5.

(i)	To a 1 cm depth of FA 5 in a test-tube add aqueous sodium hydroxide.		
	observation		
		[1]	
(ii)	You are to carry out tests to allow you to determine the anion present in FA 5 .		

Identify reagents for these tests, carry out these tests and record these tests and

© UCLES 2020 9701/35/M/J/20

observations in a table.

(iii)	Use your results to identify the ions present in FA 5 .		
	formulae of ions present in FA 5	[1]
(iv)	Write an ionic equation for the expedinclude state symbols.	cted reaction between FA 5 and aqueous ammonia	а.
(v)	Carry out the following tests and reco	ord your observations.	1]
	test	observations	
	Test 1 To a 1 cm depth of FA 5 in a test-tube, add a 1 cm depth of hydrogen peroxide, then		
	add aqueous sodium hydroxide.		
(:\			2]
(vi)	Suggest what type of reaction occurr	ed when hydrogen peroxide was added to FA 5 .	
			1]

(b)	FA 6, FA 7 and FA 8 are butan-1-ol,	butan-2-ol and methylpropan-2-ol, but not necessarily in
	that order.	

(i) Carry out the following tests and record your observations.

toot	observations						
test	FA 6	FA 7	FA 8				
Test 1 To a 1 cm depth of dilute sulfuric acid in a test-tube, add 2 or 3 drops of FA 1, KMnO ₄ , then add a few drops of the alcohol. Shake the tube and place it in the hot water bath. Shake the tube occasionally until there is no further change.							
Test 2 To a 1 cm depth of aqueous iodine in a test-tube, add a few drops of the alcohol, then add drops of aqueous sodium hydroxide until the iodine colour just disappears or remains unchanged. Place the test-tube in the hot water bath.							

[3]

(ii) Use your observations from (b)(i) to identify the alcohols.

alcohol	FA
butan-1-ol	FA
butan-2-ol	FA
methylpropan-2-ol	FA

г	-	7
	٠,	
	_	

iii)	Write an equation for the oxidation of one of these alcohols with acidified KMnO ₄ .
	Use [O] to represent the oxidising agent.

[Total: 15]

9

BLANK PAGE

Qualitative Analysis Notes

1 Reactions of aqueous cations

ia la	react	ion with
ion	NaOH(aq)	NH ₃ (aq)
aluminium, Al³+(aq)	white ppt. soluble in excess	white ppt. insoluble in excess
ammonium, NH₄⁺(aq)	no ppt. ammonia produced on heating	_
barium, Ba ²⁺ (aq)	faint white ppt. is nearly always observed unless reagents are pure	no ppt.
calcium, Ca ²⁺ (aq)	white ppt. with high [Ca ²⁺ (aq)]	no ppt.
chromium(III), Cr³+(aq)	grey-green ppt. soluble in excess	grey-green ppt. insoluble in excess
copper(II), Cu ²⁺ (aq)	pale blue ppt. insoluble in excess	blue ppt. soluble in excess giving dark blue solution
iron(II), Fe²+(aq)	green ppt. turning brown on contact with air insoluble in excess	green ppt. turning brown on contact with air insoluble in excess
iron(III), Fe³+(aq)	red-brown ppt. insoluble in excess	red-brown ppt. insoluble in excess
magnesium, Mg²+(aq)	white ppt. insoluble in excess	white ppt. insoluble in excess
manganese(II), Mn ²⁺ (aq)	off-white ppt. rapidly turning brown on contact with air insoluble in excess	off-white ppt. rapidly turning brown on contact with air insoluble in excess
zinc, Zn²+(aq)	white ppt. soluble in excess	white ppt. soluble in excess

2 Reactions of anions

ion	reaction
carbonate, CO ₃ ²⁻	CO ₂ liberated by dilute acids
chloride, C <i>l</i> ⁻ (aq)	gives white ppt. with Ag ⁺ (aq) (soluble in NH ₃ (aq))
bromide, Br ⁻ (aq)	gives cream ppt. with Ag ⁺ (aq) (partially soluble in NH ₃ (aq))
iodide, I ⁻ (aq)	gives yellow ppt. with Ag+(aq) (insoluble in NH ₃ (aq))
nitrate, NO ₃ -(aq)	NH ₃ liberated on heating with OH ⁻ (aq) and A <i>l</i> foil
nitrite, NO ₂ -(aq)	NH₃ liberated on heating with OH⁻(aq) and A <i>l</i> foil
sulfate, SO ₄ ²⁻ (aq)	gives white ppt. with Ba ²⁺ (aq) (insoluble in excess dilute strong acids)
sulfite, SO ₃ ²⁻ (aq)	gives white ppt. with Ba ²⁺ (aq) (soluble in excess dilute strong acids)

3 Tests for gases

gas	test and test result
ammonia, NH ₃	turns damp red litmus paper blue
carbon dioxide, CO ₂	gives a white ppt. with limewater (ppt. dissolves with excess CO ₂)
chlorine, Cl ₂	bleaches damp litmus paper
hydrogen, H ₂	'pops' with a lighted splint
oxygen, O ₂	relights a glowing splint

The Periodic Table of Elements

	~		a١	Ę C		رن و	Ę 0		_	ر 6	<i>'</i> ~	_	1 Pun 8	_	m	no E.	<i>"</i>	_	uc				
	18	2	Ĭ	heliu 4	101	ž	20.	18	⋖	argon 39.9	36		krypt 83.	54	×	xenc 131	86	~	radc				
	17				6	ш	fluorine 19.0	17	Cl	chlorine 35.5	32	Ŗ	bromine 79.9	53	Ι	iodine 126.9	85	Αţ	astatine -				
	16				80	0	oxygen 16.0	16	S	sulfur 32.1	34	Se	selenium 79.0	52	<u>e</u>	tellurium 127.6	84	Ро	moloulum —	116	_	livermorium	ı
	15				7	z	nitrogen 14.0	15	۵	phosphorus 31.0	33	As	arsenic 74.9	51	Sp	antimony 121.8	83	Ξ	bismuth 209.0				
	14				9	O	carbon 12.0	14	S	silicon 28.1	32	Ge	germanium 72.6	20	S	tin 118.7	82	Pb	lead 207.2	114	Εl	flerovium	
	13				2	В	boron 10.8	13	Αl	aluminium 27.0	31	Ga	gallium 69.7	49	In	indium 114.8	81	1L	thallium 204.4				
										12	30	Zu	zinc 65.4	48	В	cadmium 112.4	80	Ε̈́Η	mercury 200.6	112	ပ်	copernicium	-
										1	29	D O	copper 63.5	47	Ag	silver 107.9	62	Au	gold 197.0	111	Rg	roentgenium	-
dn										10	28	Z	nickel 58.7	46	Pq	palladium 106.4	78	₹	platinum 195.1	110	Ds	darmstadtium	-
Group										o	27	ပိ	cobalt 58.9	45	R	rhodium 102.9	77	ī	iridium 192.2	109	¥	meitnerium	-
		-	I	hydrogen	!					80	26	Fe	iron 55.8	44	Ru	ruthenium 101.1	92	SO	osmium 190.2	108	ΗS	hassium	1
					_					7	25	M	manganese 54.9	43	ပ	technetium -	75	Re	rhenium 186.2	107	뮵	bohrium	1
						0	SS			9	24	ပ်	chromium 52.0	42	Mo	molybdenum 95.9	74	≥	tungsten 183.8	106	Sg	seaborgium	1
				Kev	atomic number	atomic symbo	name relative atomic mass			2	23	>	vanadium 50.9	41	qN	niobium 92.9	73	Б	tantalum 180.9	105	op O	dubnium	-
					at	ator	relat			4	22	F	titanium 47.9	40	Zr	zirconium 91.2	72	둧	hafnium 178.5	104	꿒	rutherfordium	
								1		ဗ	21	လွ	scandium 45.0	39	>	yttrium 88.9	57-71	lanthanoids		89-103	actinoids		
	2				4	Be	beryllium 9.0	12	Mg	magnesium 24.3	20	Ca	calcium 40.1	38	Š	strontium 87.6	56	Ва	barium 137.3	88	Ra	radium	-
	_				8	:=	lithium 6.9	1	Na	sodium 23.0	19	×	potassium 39.1	37	Rb	rubidium 85.5	55	Cs	caesium 132.9	87	占	francium	-

Lu	lutetium 175.0	103	۲	lawrencium	ı
°° X	ytterbium 173.1	102	8	nobelium	ı
e9 L	thulium 168.9	101	Md	mendelevium	ı
₈₈ <u>п</u>	erbium 167.3	100	Fm	ferminm	ı
67 Ho	holmium 164.9	66	Es	einsteinium	ı
® Q	dysprosium 162.5	86	ర	californium	ı
c5 dT	terbium 158.9	26	益	berkelium	ı
² PO	gadolinium 157.3	96	S	curium	ı
En Eu	europium 152.0	96	Am	americium	ı
ss Sm	samarium 150.4	94	Pn	plutonium	1
Pm	promethium —	93	ď	neptunium	ı
⁰⁹ PN	neodymium 144.4	92	\supset	uranium	238.0
59 P	praseodymium 140.9	91	Ра	protactinium	231.0
Se Se	cerium 140.1	06	ᄕ	thorium	232.0
57 La	lanthanum 138.9	68	Ac	actinium	1

lanthanoids actinoids

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.