



Cambridge International AS & A Level

CANDIDATE
NAME

--	--	--	--	--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

* 3 3 3 0 9 5 4 0 0 9 *

CHEMISTRY

9701/33

Paper 3 Advanced Practical Skills 1

October/November 2020

2 hours

You must answer on the question paper.

You will need: The materials and apparatus listed in the confidential instructions

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You may use a calculator.
- You should show all your working, use appropriate units and use an appropriate number of significant figures.
- Give details of the practical session and laboratory, where appropriate, in the boxes provided.

Session
Laboratory

INFORMATION

- The total mark for this paper is 40.
- The number of marks for each question or part question is shown in brackets [].
- The Periodic Table is printed in the question paper.
- Notes for use in qualitative analysis are provided in the question paper.

For Examiner's Use	
1	
2	
Total	

This document has **16** pages. Blank pages are indicated.


BLANK PAGE

Quantitative Analysis

Read through the whole method before starting any practical work. Where appropriate, prepare a table for your results in the space provided on page 4.

Show your working and appropriate significant figures in the final answer to **each** step of your calculations.

- 1** In acidic solutions iron(III) ions are reduced by iodide ions to form iron(II) ions. The iodide ions are oxidised to iodine.



The rate of this reaction can be investigated by using starch indicator, which turns blue-black in the presence of iodine. Sodium thiosulfate is added to the reaction mixture to react with iodine as it is formed. The blue-black colour is seen when all the thiosulfate has reacted.



You will investigate how the rate of reaction is affected by changing the concentration of the iodide ions.

FA 1 is 0.0500 mol dm⁻³ potassium iodide, KI.

FA 2 is 0.0500 mol dm⁻³ acidified iron(III) chloride, FeCl₃.

FA 3 is 0.00500 mol dm⁻³ sodium thiosulfate, Na₂S₂O₃.

FA 4 is starch indicator.

(a) Method

Prepare a table on page 4 for your results. You will need to include the volume of **FA 1**, volume of water, reaction time and rate of reaction for each of five experiments.

Experiment 1

- Fill the burette labelled **FA 1** with **FA 1**.
- Run 20.00 cm³ of **FA 1** into the 100 cm³ beaker.
- Use the 50 cm³ measuring cylinder to add the following to the same 100 cm³ beaker:
 - 20.0 cm³ of **FA 3**
 - 10.0 cm³ of **FA 4**.
- Use the 25 cm³ measuring cylinder to measure 10.0 cm³ of **FA 2**.
- Add this **FA 2** into the same 100 cm³ beaker and start timing immediately.
- Stir once and place the beaker on the white tile.
- Stop timing as soon as the solution turns blue-black.
- Record this reaction time to the nearest second.
- Wash out the beaker and dry it with a paper towel.

Experiment 2

- Fill the second burette with distilled water.
- Run 10.00 cm³ of **FA 1** into the 100 cm³ beaker.
- Run 10.00 cm³ of distilled water into the beaker containing **FA 1**.
- Use the 50 cm³ measuring cylinder to add the following to the same 100 cm³ beaker:
 - 20.0 cm³ of **FA 3**
 - 10.0 cm³ of **FA 4**.
- Use the 25 cm³ measuring cylinder to measure 10.0 cm³ of **FA 2**.
- Add the **FA 2** to the same 100 cm³ beaker and start timing immediately.
- Stir once and place the beaker on the white tile.
- Stop timing as soon as the solution turns blue-black.
- Record this reaction time to the nearest second.
- Wash out the beaker and dry it with a paper towel.

Experiments 3–5

- Carry out three further experiments to investigate how the reaction time changes with different volumes of potassium iodide, **FA 1**.
The combined volume of **FA 1** and distilled water must always be 20.00 cm³.
Do not use a volume of **FA 1** that is less than 6.00 cm³.

Results


The rate of reaction can be calculated as shown:

$$\text{rate} = \frac{1000}{\text{reaction time}}$$

I	
II	
III	
IV	
V	
VI	
VII	
VIII	
IX	
X	

[10]

- (b) On the grid opposite, plot a graph of rate of reaction (y-axis) against volume of **FA 1** (x-axis). Include the origin, (0,0), in your scales. Circle any points you consider anomalous and draw a line of best fit. [3]



- (c) Use your graph to calculate the time that the reaction would have taken if 5.00 cm³ of **FA 1** had been used. Show on the graph how you obtained your answer.

time = s [2]

- (d) (i) Using data from **Experiments 1** and **2**, show by calculation that the volume of aqueous potassium iodide, **FA 1**, used was directly proportional to the concentration of iodide ions.

[2]

- (ii) Explain, by referring to your graph or your table of results, how the rate of reaction is affected by an increase in the concentration of aqueous potassium iodide, **FA 1**.

.....
.....
.....
.....

[2]

- (e) Thiosulfate ions can reduce iron(III) ions and also react with acid to form sulfur, sulfur dioxide and water.

- (i) Write an ionic equation for the reaction between thiosulfate ions and hydrogen ions in aqueous solution. Include state symbols.

..... [1]

- (ii) A student carries out the same investigation as in (a) but the solutions are mixed in a different order. The student places **FA 1** and an appropriate volume of distilled water in one beaker and all the other reactants in a second beaker. The student then transfers the mixture from the second beaker to the first and starts timing.

Tick the box for the statement you consider correct. Explain your answer.

The student's method is better than that in (a).

The two methods are equally good.

The student's method is not as good as that in (a).

reason

..... [1]

- (f) Another student investigates the effect of iron(III) concentration on the rate of this reaction. The student carries out another experiment, **Experiment 6**, and the rate is compared to that of **Experiment 2**. In **Experiment 2**, the volumes used were:

reagent	volume/cm ³
FA 1	10.00
FA 2	10.0
FA 3	20.0
FA 4	10.0
distilled water	10.00

- (i) Suggest the volumes the student could use for **Experiment 6**.

reagent	volume/cm ³
FA 1	
FA 2	
FA 3	
FA 4	
distilled water	

[2]

- (ii) This student records a time of 178 s for **Experiment 2**.

The rate of reaction is directly proportional to the concentration of iron(III) ions.

Suggest how long it would take the reaction mixture proposed for **Experiment 6** in (f)(i) to turn blue-black. Assume that **Experiment 6** is carried out at the same temperature as **Experiment 2**.

Do not carry out Experiment 6.

time = s [1]

[Total: 24]

Qualitative Analysis

Where reagents are selected for use in a test, the **name or correct formula** of the element or compound must be given.

At each stage of any test you are to record details of the following:

- colour changes seen
- the formation of any precipitate and its solubility in an excess of the reagent added
- the formation of any gas and its identification by a suitable test.

You should indicate clearly at what stage in a test a change occurs.

If any solution is warmed, a **boiling tube** must be used.

Rinse and reuse test-tubes and boiling tubes where possible.

No additional tests for ions present should be attempted.

2 (a) FA 5 is a salt consisting of two ions both of which are listed in the Qualitative Analysis Notes.

- (i) Place a small spatula measure of **FA 5** into a hard-glass test-tube. Heat the tube gently at first and then more strongly. Record **all** your observations.

.....
.....
.....
..... [3]

- (ii) **FA 6** is a sample of the residue obtained from **FA 5** after strong heating.

Place a small spatula measure of **FA 6** in a test-tube and add a 1 cm depth of aqueous hydrogen peroxide. Record your observations.

.....
..... [2]

- (iii) Dissolve a spatula measure of **FA 5** in a 5cm depth of distilled water in a boiling tube. Use separate 1 cm depths of this solution in test-tubes for the following tests. Record your observations.

Keep the remainder of FA 5(aq) for use in 2(b)(ii).

<i>test</i>	<i>observations</i>
Test 1 Add aqueous sodium hydroxide.	
Test 2 Add a 1 cm depth of aqueous hydrogen peroxide, then add aqueous sodium hydroxide.	

[2]

- (b) (i) **FA 7** is a solution of a different salt. The cation present in **FA 7** is not listed in the Qualitative Analysis Notes.

FA 5(aq) and **FA 7** each contain either a halide ion or an anion containing sulfur. These anions are listed in the Qualitative Analysis Notes.

For both of these anions, select reagents that you would use in order to carry out tests that give positive results.

Record the reagents and the ions for which they would test.

[2]

- (ii) Carry out both of your tests on **FA 5(aq)** and **FA 7** and record your results in the space below.

[3]

10

- (iii) Use your observations in (a) and (b)(ii) to identify the ions present in **FA 5** and **FA 7**. Write the formula of each ion in the table. If the tests you carry out do not allow you to identify any of the ions, write ‘unknown’.

	FA 5	FA 7
cation		
anion		

[2]

- (iv) Suggest what you would observe if you added aqueous chlorine to separate portions of aqueous solutions of **FA 5** and **FA 7**.

Do not carry out this test.

aqueous chlorine and **FA 5(aq)**

aqueous chlorine and **FA 7(aq)**

[2]

[Total: 16]

12

BLANK PAGE

13

BLANK PAGE

Qualitative Analysis Notes

1 Reactions of aqueous cations

ion	<i>reaction with</i>	
	NaOH(aq)	NH ₃ (aq)
aluminium, Al ³⁺ (aq)	white ppt. soluble in excess	white ppt. insoluble in excess
ammonium, NH ₄ ⁺ (aq)	no ppt. ammonia produced on heating	—
barium, Ba ²⁺ (aq)	faint white ppt. is nearly always observed unless reagents are pure	no ppt.
calcium, Ca ²⁺ (aq)	white ppt. with high [Ca ²⁺ (aq)]	no ppt.
chromium(III), Cr ³⁺ (aq)	grey-green ppt. soluble in excess	grey-green ppt. insoluble in excess
copper(II), Cu ²⁺ (aq)	pale blue ppt. insoluble in excess	blue ppt. soluble in excess giving dark blue solution
iron(II), Fe ²⁺ (aq)	green ppt. turning brown on contact with air insoluble in excess	green ppt. turning brown on contact with air insoluble in excess
iron(III), Fe ³⁺ (aq)	red-brown ppt. insoluble in excess	red-brown ppt. insoluble in excess
magnesium, Mg ²⁺ (aq)	white ppt. insoluble in excess	white ppt. insoluble in excess
manganese(II), Mn ²⁺ (aq)	off-white ppt. rapidly turning brown on contact with air insoluble in excess	off-white ppt. rapidly turning brown on contact with air insoluble in excess
zinc, Zn ²⁺ (aq)	white ppt. soluble in excess	white ppt. soluble in excess

2 Reactions of anions

<i>ion</i>	<i>reaction</i>
carbonate, CO_3^{2-}	CO_2 liberated by dilute acids
chloride, Cl^- (aq)	gives white ppt. with Ag^+ (aq) (soluble in NH_3 (aq))
bromide, Br^- (aq)	gives cream ppt. with Ag^+ (aq) (partially soluble in NH_3 (aq))
iodide, I^- (aq)	gives yellow ppt. with Ag^+ (aq) (insoluble in NH_3 (aq))
nitrate, NO_3^- (aq)	NH_3 liberated on heating with OH^- (aq) and Al foil
nitrite, NO_2^- (aq)	NH_3 liberated on heating with OH^- (aq) and Al foil
sulfate, SO_4^{2-} (aq)	gives white ppt. with Ba^{2+} (aq) (insoluble in excess dilute strong acids)
sulfite, SO_3^{2-} (aq)	gives white ppt. with Ba^{2+} (aq) (soluble in excess dilute strong acids)

3 Tests for gases

<i>gas</i>	<i>test and test result</i>
ammonia, NH_3	turns damp red litmus paper blue
carbon dioxide, CO_2	gives a white ppt. with limewater (ppt. dissolves with excess CO_2)
chlorine, Cl_2	bleaches damp litmus paper
hydrogen, H_2	'pops' with a lighted splint
oxygen, O_2	relights a glowing splint

The Periodic Table of Elements

		Group																																																	
1	2																																																		
Key																																																			
3 Li lithium 6.9	4 Be beryllium 9.0	5 B boron 10.8	6 C carbon 12.0	7 N nitrogen 14.0	8 O oxygen 16.0	9 F fluorine 19.0	10 Ne neon 20.2	11 Na sodium 23.0	12 Mg magnesium 24.3	13 Al aluminum 27.0	14 Si silicon 28.1	15 P phosphorus 31.0	16 S sulfur 32.1	17 Cl chlorine 35.5	18 Ar argon 39.9	19 K potassium 39.1	20 Ca calcium 40.1	21 Sc scandium 45.0	22 Ti titanium 48.0	23 Cr chromium 52.0	24 Mn manganese 54.9	25 Fe iron 55.8	26 Co cobalt 58.9	27 Ni nickel 58.7	28 Cu copper 63.5	29 Zn zinc 65.4	30 Ga gallium 69.7	31 Ge germanium 72.6	32 As arsenic 74.9	33 Se selenium 79.0	34 Br bromine 79.9	35 Kr krypton 83.8	36 Xe xenon 131.3	37 Rb rubidium 85.5	38 Sr strontium 87.6	39 Y yttrium 88.9	40 Nb niobium 92.9	41 Mo molybdenum 95.9	42 Tc technetium –	43 Ru ruthenium 101.1	44 Rh rhodium 102.9	45 Pd palladium 106.4	46 Ag silver 107.9	47 Cd cadmium 112.4	48 In indium 114.8	49 Sn tin 118.7	50 Sb antimony 121.8	51 Te tellurium 127.6	52 I iodine 126.9	53 At astatine –	54 Rn radon –
55 Cs caesium 132.9	56 Ba barium 137.3	57–71 lanthanoids 137.3	72 Hf hafnium 178.5	73 Ta tantalum 180.9	74 W tungsten 183.8	75 Re rhenium 186.2	76 Os osmium 190.2	77 Ir iridium 192.2	78 Pt platinum 195.1	79 Au gold 197.0	80 Hg mercury 200.6	81 Tl thallium 204.4	82 Pb lead 207.2	83 Bi bismuth 209.0	84 Po polonium –	85 Lv Livermorium –	86 Lu Lutetium 175.0	87 Fr francium –	88 Ra radium –	89–103 actinoids 137.3	104 Rf rutherfordium –	105 Dy dubnium –	106 Sg seaborgium –	107 Bh bohrium –	108 Hs hassium –	109 Mt meitnerium –	110 Ds darmstadtium –	111 Rg roentgenium –	112 Cn copernicium –	114 Fl flerovium –	116 Lv Livermorium –	117 Mc mendelevium –	118 Nh nobelium –	119 Lr lawrencium –																	
lanthanoids		57 La lanthanum 138.9	58 Ce cerium 140.1	59 Pr praseodymium 140.9	60 Nd neodymium 144.4	61 Pm promethium –	62 Sm samarium 150.4	63 Eu europium 152.0	64 Gd gadolinium 157.3	65 Tb terbium 158.9	66 Dy dysprosium 162.5	67 Ho holmium 164.9	68 Er erbium 167.3	69 Tm thulium 168.9	70 Yb ytterbium 173.1	71 Lu lutetium 175.0	72 Cf californium –	73 Ef einsteinium –	74 Md mendelevium –	75 No nobelium –	76 Lr lawrencium –																														
actinoids		88 Ac actinium –	90 Th thorium 232.0	91 Pa protactinium 231.0	92 U uranium 238.0	93 Np neptunium –	94 Pu plutonium –	95 Am americium –	96 Cm curium –	97 Bk berkelium –	98 Cf californium –	99 Fm einsteinium –	100 Md mendelevium –	101 Md mendelevium –	102 No nobelium –	103 Lr lawrencium –	104 Lu lutetium 175.0	105 Yb ytterbium 173.1	106 Lu lutetium 175.0																																

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.