



Cambridge International AS & A Level

CANDIDATE
NAME



CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

CHEMISTRY

9701/43

Paper 4 A Level Structured Questions

October/November 2024

2 hours

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 100.
- The number of marks for each question or part question is shown in brackets [].
- The Periodic Table is printed in the question paper.
- Important values, constants and standards are printed in the question paper.

This document has **24** pages. Any blank pages are indicated.



- 1 (a) Disodium phosphate, $(\text{Na}^+)_2(\text{HPO}_4^{2-})$, reacts with an acid to form monosodium phosphate, $\text{Na}^+(\text{H}_2\text{PO}_4^-)$.

- (i) Identify the ions that are a conjugate acid–base pair in this reaction, using the formulae of the species involved.

conjugate acid

conjugate base

..... [1]

- (ii) Define buffer solution.

..... [2]

- (iii) Write **two** equations to show how a mixture of $(\text{Na}^+)_2(\text{HPO}_4^{2-})$ and $\text{Na}^+(\text{H}_2\text{PO}_4^-)$ can act as a buffer solution.

equation 1

equation 2 [2]

- (iv) Identify **one** inorganic ion that acts as a buffer in blood.

..... [1]

- (b) Compound **E** is the hydroxide of a Group 2 element. Compound **E** is a strong alkali.

2.63 g of **E** is dissolved in water to make 250 cm^3 of solution **F**. Solution **F** has a pH of 13.09 at 298 K.

- (i) Show that the concentration of hydroxide ions in solution **F** is 0.123 mol dm^{-3} .

[2]

- (ii) Explain why the concentration of compound **E** in solution **F** is $0.0615\text{ mol dm}^{-3}$.

..... [1]

- (iii) Use the concentration given in (ii) to identify compound **E**.

compound **E** [1]





- (c) Compound **E** is much more soluble than magnesium hydroxide.

A saturated solution of magnesium hydroxide in water has a concentration of $1.40 \times 10^{-4} \text{ mol dm}^{-3}$ at 298 K.

Calculate the solubility product, K_{sp} , of magnesium hydroxide. Include units.

$K_{\text{sp}} = \dots\dots\dots \text{ units } \dots\dots\dots [3]$

- (d) Explain why compound **E** is much more soluble than magnesium hydroxide.

.....

.....

.....

.....

.....

.....

.....

..... [3]

[Total: 16]



- 2 (a) Predict and explain the variation in enthalpy change of hydration for the ions F^- , Cl^- , Br^- and I^- .

.....

 [2]

- (b) Fig. 2.1 shows an incomplete energy cycle involving calcium fluoride, CaF_2 .





Fig. 2.1

- (i) Complete line D. Include state symbols. [1]
- (ii) The value of the enthalpy change for process 1 can be calculated using the values of **five** other enthalpy changes which are **not** referred to in Fig. 2.1.



Identify these **five** other enthalpy changes, using either names or symbols.

.....

 [2]





(iii) Define lattice energy, ΔH_{latt} .

.....

 [2]

(iv) Complete the expression to give the mathematical relationship between ΔH_{latt} of calcium fluoride and the enthalpy changes for processes 1 and 3.

$$\Delta H_{\text{latt}} =$$

[1]

(c) Use data from Table 2.1 to calculate a value for the hydration energy, ΔH_{hyd} , of fluoride ions, $\text{F}^-(\text{g})$.

Table 2.1

	value / kJ mol^{-1}
enthalpy change of solution of calcium fluoride, $\text{CaF}_2(\text{s})$	+13
overall enthalpy change of process 1 in Fig. 2.1	+1395
enthalpy change of formation of calcium fluoride	−1214
enthalpy change of hydration of $\text{Ca}^{2+}(\text{g})$	−1650

$$\Delta H_{\text{hyd}} \text{F}^-(\text{g}) = \dots\dots\dots \text{kJ mol}^{-1} \quad [2]$$

(d) Define entropy.

.....
 [1]





- (e) At 298 K, the Gibbs free energy change, ΔG , for the solution of compound **T** is $+6.00 \text{ kJ mol}^{-1}$.

The enthalpy change of solution, ΔH_{sol} , of compound **T** is $+30.0 \text{ kJ mol}^{-1}$ at 298 K.

Calculate the value of the entropy change, ΔS , for the solution of compound **T** at 298 K.

$$\Delta S = \dots\dots\dots \text{ J K}^{-1} \text{ mol}^{-1} \quad [2]$$

- (f) Predict whether compound **T** becomes more or less soluble as the water is heated from 298 K to 360 K. Explain your answer.

.....


..... [1]

[Total: 14]





- 3 (a) A and B react together to give product AB.



When the concentrations of A and B are both $0.0100 \text{ mol dm}^{-3}$, the rate of formation of AB is $7.62 \times 10^{-4} \text{ mol dm}^{-3} \text{ s}^{-1}$. When the concentrations of A and B are both $0.0200 \text{ mol dm}^{-3}$, the rate of formation of AB is $3.05 \times 10^{-3} \text{ mol dm}^{-3} \text{ s}^{-1}$.

- (i) Complete the **three** possible rate equations that are consistent with these data.

rate =

rate =

rate =

[2]

- (ii) Choose **one** of the rate equations you have written in (i), and calculate the value of the rate constant, k . Include the units of k .

$k = \dots$ units $\dots$ [2]

- (iii) Explain why it is **not** possible to calculate a value for the half-life, $t_{\frac{1}{2}}$, of this reaction using the value of the rate constant k calculated in (ii) and the equation $k = 0.693/t_{\frac{1}{2}}$.

.....

.....

..... [1]





(b) Catalysts may be homogeneous or heterogeneous.

- (i) Identify **two** metals that act as heterogeneous catalysts in the removal of NO_2 from the exhaust gases of car engines.

..... and [1]

- (ii) Iron acts as a heterogeneous catalyst in the Haber process.

Describe the mode of action of this iron catalyst.

.....

 [2]

- (iii) Fe^{2+} ions act as a homogeneous catalyst in the reaction between $\text{I}^-(\text{aq})$ and $\text{S}_2\text{O}_8^{2-}(\text{aq})$.

Write equations for the **two** reactions that occur when $\text{Fe}^{2+}(\text{aq})$ is added to a mixture of $\text{I}^-(\text{aq})$ and $\text{S}_2\text{O}_8^{2-}(\text{aq})$.

equation 1 $\text{S}_2\text{O}_8^{2-} + \dots\dots\dots$

equation 2 $\dots\dots\dots$ [2]

- (iv) Explain the difference between a homogeneous catalyst and a heterogeneous catalyst.

.....

 [1]






(c) Fe^{2+} ions can be oxidised to Fe^{3+} ions under alkaline conditions by suitable oxidising agents.




- (i) Iron is a transition element. Explain why iron forms stable compounds in both the +2 and the +3 oxidation states.

..... [1]

- (ii) The half-equation for the reduction of Fe^{3+} under alkaline conditions, and its $E^\ominus$ value, are shown.



Four more half-equations for reactions under alkaline conditions, and their $E^\ominus$ values, are shown.



Select **two** oxidising agents that can oxidise Fe^{2+} ions to Fe^{3+} ions under alkaline conditions.

Write an equation, and give the $E^\ominus_{\text{cell}}$ value, for **each** of the **two** reactions that occur.

oxidising agent 1:

equation:

$$E^\ominus_{\text{cell}} = \dots\dots\dots \text{V}$$

oxidising agent 2:

equation:

$$E^\ominus_{\text{cell}} = \dots\dots\dots \text{V} \quad [4]$$

[Total: 16]





4 Transition metal atoms and transition metal ions form complexes by combining with ligands.

(a) Explain why transition elements form complex ions.

.....
 [1]

(b) Co^{2+} ions form complex ion **G**.

Each **G** ion contains **two** Co^{2+} ions, both of which are octahedrally coordinated.

Each **G** ion contains one O_2 molecule, which donates one pair of electrons to **each** Co^{2+} ion, and one NH_2^- ion, which donates one pair of electrons to **each** Co^{2+} ion.

The remaining ligands are NH_3 molecules.

(i) Deduce the formula of complex ion **G**. Include its overall charge.

formula of **G** [2]

(ii) The d-orbitals of the Co^{2+} ions present in complex ion **G** are split. State the number of d-orbitals that are at a higher energy level and the number of d-orbitals that are at a lower energy level in **each** Co^{2+} ion.

number of d-orbitals at a higher energy level	
number of d-orbitals at a lower energy level	

[1]

(iii) Co^{2+} ions form a different complex ion, **M**.

Each **M** ion contains **two** Co^{2+} ions, both of which are octahedrally coordinated, but the ligands are different from the ligands in **G**.

Explain why **G** and **M** have different colours.

.....

 [2]





- (c) Cadmium forms complex ion **X**, $[\text{Cd}(\text{NH}_3)_4]^{2+}$.

When a solution containing CN^- ions is added to an aqueous solution of **X**, a ligand exchange reaction takes place, forming complex ion **Y**. **Y** contains no NH_3 ligands and no H_2O ligands.

Y is in a much higher concentration in the mixture than **X**.

The oxidation state and coordination number of cadmium do **not** change in this reaction.

- (i) Write an ionic equation for this reaction, using the formulae of the complex ions.

..... [2]

- (ii) Cadmium forms complex ion **Z** in the same oxidation state and with the same coordination number as in **X**. All the ligands in **Z** are Cl^- ions.

When $\text{NaCl}(\text{aq})$ is added to a solution of **X**, very little **Z** forms.



Write the **three** cadmium complexes, **X**, **Y** and **Z**, in order of increasing stability constant, K_{stab} .

.....
smallest value of K_{stab} largest value of K_{stab} [1]

- (d) Ethanedioate ions, $\text{C}_2\text{O}_4^{2-}$, form complexes with transition element ions.

The concentration of $\text{C}_2\text{O}_4^{2-}$ ions can be found by reaction with acidified $\text{Cr}_2\text{O}_7^{2-}$ ions. $\text{C}_2\text{O}_4^{2-}$ ions are protonated and form HOOCCOOH molecules which are oxidised by $\text{Cr}_2\text{O}_7^{2-}$.

The half-equations are shown.



- (i) Construct an equation for the reaction between acidified $\text{Cr}_2\text{O}_7^{2-}$ and HOOCCOOH .

..... [1]

- (ii) A 25.0 cm^3 sample of a solution of $\text{Na}_2\text{C}_2\text{O}_4$ reacts with exactly 16.20 cm^3 of an acidified solution of $0.0500\text{ mol dm}^{-3}$ $\text{K}_2\text{Cr}_2\text{O}_7$.

Calculate the concentration of the solution of $\text{Na}_2\text{C}_2\text{O}_4$.

$[\text{Na}_2\text{C}_2\text{O}_4] = \dots\dots\dots \text{mol dm}^{-3}$ [2]

[Total: 12]





- 5 The shapes of four different complexes, **P**, **Q**, **R** and **S**, are shown in Table 5.1.

The symbol **J** represents an atom or ion of a transition element.

The symbol **L** is used to represent a monodentate ligand.

Table 5.1

P	Q
R	S

- (a) Label **one** bond angle on **each** of complexes **P**, **Q**, **R** and **S**, and identify the size of the angle in degrees. [2]

- (b) Identify the shapes of complexes **P**, **Q**, **R** and **S**.

P

Q

R

S [2]

- (c) Two **L** ligands are exchanged with two **different** monodentate ligands **X** and **Y** in each of complexes **P**, **Q**, **R** and **S**.

Identify all the complexes which form new complexes that show geometrical isomerism.

..... [1]

- (d) Three **L** ligands are exchanged with three **different** monodentate ligands **X**, **Y** and **Z** in each of complexes **P**, **Q** and **R**.

Identify all the complexes which form new complexes that show optical isomerism.

..... [1]

[Total: 6]



* 0000800000013 *



13



BLANK PAGE


DO NOT WRITE IN THIS MARGIN






- 6 Benzene, C_6H_6 , reacts with chloroethane, C_2H_5Cl , in the presence of a suitable catalyst to form ethylbenzene, $C_6H_5C_2H_5$. In the presence of the catalyst, the ion $C_2H_5^+$ is formed. This ion reacts with benzene.

- (a) Complete the equation for the reaction of C_2H_5Cl with this catalyst to form $C_2H_5^+$ as one product.



- (b) Ethylbenzene reacts with more C_2H_5Cl , forming a mixture containing 1,2-diethylbenzene and 1,4-diethylbenzene.

- (i) Draw the structures of 1,2-diethylbenzene and 1,4-diethylbenzene.



[1]

- (ii) Explain why there is very little 1,3-diethylbenzene in the product mixture.

.....


..... [1]





(c) 1,2-diethylbenzene can be oxidised to benzene-1,2-dioic acid, $\text{C}_6\text{H}_4(\text{COOH})_2$.

benzene-1,2-dioic acid



(i) State the reagent and conditions used for this reaction.

..... [1]

(ii) Complete the overall equation for this reaction.

An atom of oxygen from the oxidising agent is represented as [O].

All of the atoms in the two ethyl groups are fully oxidised in this reaction.

..... + [O] $\rightarrow$ $\text{C}_6\text{H}_4(\text{COOH})_2$ + +
(1,2-diethylbenzene) [2]

(iii) Predict the number of peaks in the carbon-13 NMR spectrum of benzene-1,2-dioic acid.

..... [1]





- (d) The proton (^1H) NMR spectra of ethylbenzene, $\text{C}_6\text{H}_5\text{C}_2\text{H}_5$, in CDCl_3 and of benzene-1,2-dioic acid, $\text{C}_6\text{H}_4(\text{COOH})_2$, in CDCl_3 are shown. They have **not** been identified.




Fig. 6.1




Fig. 6.2

- (i) Explain the use of CDCl_3 , instead of CHCl_3 , as the solvent when obtaining these spectra.

..... [1]



(ii) Identify the substance shown by the spectrum in Fig. 6.1, and complete Table 6.1.

substance

Table 6.1

	peak at $\delta = 1.2$	peak at $\delta = 2.6$
name of splitting pattern		
group responsible for peak		
explanation of splitting pattern		

[3]

(iii) Identify the substance shown by the spectrum in Fig. 6.2, and complete Table 6.2.

substance

Table 6.2

	peak at $\delta = 7.8$	peak at $\delta = 13.1$
group responsible for peak		

[1]


(iv) When D_2O is used as a solvent, the spectrum obtained is different from the spectrum in Fig. 6.2.

Describe this difference and explain your answer.

.....

 [1]

(e) Benzene-1,2-dioic acid can be used to produce **K**.



Suggest the name of this type of reaction.

..... [1]





7 A reaction scheme is shown in Fig. 7.1.

The reagents needed for reaction 2 and reaction 3 are stated.

Reaction 5 takes place when $\text{C}_2\text{H}_5\text{NH}_2$ is mixed with compound **V**. No special conditions are required.




Fig. 7.1

(a) Identify compound **U** which contains only three elements.

..... [1]

(b) Describe the reagents and conditions for reaction 1.

..... [1]

(c) Identify compound **V**.

..... [1]

(d) Complete the equation for reaction 3.

$\text{CH}_3\text{COOH} + \text{SOCl}_2 \rightarrow$ [1]

(e) Identify compound **W**.

..... [1]

(f) Describe the conditions for reaction 4.

..... [1]





(g) Suggest the reagent needed for reaction 6.

..... [1]

(h) Complete Table 7.1 by adding the reaction numbers, 1, 2, 3, 4, 5 and 6, to the right-hand column. Use the reaction numbers given in Fig. 7.1.

Each of the numbers 1, 2, 3, 4, 5 and 6 should be used once only.

Table 7.1

type of reaction	reaction number(s)
hydrolysis	
addition	
reduction	
substitution	

[4]

(i) Compare the basicities of $\text{C}_2\text{H}_5\text{NHCOCH}_3$, $\text{C}_2\text{H}_5\text{NHC}_2\text{H}_5$ and NH_3 .

Explain your answer.

.....
 most basic least basic

 [4]

[Total: 15]





- 8 (a) An aqueous solution of phenol, $\text{C}_6\text{H}_5\text{OH}$, is acidic at 298 K.

Explain why phenol is more acidic than water.

.....

 [2]

- (b) (i) Name the **two** products formed when phenol reacts with an excess of $\text{Br}_2(\text{aq})$.

..... and [1]

- (ii) Draw the structures of the **two** isomeric organic products, with $M_r = 139$, that are formed when phenol reacts with $\text{HNO}_3(\text{aq})$ at room temperature.



[1]

- (iii) Write the equation for the reaction between phenol, $\text{C}_6\text{H}_5\text{OH}$, and sodium metal.

..... [1]

- (c) Phenol can be produced from phenylamine in a two-step synthesis.



Describe the reagents and conditions needed in each step.

step one:

reagents

conditions

step two:

reagents

conditions

[2]

[Total: 7]



* 0000800000021 *



21



BLANK PAGE

DO NOT WRITE IN THIS MARGIN





BLANK PAGE

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN





Important values, constants and standards

molar gas constant	$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
Faraday constant	$F = 9.65 \times 10^4 \text{ C mol}^{-1}$
Avogadro constant	$L = 6.022 \times 10^{23} \text{ mol}^{-1}$
electronic charge	$e = -1.60 \times 10^{-19} \text{ C}$
molar volume of gas	$V_m = 22.4 \text{ dm}^3 \text{ mol}^{-1}$ at s.t.p. (101 kPa and 273 K) $V_m = 24.0 \text{ dm}^3 \text{ mol}^{-1}$ at room conditions
ionic product of water	$K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ (at 298 K (25 °C))
specific heat capacity of water	$c = 4.18 \text{ kJ kg}^{-1} \text{ K}^{-1}$ (4.18 J g ⁻¹ K ⁻¹)





The Periodic Table of Elements

Group																					
1	2															13	14	15	16	17	18
1 H hydrogen 1.0																					
2 He helium 4.0																					
3 Li lithium 6.9																					
4 Be beryllium 9.0																					
5 B boron 10.8																					
6 C carbon 12.0																					
7 N nitrogen 14.0																					
8 O oxygen 16.0																					
9 F fluorine 19.0																					
10 Ne neon 20.2																					
11 Na sodium 23.0																					
12 Mg magnesium 24.3																					
13 Al aluminium 27.0																					
14 Si silicon 28.1																					
15 P phosphorus 31.0																					
16 S sulfur 32.1																					
17 Cl chlorine 35.5																					
18 Ar argon 39.9																					
19 K potassium 39.1																					
20 Ca calcium 40.1																					
21 Sc scandium 45.0																					
22 Ti titanium 47.9																					
23 V vanadium 50.9																					
24 Cr chromium 52.0																					
25 Mn manganese 54.9																					
26 Fe iron 55.8																					
27 Co cobalt 58.9																					
28 Ni nickel 58.7																					
29 Cu copper 63.5																					
30 Zn zinc 65.4																					
31 Ga gallium 69.7																					
32 Ge germanium 72.6																					
33 As arsenic 74.9																					
34 Se selenium 79.0																					
35 Br bromine 79.9																					
36 Kr krypton 83.8																					
37 Rb rubidium 85.5																					
38 Sr strontium 87.6																					
39 Y yttrium 88.9																					
40 Zr zirconium 91.2																					
41 Nb niobium 92.9																					
42 Mo molybdenum 95.9																					
43 Tc technetium —																					
44 Ru ruthenium 101.1																					
45 Rh rhodium 102.9																					
46 Pd palladium 106.4																					
47 Ag silver 107.9																					
48 Cd cadmium 112.4																					
49 In indium 114.8																					
50 Sn tin 118.7																					
51 Sb antimony 121.8																					
52 Te tellurium 127.6																					
53 I iodine 126.9																					
54 Xe xenon 131.3																					
55 Cs caesium 132.9																					
56 Ba barium 137.3																					
57–71 lanthanoids																					
72 Hf hafnium 178.5																					
73 Ta tantalum 180.9																					
74 W tungsten 183.8																					
75 Re rhenium 186.2																					
76 Os osmium 190.2																					
77 Ir iridium 192.2																					
78 Pt platinum 195.1																					
79 Au gold 197.0																					
80 Hg mercury 200.6																					
81 Tl thallium 204.4																					
82 Pb lead 207.2																					
83 Bi bismuth 209.0																					
84 Po polonium —																					
85 At astatine —																					
86 Rn radon —																					
87 Fr francium —																					
88 Ra radium —																					
89–103 actinoids																					
104 Rf rutherfordium —																					
105 Db dubnium —																					
106 Sg seaborgium —																					
107 Bh bohrium —																					
108 Hs hassium —																					
109 Mt meitnerium —																					
110 Ds darmstadtium —																					
111 Rg roentgenium —																					
112 Cn copernicium —																					
113 Nh nihonium —																					
114 Fl flerovium —																					
115 Mc moscovium —																					
116 Lv livermorium —																					
117 Ts tennessine —																					
118 Og oganeson —																					
119 Uue unbinilium —																					
120 Uuh unhexium —																					
121 Uus unseptium —																					
122 Uuq unquadium —																					
123 Uub unbibium —																					
124 Uut untrium —																					
125 Uuq unquadium —																					
126 Uub unbibium —																					
127 Uut untrium —																					
128 Uuq unquadium —																					
129 Uub unbibium —																					
130 Uut untrium —																					
131 Uuq unquadium —																					
132 Uub unbibium —																					
133 Uut untrium —																					
134 Uuq unquadium —																					
135 Uub unbibium —																					
136 Uut untrium —																					
137 Uuq unquadium —																					
138 Uub unbibium —																					
139 Uut untrium —																					
140 Uuq unquadium —																					
141 Uub unbibium —																					
142 Uut untrium —																					
143 Uuq unquadium —																					
144 Uub unbibium —																					
145 Uut untrium —																					
146 Uuq unquadium —																					
147 Uub unbibium —																					
148 Uut untrium —																					
149 Uuq unquadium —																					
150 Uub unbibium —																					
151 Uut untrium —																					
152 Uuq unquadium —																					
153 Uub unbibium —																					
154 Uut untrium —																					
155 Uuq unquadium —																					
156 Uub unbibium —																					
157 Uut untrium —																					
158 Uuq unquadium —																					
159 Uub unbibium —																					
160 Uut untrium —																					
161 Uuq unquadium —																					
162 Uub unbibium —																					
163 Uut untrium —																					
164 Uuq unquadium —																					
165 Uub unbibium —																					
166 Uut untrium —																					
167 Uuq unquadium —																					
168 Uub unbibium —																					
169 Uut untrium —																					
170 Uuq unquadium —																					
171 Uub unbibium —																					
172 Uut untrium —																					
173 Uuq unquadium —																					
174 Uub unbibium —																					
175 Uut untrium —																					
176 Uuq unquadium —																					
177 Uub unbibium —																					
178 Uut untrium —																					
179 Uuq unquadium —																					
180 Uub unbibium —																					
181 Uut untrium —																					
182 Uuq unquadium —																					
183 Uub unbibium —																					
184 Uut untrium —																					
185 Uuq unquadium —																					
186 Uub unbibium —																					
187 Uut untrium —																					
188 Uuq unquadium —																					
189 Uub unbibium —																					
190 Uut untrium —																					
191 Uuq unquadium —																					
192 Uub unbibium —																					
193 Uut untrium —																					
194 Uuq unquadium —																					
195 Uub unbibium —																					
196 Uut untrium —																					
197 Uuq unquadium —																					
198 Uub unbibium —																					
199 Uut untrium —																					
200 Uuq unquadium —																					
201 Uub unbibium —																					
202 Uut untrium —																					
203 Uuq unquadium —																					
204 Uub unbibium —																					
205 Uut untrium —																					
206 Uuq unquadium —																					
207 Uub unbibium —																					
208 Uut untrium —																					
209 Uuq unquadium —																					
210 Uub unbibium —																					
211 Uut untrium —																					
212 Uuq unquadium —																					
213 Uub unbibium —																					
214 Uut untrium —																					
215 Uuq unquadium —																					
216 Uub unbibium —																					
217 Uut untrium —																					
218 Uuq unquadium —																					
219 Uub unbibium —																					
220 Uut untrium —																					
221 Uuq unquadium —																					
222 Uub unbibium —																					
223 Uut untrium —																					
224 Uuq unquadium —																					
225 Uub unbibium —																					
226 Uut untrium —																					
227 Uuq unquadium —																					
228 Uub unbibium —																					
229 Uut untrium —																					
230 Uuq unquadium —																					
231 Uub unbibium —																					
232 Uut untrium —																					
233 Uuq unquadium —																					
234 Uub unbibium —																					
235 Uut untrium —																					
236 Uuq unquadium —																					
237 Uub unbibium —																					
238 Uut untrium —																					
239 Uuq unquadium —																					
240 Uub unbibium —																					
241 Uut untrium —																					
242 Uuq unquadium —																					
243 Uub unbibium —																					
244 Uut untrium —																					
245 Uuq unquadium —																					
246 Uub unbibium —																					
247 Uut untrium —																					
248 Uuq unquadium —																					
249 Uub unbibium —																					
250 Uut untrium —																					
251 Uuq unquadium —																					
252 Uub unbibium —																					
253 Uut untrium —																					
254 Uuq unquadium —																					
255 Uub unbibium —																					
256 Uut untrium —																					
257 Uuq unquadium —																					
258 Uub unbibium —																					
259 Uut untrium —																					
260 Uuq unquadium —																					
261 Uub unbibium —																					
262 Uut untrium —																					
263 Uuq unquadium —																					
264 Uub unbibium —																					
265 Uut untrium —																					
266 Uuq unquadium —																					
267 Uub unbibium —																					
268 Uut untrium —																					
269 Uuq unquadium —																					
270 Uub unbibium —																					
271 Uut untrium —																					
272 Uuq unquadium —																					
273 Uub unbibium —																					
274 Uut untrium —																					
275 Uuq unquadium —																					
276 Uub unbibium —																					
277 Uut untrium —																					
278 Uuq unquadium —																					
279 Uub unbibium —																					
280 Uut untrium —																					
281 Uuq unquadium —																					
282 Uub unbibium —																					
283 Uut untrium —																					
284 Uuq unquadium —																					
285 Uub unbibium —																					
286 Uut untrium —																					
287 Uuq unquadium —																					
288 Uub unbibium —																					
289 Uut untrium —																					
290 Uuq unquadium —																					
291 Uub unbibium —																					
292 Uut untrium —																					
293 Uuq unquadium —																					
294 Uub unbibium —																					
295 Uut untrium —																					
296 Uuq unquadium —																					
297 Uub unbibium —																					
298 Uut untrium —																					
299 Uuq unquadium —																					
300 Uub unbibium —																					
301 Uut untrium —																					