
CHEMISTRY**9701/52**

Paper 5 Planning, Analysis and Evaluation

October/November 2017

MARK SCHEME

Maximum Mark: 30

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.


Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2017 series for most Cambridge IGCSE[®], Cambridge International A and AS Level components and some Cambridge O Level components.

© IGCSE is a registered trademark.

This document consists of **7** printed pages.

Question	Answer	Marks
1(a)(i)	CuCO_3 and Cu(OH)_2 both react (with HCl) or both form copper(II) chloride	1
1(a)(ii)	(Transfer) $12.5(0) \text{ cm}^3$ of $(10.0 \text{ mol dm}^{-3}) \text{ HCl}$ using a (graduated) pipette or a burette	1
	add to a 250 cm^3 volumetric flask AND make to mark with distilled water	1
1(a)(iii)	Measure a volume of gas from the carbonate reaction or measure the (loss of) mass from the carbonate reaction	1
1(a)(iv)	Suitable apparatus for production of CO_2	1
	Suitable means of measuring CO_2 evolved	1
1(a)(v)	Correct labels on axes y-axis: volume (of gas) or mass loss or mass of 'limewater' and x-axis: time or t	1
	<p>curved line (from origin) to reach a plateau, e.g.</p> 	1

Question	Answer	Marks
1(a)(vi)	<p>Any sensible attempt seen to make the experiment accurate</p> <p>If mass loss Reduce risk of mass loss through spraying Insert cotton wool plug</p> <p>If gas collection Any method to reduce risk of gas loss Check apparatus is sealed Insert bung quickly</p> <p>Any attempt to measure temperature Check apparatus is at room temperature</p> <p>Apparatus accuracy Use an accurate or 2dp (or more) balance / gas syringe / measuring cylinder</p>	1
1(a)(vii)	<p>mol of $\text{CuCO}_3 = 0.5 \div 123.5 = 4.05 \times 10^{-3} \text{ mol}$</p>	1
	<p>moles of $\text{HCl} = 2 \times 4.05 \times 10^{-3} = 8.10 \times 10^{-3} \text{ mol}$ and volume of $\text{HCl} = 8.10 \times 10^{-3} \div 0.500 = 0.0162 \text{ dm}^3$ $= 16.2 \text{ cm}^3$</p>	1

Question	Answer	Marks
1(b)	<p>Any suitable precaution relating to stated hazard of given chemical</p> <p>For HCl Precaution (lab) gloves</p> <p>Explanation (10 mol dm⁻³) HCl is corrosive</p> <p>For CuCO₃ Precaution (lab) gloves / wash hands (after use) / face or mouth mask</p> <p>Explanation Harmful if swallowed</p>	1
1(c)(i)	$\text{moles of H}_2\text{SO}_4 = 0.40 \times \frac{24.15}{1000} = 9.66 \times 10^{-3} \text{ mol}$	1
	$\text{mass of Cu}_3(\text{CO}_3)_2(\text{OH})_2 = 344.5 \times 9.66 \times 10^{-3} \div 3 = 1.11 \text{ g}$	1
	$\% \text{ by mass} = \frac{1.11}{1.50} \times 100\% = 74.0\%$	1

Question	Answer	Marks
1(c)(ii)	<p>Problem 1 titres are not concordant / are too far apart / are 0.5(0) cm³ apart / difference is too large</p> <p>Improvement Repeat until (two) concordant titres have been achieved / two readings within 0.1(0) cm³</p> <p>Problem 2 colour change (of indicator) will be masked</p> <p>Improvement 2 Use an alternative indicator / named indicator</p> <p>[1] for each problem, [1] for an improvement</p>	3

Question	Answer				Marks																																								
2(a)(i)	<table border="1" data-bbox="349 217 994 767"> <thead> <tr> <th data-bbox="349 217 539 312">Difference in conc. D</th> <th data-bbox="546 217 692 312">$\frac{D}{m}$</th> <th data-bbox="698 217 844 312">$\log\left(\frac{D}{m}\right)$</th> <th data-bbox="851 217 994 312">$\log[X]$</th> </tr> </thead> <tbody> <tr><td data-bbox="349 317 539 360">24.04</td><td data-bbox="546 317 692 360">120.20</td><td data-bbox="698 317 844 360">2.08</td><td data-bbox="851 317 994 360">-0.02</td></tr> <tr><td data-bbox="349 365 539 408">24.31</td><td data-bbox="546 365 692 408">97.24</td><td data-bbox="698 365 844 408">1.99</td><td data-bbox="851 365 994 408">-0.16</td></tr> <tr><td data-bbox="349 413 539 456">24.40</td><td data-bbox="546 413 692 456">81.33</td><td data-bbox="698 413 844 456">1.91</td><td data-bbox="851 413 994 456">-0.22</td></tr> <tr><td data-bbox="349 461 539 504">24.59</td><td data-bbox="546 461 692 504">70.26</td><td data-bbox="698 461 844 504">1.85</td><td data-bbox="851 461 994 504">-0.39</td></tr> <tr><td data-bbox="349 509 539 552">24.67</td><td data-bbox="546 509 692 552">61.68</td><td data-bbox="698 509 844 552">1.79</td><td data-bbox="851 509 994 552">-0.48</td></tr> <tr><td data-bbox="349 557 539 600">24.73</td><td data-bbox="546 557 692 600">54.96</td><td data-bbox="698 557 844 600">1.74</td><td data-bbox="851 557 994 600">-0.57</td></tr> <tr><td data-bbox="349 604 539 647">24.77</td><td data-bbox="546 604 692 647">49.54</td><td data-bbox="698 604 844 647">1.69</td><td data-bbox="851 604 994 647">-0.64</td></tr> <tr><td data-bbox="349 652 539 695">24.80</td><td data-bbox="546 652 692 695">45.09</td><td data-bbox="698 652 844 695">1.65</td><td data-bbox="851 652 994 695">-0.70</td></tr> <tr><td data-bbox="349 700 539 743">24.83</td><td data-bbox="546 700 692 743">41.38</td><td data-bbox="698 700 844 743">1.62</td><td data-bbox="851 700 994 743">-0.77</td></tr> </tbody> </table> <p data-bbox="349 802 629 903">D data correct [1] log[X] data correct [1] All data to 2 dp [1]</p>				Difference in conc. D	$\frac{D}{m}$	$\log\left(\frac{D}{m}\right)$	$\log[X]$	24.04	120.20	2.08	-0.02	24.31	97.24	1.99	-0.16	24.40	81.33	1.91	-0.22	24.59	70.26	1.85	-0.39	24.67	61.68	1.79	-0.48	24.73	54.96	1.74	-0.57	24.77	49.54	1.69	-0.64	24.80	45.09	1.65	-0.70	24.83	41.38	1.62	-0.77	3
Difference in conc. D	$\frac{D}{m}$	$\log\left(\frac{D}{m}\right)$	$\log[X]$																																										
24.04	120.20	2.08	-0.02																																										
24.31	97.24	1.99	-0.16																																										
24.40	81.33	1.91	-0.22																																										
24.59	70.26	1.85	-0.39																																										
24.67	61.68	1.79	-0.48																																										
24.73	54.96	1.74	-0.57																																										
24.77	49.54	1.69	-0.64																																										
24.80	45.09	1.65	-0.70																																										
24.83	41.38	1.62	-0.77																																										
2(a)(ii)	greater adsorption				1																																								
	greater surface area available				1																																								
2(b)	all nine points plotted correctly				1																																								
	best-fit straight line drawn				1																																								
2(c)	Correct point (at -0.22, 1.91) identified				1																																								
	Statement explaining lack of adsorption, e.g. not enough stirring, mass of activated charcoal too low, surface area not high enough / too low / coagulation of charcoal / bulkier particles used not left long enough				1																																								

Question	Answer	Marks
2(d)(i)	co-ordinates read and recorded correctly	1
	gradient determined and same value for b	1
2(d)(ii)	intercept on y -axis read and recorded correctly	1