



Mark Scheme (Results)

October 2020

Pearson Edexcel International Advanced
Subsidiary Level
In Chemistry (WCH12)
Paper 1: Energetics, Group Chemistry,
Halogenoalkanes and Alcohols

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

October 2020

Publications Code WCH12_01_2010_MS

All the material in this publication is copyright

© Pearson Education Ltd 2020

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Using the mark scheme

Examiners should look for qualities to reward rather than faults to penalise. This does NOT mean giving credit for incorrect or inadequate answers, but it does mean allowing candidates to be rewarded for answers showing correct application of principles and knowledge. Examiners should therefore read carefully and consider every response: even if it is not what is expected it may be worthy of credit.

The mark scheme gives examiners:

- an idea of the types of response expected
- how individual marks are to be awarded
- the total mark for each question
- examples of responses that should NOT receive credit.

/ means that the responses are alternatives and either answer should receive full credit. () means that a phrase/word is not essential for the award of the mark, but helps the examiner to get the sense of the expected answer.

Phrases/words in **bold** indicate that the meaning of the phrase or the actual word is **essential** to the answer. ecf/TE/cq (error carried forward) means that a wrong answer given in an earlier part of a question is used correctly in answer to a later part of the same question.

Candidates must make their meaning clear to the examiner to gain the mark. Make sure that the answer makes sense. Do not give credit for correct words/phrases which are put together in a meaningless manner. Answers must be in the correct context.

Quality of Written Communication

Questions which involve the writing of continuous prose will expect candidates to:

- write legibly, with accurate use of spelling, grammar and punctuation in order to make the meaning clear
- select and use a form and style of writing appropriate to purpose and to complex subject matter
- organise information clearly and coherently, using specialist vocabulary when appropriate.

Full marks will be awarded if the candidate has demonstrated the above abilities. Questions where QWC is likely to be particularly important are indicated (QWC) in the mark scheme, but this does not preclude others.

Section A

Question Number	Answer	Mark
1	<p>The only correct answer is C(+121.5)</p> <p><i>A is incorrect because this is the enthalpy of atomisation of 2 mol of chlorine atoms</i></p> <p><i>B is incorrect because the enthalpy of atomisation is endothermic and refers to 2 mol of chlorine atoms</i></p> <p><i>D is incorrect because the enthalpy of atomisation is endothermic</i></p>	(1)

Question Number	Answer	Mark
2	<p>The only correct answer is A (exothermic, increases)</p> <p><i>B is incorrect because in exothermic reactions, temperatures increase</i></p> <p><i>C is incorrect because reactions with a negative enthalpy change are not endothermic</i></p> <p><i>D is incorrect because reactions with a negative enthalpy change are not endothermic and the temperature increases</i></p>	(1)

Question Number	Answer	Mark
3	<p>The only correct answer is D (water is a good solvent for ionic compounds)</p> <p><i>A is incorrect because the less open structure of water compared to ice is due to hydrogen bonding</i></p> <p><i>B is incorrect because the reduction in density of ice compared to water is due to hydrogen bonding</i></p> <p><i>C is incorrect because the expansion of volume of ice compared to water is due to hydrogen bonding</i></p>	(1)

Question Number	Answer	Mark
4	<p>The only correct answer is D (X,Z,Y)</p> <p><i>A is incorrect because Y has the highest boiling temperature and X will have a boiling temperature below that of the singly branched chain alkane Z</i></p> <p><i>B is incorrect because although X has the lowest boiling temperature, Y will have a boiling temperature above that of the singly branched chain alkane, Z</i></p> <p><i>C is incorrect because Y has the highest boiling temperature and X has the lowest boiling temperature</i></p>	(1)

Question Number	Answer	Mark
5(a)	<p>The only correct answer is B (disproportionation)</p> <p><i>A is incorrect because no species has been replaced/displaced</i></p> <p><i>C is incorrect because no species has been eliminated</i></p> <p><i>D is incorrect because water is produced and no species is hydrolysed</i></p>	(1)

Question Number	Answer	Mark
5(b)	<p>The only correct answer is B ($4.2 \times 10^{-5} \text{ mol s}^{-1}$)</p> <p><i>A is incorrect because incorrect units for the Molar Volume of a gas has been used</i></p> <p><i>C is incorrect because the stoichiometry has not been taken into account</i></p> <p><i>D is incorrect because the stoichiometry is incorrect</i></p>	(1)

Question Number	Answer	Mark
6	<p>The only correct answer is A (14,6,7)</p> <p><i>B is incorrect because the charges do not balance and the number of electrons is incorrect</i></p> <p><i>C is incorrect because the oxygen atoms do not balance and the charges do not balance</i></p> <p><i>D is incorrect because the charges do not balance and the oxygen atoms do not balance</i></p>	(1)

Question Number	Answer	Mark
7	<p>The only correct answer is D (KO_2)</p> <p><i>A is incorrect because in this oxide potassium would have an oxidation number of +1/4</i></p> <p><i>B is incorrect because in this oxide potassium would have an oxidation number of +1/2</i></p> <p><i>C is incorrect because in this oxide potassium would have an oxidation number of +3/4</i></p>	(1)
8(a)	<p>The only correct answer is B (lithium nitrate)</p> <p><i>A is incorrect because barium has a green colour in the flame test</i></p> <p><i>C is incorrect because magnesium does not have a red colour in the flame test</i></p> <p><i>D is incorrect because rubidium nitrate does not produce nitrogen dioxide when heated</i></p>	(1)

Question Number	Answer	Mark
8(b)	<p>The only correct answer is A (brown and blue to red)</p> <p><i>B is incorrect because nitrogen dioxide would have no effect on damp red litmus paper</i></p> <p><i>C is incorrect because nitrogen dioxide is brown</i></p> <p><i>D is incorrect because nitrogen dioxide is brown and would have no effect on damp red litmus paper</i></p>	(1)


Question Number	Answer	Mark
9	<p>The only correct answer is A (hydrogen fluoride only)</p> <p><i>B is incorrect because sulfuric acid is not a strong enough oxidising agent to oxidise the fluoride ion</i></p> <p><i>C is incorrect because the fluoride ion is not a strong enough reducing agent to reduce sulfuric acid</i></p> <p><i>D is incorrect because the fluoride ion is not a strong enough reducing agent to reduce sulfuric acid and sulfuric acid is not a strong enough oxidising agent to oxidise the fluoride ion</i></p>	(1)


Question Number	Answer	Mark
10(a)	<p>The only correct answer is B (v)</p> <p><i>A is incorrect because u is the activation energy for the reaction $R \rightarrow P$</i></p> <p><i>C is incorrect because w is the activation energy for the reaction $P \rightarrow R$</i></p> <p><i>D is incorrect because x is the enthalpy change for the reaction $P \rightarrow R$</i></p>	(1)

Question Number	Answer	Mark
10(b)	<p>The only correct answer is C (w)</p> <p><i>A is incorrect because u is the activation energy for the reaction $R \rightarrow P$</i></p> <p><i>B is incorrect because v is the enthalpy change for the reaction $R \rightarrow P$</i></p> <p><i>D is incorrect because x is the enthalpy change for the reaction $P \rightarrow R$</i></p>	(1)

Question Number	Answer	Mark
10(c)	<p>The only correct answer is C ($u - w = v$)</p> <p><i>A is incorrect because $v+x = 0$ and $u+w$ is the sum of the activation energies</i></p> <p><i>B is incorrect because $w-x = u$</i></p> <p><i>D is incorrect because $u-v = w$ (not x)</i></p>	(1)

Question Number	Answer	Mark
11(a)	<p>The only correct answer is B (1.20 dm^3)</p> <p><i>A is incorrect because the limiting factor of the hydrochloric acid has been neglected</i></p> <p><i>C is incorrect because the limiting factor of the hydrochloric acid has been neglected and the answer would be correct if the units were cm^3</i></p> <p><i>D is incorrect because the answer would be correct if the units were cm^3</i></p>	(1)

Question Number	Answer	Mark
11(b)	<p>The only correct answer is B</p>  <p><i>A is incorrect because the volume of gas is shown decreasing with time</i></p> <p><i>C is incorrect because the gradient is shown increasing with time</i></p> <p><i>D is incorrect because the gradient is shown as constant</i></p>	(1)

Question Number	Answer	Mark
11(c)	<p>The only correct answer is A</p>  <p><i>B is incorrect because the rate is shown increasing with time</i></p> <p><i>C is incorrect because the rate is shown as constant</i></p> <p><i>D is incorrect because the rate is shown increasing with time</i></p>	(1)

Question Number	Answer	Mark
12(a)	<p>The only correct answer is D (<i>E</i>-oct-5-en-3-ol)</p> <p><i>A is incorrect because the wrong priority has been assigned and the E/Z nomenclature is incorrect</i></p> <p><i>B is incorrect because the wrong priority has been assigned</i></p> <p><i>C is incorrect because this is the name of the Z isomer</i></p>	(1)

Question Number	Answer	Mark
12(b)	<p>The only correct answer is A ($\text{Cl}_2(\text{g})$)</p> <p><i>B is incorrect because only one chlorine atom would be added</i></p> <p><i>C is incorrect because PCl_5 does not react with a C=C bond</i></p> <p><i>D is incorrect because potassium chloride does not react with either a C=C bond or -OH group</i></p>	(1)

Question Number	Answer	Mark
12(c)	<p>The only correct answer is C ($\text{PCl}_5(\text{s})$)</p> <p><i>A is incorrect because chlorine does not react with an alcohol</i></p> <p><i>B is incorrect because hydrogen chloride does not react with an alcohol</i></p> <p><i>D is incorrect because potassium chloride does not react with either a C=C bond or -OH group</i></p>	(1)

(Total for Section A = 20 marks)

Section B

Question Number	Answer	Additional Guidance	Mark
13(a)(i)	All three arrowheads down at any point in/on the line (1) AND coefficients are 3 for CO ₂ and 4 for H ₂ O		(1)

Question Number	Answer	Additional Guidance	Mark
13(a)(ii)	<ul style="list-style-type: none"> calculation for enthalpy change on LHS (1) correct application of Hess's Law and answer with correct sign (1) 	Example of calculation; $(3 \times -393.5) + (4 \times -285.8) =$ $(-1180.5) + (-1143.2) = -2323.7(\text{kJ mol}^{-1})$ $+2219 -2323.7 = -104.7/-105 (\text{kJ mol}^{-1})$ Correct answer with sign (2)	(2)

Question Number	Answer	Additional Guidance	Mark
13(b)(i)	<p>An explanation that makes reference to the following points:</p> <ul style="list-style-type: none"> • each alkane differs from the next by $-CH_2$/one carbon atom and two hydrogen atoms/same number of C-C bonds and same number of C-H bonds/ forms a homologous series/ has a general formula C_nH_{2n+2} (1) • bonds broken and/or made increase regularly/ each $-CH_2$ combusted adds almost the same amount of energy/ bond energies are very similar (1) 	Ignore references to intermolecular forces/boiling points	(2)

Question Number	Answer	Additional Guidance	Mark
13(b)(ii)	<p>An explanation that makes reference to two of the following points:</p> <ul style="list-style-type: none"> • butane is a gas and pentane is a liquid (at 298 K / under standard conditions) (1) • liquids vaporise before burning/combustion (1) • (pentane is a liquid so) some of the energy released by combustion is used to vaporise / vaporisation is endothermic OR More energy is needed to break intermolecular forces in pentane or reverse argument (1) 	Pentane vaporises before combustion and butane is (already) a gas would score M1 and M2	(2)

Question Number	Answer	Additional Guidance	Mark
13(b)(iii)	<p>An explanation that makes reference to the following points:</p> <ul style="list-style-type: none"> • the intermolecular forces are London / dispersion forces / instantaneous dipole - induced dipole forces • these forces depend on the number of electrons which increase (as the number of carbon atoms increase) • more energy is needed to break these intermolecular forces/to separate the molecules (so the boiling temperature increases) 	<p>Any reference to hydrogen bonding or permanent dipole interaction loses M1 and M2</p> <p>(1) ALLOW: Van der Waals forces for London forces</p> <p>(1) ALLOW: increase in surface area/more points of contact (as the number of carbon atoms increase)</p> <p>(1) Any reference to breaking of covalent bonds loses M3</p>	(3)

(Total for Question 13 = 10 marks)

Question Number	Answer	Additional Guidance	Mark
14(a)	<ul style="list-style-type: none"> balanced equation state symbols correct 	<p>(1) <u>Example of equation</u> $\text{Ca(OH)}_2(\text{aq}) + \text{CO}_2(\text{g}) \rightarrow \text{CaCO}_3(\text{s}) + \text{H}_2\text{O}(\text{l})$</p> <p>(1) ALLOW 1 mark for LHS or RHS totally correct</p>	(2)

Question Number	Answer	Additional Guidance	Mark
14(b)	<ul style="list-style-type: none"> moles of hydrochloric acid in titre moles Ca(OH)_2 in 1 dm^3 mass Ca(OH)_2 in 1 dm^3 answer to 2 or 3 SF 	<p>(1) <u>Example of calculations:</u> $23.40 \times 0.0500 \div 1000 = 0.00117 / 1.17 \times 10^{-3}$ (moles)</p> <p>(1) $0.00117 \div 2 = 0.000585$ (moles) in 25 cm^3 $0.000585 \times 1000 \div 25 = 0.0234 / 2.34 \times 10^{-2}$ (moles/moles dm^{-3})</p> <p>(1) $0.0234 \times 74.1 / 74 = 1.7339 / 1.7316$ (g)</p> <p>(1) 1.7/1.73 (g dm^{-3}) to 2 or 3 SF Penalise incorrect units once only Answer to 2 or 3 SF from a calculated mass</p> <p>Correct answer with no working scores 4</p> <p>Allow TE throughout</p>	(4)

Question Number	Answer	Additional Guidance	Mark
14(c)	<ul style="list-style-type: none"> as magnesium hydroxide is less soluble/concentration of hydroxide ions would be smaller (1) titre value would be smaller (1) 	<p>Allow reverse argument Do not award magnesium hydroxide is insoluble</p> <p>M2 depends on correct M1 except when magnesium hydroxide is described as insoluble ALLOW a value less than 23.00</p>	(2)

(Total for Question 14 = 8 marks)

Question Number	Answer	Additional Guidance	Mark
15(a)	<ul style="list-style-type: none"> Silver is oxidised from 0 to +1 (1) Nitrogen is reduced from +5 to +4 (1) 	<p>These may be shown on the equation</p> <p>ALLOW one mark for silver is oxidised and nitrogen is reduced or all oxidation numbers correct ALLOW 1 or 1+ for +1</p>	(2)

Question Number	Answer	Additional Guidance	Mark
15(b)(i)	<ul style="list-style-type: none"> • calculation of energy change 	(1) Examples of calculation $Q = 50.0 \div 1000 \times 4.18 \times 5.2$ $= 1.0868 \text{ (kJ)}/1086.8\text{(J)}$	(3)
	<ul style="list-style-type: none"> • calculation of no of moles 	(1) $1.0868 \div 36.1 = 0.030105/3.0105 \times 10^{-2} \text{ (moles)}$	
	<ul style="list-style-type: none"> • calculation of concentration 	(1) $0.030105 \div 50 \times 1000 = 0.602105\text{(mol dm}^{-3}\text{)}$ $= 0.602/0.60/0.6$ $\text{(mol dm}^{-3}\text{)}$ Ignore SF	


Question Number	Answer	Additional Guidance	Mark
15(b)(ii)	<ul style="list-style-type: none"> • calculation of moles of ppt 	(1) Example of calculation $5.96 \div 187.8 = 0.0317359 \text{ (moles)}$	(2)
	<ul style="list-style-type: none"> • calculation of concentration 	(1) $0.00317359 \div 50 \times 1000 = 0.634718\text{(mol dm}^{-3}\text{)}$ Ignore SF except 1 SF ALLOW use of 188 as M_r Penalise incorrect rounding once only	

Question Number	Answer	Additional Guidance	Mark
15(b)(iii)	An explanation that makes reference to: <ul style="list-style-type: none"> <li data-bbox="421 347 1126 467">• the calorimetry method (student A) is lower because heat may be lost to the surroundings /absorbed by the container or apparatus (1) <li data-bbox="421 523 1115 627">• the gravimetric method (student B) is higher because the product may have been wet/ have impurities (1) 	Do not award: non-standard conditions, just "incomplete reaction", human error, incorrect solution density/heat capacity ALLOW: not all the solution reacted	(2)

(Total for Question 15 = 9 marks)

Question Number	Answer				Additional Guidance	Mark																				
16(a)	<table border="1" data-bbox="479 379 1496 986"> <thead> <tr> <th data-bbox="479 379 631 497">Reaction</th> <th data-bbox="631 379 981 497">Reagent</th> <th data-bbox="981 379 1272 497">Solvent</th> <th data-bbox="1272 379 1496 497">Type of reaction</th> </tr> </thead> <tbody> <tr> <td data-bbox="479 497 631 616">1</td> <td data-bbox="631 497 981 616">(potassium Hydroxide)</td> <td data-bbox="981 497 1272 616">Water/aqueous</td> <td data-bbox="1272 497 1496 616">substitution</td> </tr> <tr> <td data-bbox="479 616 631 734">2</td> <td data-bbox="631 616 981 734">(ammonia)</td> <td data-bbox="981 616 1272 734">Ethanol/C₂H₅OH</td> <td data-bbox="1272 616 1496 734">substitution</td> </tr> <tr> <td data-bbox="479 734 631 868">3</td> <td data-bbox="631 734 981 868">sodium/potassium cyanide NaCN/KCN</td> <td data-bbox="981 734 1272 868">(ethanol)</td> <td data-bbox="1272 734 1496 868">substitution</td> </tr> <tr> <td data-bbox="479 868 631 986">4</td> <td data-bbox="631 868 981 986">potassium/sodium hydroxide KOH/NaOH</td> <td data-bbox="981 868 1272 986">Ethanol/C₂H₅OH</td> <td data-bbox="1272 868 1496 986">(elimination)</td> </tr> </tbody> </table>				Reaction	Reagent	Solvent	Type of reaction	1	(potassium Hydroxide)	Water/aqueous	substitution	2	(ammonia)	Ethanol/C ₂ H ₅ OH	substitution	3	sodium/potassium cyanide NaCN/KCN	(ethanol)	substitution	4	potassium/sodium hydroxide KOH/NaOH	Ethanol/C ₂ H ₅ OH	(elimination)	<p data-bbox="1532 529 1581 564">(1)</p> <p data-bbox="1532 641 1581 676">(1)</p> <p data-bbox="1532 791 1581 826">(1)</p> <p data-bbox="1532 906 1581 941">(1)</p> <p data-bbox="1612 497 2024 711">IGNORE nucleophilic in first three reactions but penalise electrophilic/free radical once only ALLOW alcoholic for reactions 2 and 4</p> <p data-bbox="1612 759 1872 826">Do not award: cyanide/CN⁻/HCN</p> <p data-bbox="1612 906 1872 941">Do not award OH⁻</p> <p data-bbox="1612 983 2007 1123">8 correct scores 4 marks 6/7 correct scores 3 marks 4/5 correct scores 2 marks 2/3 correct scores 1 mark</p>	(4)
Reaction	Reagent	Solvent	Type of reaction																							
1	(potassium Hydroxide)	Water/aqueous	substitution																							
2	(ammonia)	Ethanol/C ₂ H ₅ OH	substitution																							
3	sodium/potassium cyanide NaCN/KCN	(ethanol)	substitution																							
4	potassium/sodium hydroxide KOH/NaOH	Ethanol/C ₂ H ₅ OH	(elimination)																							

Question Number	Answer	Additional Guidance	Mark
16(b)	butanenitrile (1)	ALLOW butane nitrile Butan(e)(1) nitrile Butanenitrile Do not award 1-nitrile butane	(1)

Question Number	Answer	Additional Guidance	Mark
16(c)	<ul style="list-style-type: none"> • Curly arrow to show attack on C-Br carbon by lone pair of electrons on the N of ammonia and dipole C-Br (1) • Curly arrow to show loss of Br⁻ (1) • Curly arrow to show loss of H⁺ from intermediate (1) 	<p>Example of mechanism:</p>  <p>Leaving groups must be shown for M2 and M3 but penalise only once Could show another molecule of ammonia or bromide ion acting as a base to remove H⁺</p>	(3)

Question Number	Answer	Additional Guidance	Mark																
16(d)	<p>This question assesses a student's ability to show a coherent and logically structured answer with linkages and fully-sustained reasoning.</p> <p>Marks are awarded for indicative content and for how the answer is structured and shows lines of reasoning.</p> <p>The following table shows how the marks should be awarded for indicative content.</p> <table border="1" data-bbox="344 678 1382 949"> <thead> <tr> <th>Number of indicative marking points seen in answer</th> <th>Number of marks awarded for indicative marking points</th> </tr> </thead> <tbody> <tr> <td>6</td> <td>4</td> </tr> <tr> <td>5-4</td> <td>3</td> </tr> <tr> <td>3-2</td> <td>2</td> </tr> <tr> <td>1</td> <td>1</td> </tr> <tr> <td>0</td> <td>0</td> </tr> </tbody> </table> <p>The following table shows how the marks should be awarded for structure and lines of reasoning.</p> <table border="1" data-bbox="338 1098 1386 1361"> <thead> <tr> <th></th> <th>Number of marks awarded for structure and sustained lines of reasoning</th> </tr> </thead> <tbody> <tr> <td>Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning demonstrated throughout.</td> <td>2</td> </tr> </tbody> </table>	Number of indicative marking points seen in answer	Number of marks awarded for indicative marking points	6	4	5-4	3	3-2	2	1	1	0	0		Number of marks awarded for structure and sustained lines of reasoning	Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning demonstrated throughout.	2	<p>Guidance on how the mark scheme should be applied.</p> <p>The mark for indicative content should be added to the mark for lines of reasoning. For example, an answer with five indicative marking points that is partially structured with some linkages and lines of reasoning, scores 4 marks (3 marks for indicative content and 1 mark for partial structure and some linkages and lines of reasoning).</p> <p>If there are no linkages between points, the same five indicative marking points would yield an overall score of 3 marks (3 marks for indicative content and no marks for linkages).</p> <p>In general, it would be expected that 5 or 6 indicative points would get 2 reasoning marks, and 3 or 4 indicative points would get 1 mark for reasoning, and 0, 1 or 2 indicative points would score zero marks for reasoning.</p>	(6)
Number of indicative marking points seen in answer	Number of marks awarded for indicative marking points																		
6	4																		
5-4	3																		
3-2	2																		
1	1																		
0	0																		
	Number of marks awarded for structure and sustained lines of reasoning																		
Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning demonstrated throughout.	2																		

Answer is partially structured with some linkages and lines of reasoning.	1	If there is any incorrect chemistry, deduct mark(s) from the reasoning. If no reasoning mark(s) awarded, do not deduct mark(s).	
Answer has no linkages between points and is unstructured.	0		<p>Ignore hydrolysis as given in the question</p> <p>Can be shown in an equation Incorrect colours of ppts loses 1RP</p> <p>Ignore length of bond in IP3 and IP4 Incorrect reason for decrease loses 1 RP. Ignore references to electronegativity.</p> <p>Consideration of intermolecular forces loses 1 RP</p> <p>IP5 could be scored by correct comparison of the reactivity/rate of reaction of any 2 bromoalkanes</p> <p>ALLOW the production of the appropriate carbocation</p>
<p>Indicative points:</p> <ul style="list-style-type: none"> • (mechanism is nucleophilic) substitution • (Precipitation/precipitate of) silver halide forms • the time taken for the precipitate to form/rate of reaction depends on the carbon - halogen bond strength • Strength/bond enthalpy of the C-Halogen bond decreases going down group 7/any correct comparison of at least two Hal-C bond strengths (because the atomic radius increases) • The reactivity/rate of reaction of the bromoalkanes increases $1^\circ < 2^\circ < 3^\circ$ • 1-bromobutane is 1°/2-bromobutane is 2° /2-bromo-2-methyl butane is 3° 			

(Total for Question 16 = 14 Marks)
(Total for Section B = 41 Marks)

Section C

Question Number	Answer	Additional Guidance	Mark
17(a)(i)	<p>An explanation that makes reference to the following points:</p> <p>EQUILIBRIUM</p> <ul style="list-style-type: none"> • there are more (gaseous) moles/molecules on LHS / less (gaseous) moles on RHS (1) • (so high pressure) moves equilibrium to RHS / increases yield of methanol (1) <p>OR</p> <p>KINETIC</p> <ul style="list-style-type: none"> • an increase in pressure increases the number/rate of collisions (of molecules) (1) • (So) increases the rate of reaction (1) 	<p>NOTE If two marks are scored they must be derived from either the kinetic or equilibrium explanations. If both explanations are used any incorrect statement negates one mark. Any reference to atoms scores 0 for M1</p> <p>If numbers of moles given, they must be correct</p> <p>M2 depends on M1 for both explanations</p>	(2)

Question Number	Answer	Additional Guidance	Mark
17(a)(ii)	<p>An explanation that makes reference to the following points:</p> <ul style="list-style-type: none"> • the (forward) reaction is exothermic (1) • a lower temperature would increase the yield /move equilibrium to RHS or reverse argument (1) • a (compromise) temperature is used to ensure the rate is fast enough (without the equilibrium position moving too far to the LHS) (1) ALLOW (increased/high/stated) temperature increases the rate of reaction 	<p>Note: if (forward) reaction is identified as endothermic then neither M1 nor M2 can be scored</p> <p>Ignore references to cost/economics/environmental</p>	(3)

Question Number	Answer	Additional Guidance	Mark
17(a)(iii)	<p>An explanation that makes reference to one of the following points:</p> <p>(a catalyst) allows a lower temperature to be used / less energy is needed (1)</p> <p>(a copper catalyst is specific for methanol) so there is less waste/fewer other products/improves atom economy (1)</p> <p>(a catalyst) is not used up/can be used again (1)</p>	Ignore just increases rate/ lowers activation energy	(1)

Question Number	Answer	Additional Guidance	Mark
17(b)(i)	<p>Reagents</p> <ul style="list-style-type: none"> sodium / potassium dichromate ((VI)) / $\text{Na}_2\text{Cr}_2\text{O}_7/\text{K}_2\text{Cr}_2\text{O}_7/\text{Cr}_2\text{O}_7^{2-}$ and acidified / H^+ / sulfuric acid (1) <p>Reaction conditions</p> <ul style="list-style-type: none"> (Heat under) reflux (1) 	<p>Do not award potassium manganate(VII)</p> <p>M2 dependent on M1 or near miss</p>	(2)

Question Number	Answer	Additional Guidance	Mark
17(b)(ii)	$\text{CH}_3\text{OH} + 2[\text{O}] \longrightarrow \text{HCOOH} + \text{H}_2\text{O} \quad (1)$	Allow; CH ₂ O ₂ , CHOOH, HCO ₂ H	(1)

Question Number	Answer	Additional Guidance	Mark
17(b)(iii)	<p>Functional group test add any carbonate /hydrogencarbonate OR add magnesium (powder/ribbon) (1) OR addition of any alcohol (and strong acid to form an ester)</p> <p>Positive result (1) effervescence / bubbles / gas given off (turns lime water cloudy OR burns with a "pop") OR Fruity smell</p>	<p>Do not award the addition of Na or PCl₅ Ignore reference to indicator</p> <p>M2 depends on M1 or near miss</p>	(2)

Question Number	Answer	Additional Guidance	Mark
17(b)(iv)	<ul style="list-style-type: none"> infra red absorption at $3750 - 3200 \text{ cm}^{-1}$ / 3300 cm^{-1} due to O-H of alcohol/methanol (1) infra red absorption at $1725 - 1700 \text{ cm}^{-1}$ / 1700 cm^{-1} due to C=O group (of carboxylic acid/methanoic acid) (1) peaks due to methanol/alcohol would disappear/be absent (1) 	<p>Allow the peak at 1020 for methanol in the fingerprint region</p> <p>Accept infra red absorption at $3300 - 2500 \text{ cm}^{-1}$ / 2500 cm^{-1} showing O-H group of COOH</p> <p>Allow the fingerprint region would change</p> <p>Penalise use of OH⁻ / -OH once only. In M1 and M2 penalise absence of bonds once only</p>	(3)

Question Number	Answer	Additional Guidance	Mark
17(c)(i)	<p>An explanation that makes reference to two of the following points:</p> <ul style="list-style-type: none"> burning (petrol/methanol/fuel) produces CO₂ /greenhouse gases/contributes to global warming (1) the methanol is made from captured CO₂ / is made from CO₂ that would have been released anyway (1) the increase in global temperatures will be reduced (1) 	<p>IGNORE references to ozone depletion</p> <p>Do not award : methane is produced</p>	(2)

Question Number	Answer	Additional Guidance	Mark
17(c)(ii)	<ul style="list-style-type: none"> 352g carbon dioxide (released per mole octane) (1) 	Example calculation: $8 \times 44 = 352 \text{ (g)}$	(1)

Question Number	Answer	Additional Guidance	Mark
17(c)(iii)	<ul style="list-style-type: none"> Mass petrol reduction per year (1) Mass CO₂ reduction (1) OR <ul style="list-style-type: none"> Moles of petrol used per year (1) 5% reduction in CO₂ (1) OR <ul style="list-style-type: none"> 5% reduction in moles of petrol used per year (1) Mass CO₂ produced (1) 	Example of calculation: $1200 \times 0.05 = 60 \text{ (kg)}$ $60 \div 114 \times 352 = 185.26 \text{ kg}$ $(1200 \times 1000) \div 114 = 10526 \text{ (moles)}$ $352 \times 10526 \times 5 \div 100 = 185260 \text{ g}$ $= 185.26 \text{ kg}$ $1200 \times 1000 \times 0.05 \div 114 = 526.3 \text{ (moles)}$ $526.3 \times 352 \div 1000 = 185.26 \text{ kg}$ IGNORE SF except 1 SF correct answer with no working scores 2	(2)

(Total for Section C = 19 Marks)

Total for paper = 80 Marks

