

Please check the examination details below before entering your candidate information

Candidate surname					Other names			
Pearson Edexcel		Centre Number			Candidate Number			
International		[][][][][]			[][][][][]			
Advanced Level								
Wednesday 20 January 2021								
Morning (Time: 1 hour 20 minutes)					Paper Reference WCH13/01			
Chemistry								
International Advanced Subsidiary/Advanced Level								
Unit 3: Practical Skills in Chemistry I								
You must have: Scientific calculator, ruler							Total Marks	

Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
– *there may be more space than you need.*
- Show all your working in calculations and include units where appropriate.

Information

- The total mark for this paper is 50.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*
- You will be assessed on your ability to organise and present information, ideas, descriptions and arguments clearly and logically, including your use of grammar, punctuation and spelling.
- A Periodic Table is printed on the back cover of this paper.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P67747A

©2021 Pearson Education Ltd.

1/1/1



Pearson

Answer ALL questions. Write your answers in the spaces provided.

- 1 (a) A student was provided with five test tubes labelled **A**, **B**, **C**, **D** and **E**, each containing a colourless aqueous solution.

The five solutions were known to be

barium chloride

nitric acid

potassium bromide

silver nitrate

sodium carbonate

The student carried out a series of tests to identify which test tube contained which solution.

- (i) The student tested each solution using universal indicator paper. Only solution **A** turned the paper red.

Identify solution **A**.

(1)

- (ii) The student mixed 1 cm^3 of solution **A** separately with 1 cm^3 of each of the other solutions.

There was no change for three of the mixtures but effervescence was observed when solution **A** was added to solution **C**.

Identify solution **C**.

(1)

- (iii) Write an **ionic** equation for the reaction between solution **A** and solution **C**. Include state symbols.

(2)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



(iv) The student then mixed 1 cm^3 samples of the remaining solutions as shown in **Table 1**.

Solutions mixed	Observation
B and D	no change
B and E	cream precipitate
D and E	white precipitate

Table 1

Identify the three remaining solutions.

(3)

Solution **B**

Solution **D**

Solution **E**

(b) Three of the cations in the compounds in (a) can be identified using flame tests.

Complete **Table 2**.

(3)

Cation formula	Flame colour

Table 2

(Total for Question 1 = 10 marks)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



2 Sodium hydroxide solution reacts with carbon dioxide in the air and should be standardised before use. Ethanedioic acid may be used for this standardisation.

(a) A standard solution of ethanedioic acid, $(\text{COOH})_2$, is prepared.

- 2.40 g of solid ethanedioic acid is dissolved in approximately 100 cm^3 of deionised water in a beaker.
 - The solution is transferred into a 250.0 cm^3 volumetric flask and made up to the mark with deionised water.
- (i) Give a possible reason why any solution remaining in the beaker is washed into the volumetric flask before making up to the mark.

(1)

(ii) Calculate the concentration of this standard solution of ethanedioic acid in mol dm^{-3} .

Give your answer to an appropriate number of significant figures.

[Molar mass of ethanedioic acid = 90.0 g mol^{-1}]

(2)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



- (b) A **different** standard solution of ethanedioic acid is used to determine the concentration of a sodium hydroxide solution **J**.

Procedure

Step 1 A burette is rinsed with deionised water.

Step 2 The burette is then rinsed with $0.0900 \text{ mol dm}^{-3}$ ethanedioic acid and filled with this acid solution.

Step 3 A pipette is used to transfer 25.0 cm^3 portions of solution **J** to conical flasks.

Step 4 The portions are titrated with the ethanedioic acid solution using phenolphthalein indicator.


- (i) Explain why the burette is rinsed with ethanedioic acid solution in Step 2.

(1)

- (ii) The diagram shows how the student read the filled pipette in Step 3.

Identify the **two** mistakes the student made.

(2)



(iii) The student completely emptied the pipette for each transfer in Step 3.

Explain the effect **on the titre** of completely emptying the pipette rather than leaving a small amount of solution in the tip.

(2)

.....

.....

.....

.....

(iv) State the colour **change** in the conical flask at the end-point.

(2)

From to

(c) The titration results are shown.

Titration	1	2	3
Final reading / cm ³	25.05	26.60	25.50
Initial reading / cm ³	0.00	2.00	1.00
Titre / cm ³			
Titres used in calculation of mean			

(i) Complete the table and calculate the mean titre.

(2)

DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



(ii) Calculate the concentration of the sodium hydroxide solution in mol dm^{-3} .

The equation for the titration is



(3)

(Total for Question 2 = 15 marks)

DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



3 This question is about the thermal decomposition of Group 2 carbonates.

A student heated a sample of a Group 2 carbonate until no more gas was produced. The equation for the decomposition is



(a) Give a reason why the delivery tube must be removed from the water bath before removing the test tube from the heat source.

(1)

(b) The results of the experiment are shown.

Measurement	Value
Volume of carbon dioxide / cm ³	95
Mass of test tube + carbonate / g	21.69
Mass of test tube / g	21.36
Mass of carbonate / g	0.33



- (i) Using the results of the experiment identify the Group 2 metal.

[Molar volume of gas at room temperature and pressure = $24.0 \text{ dm}^3 \text{ mol}^{-1}$]

(3)

- (ii) The student suggested that the experiment could be made more accurate by increasing the mass of carbonate from 0.33 g to 1.00 g.

No changes to the size of the apparatus or the method of measurement of the gas produced would be made.

Comment on this suggestion.

(2)

.....

.....

.....

.....

.....

.....

.....


.....

.....



- (c) The enthalpy change for the thermal decomposition of a carbonate, $\Delta_r H$, is difficult to measure directly.

An example of a Hess's Law cycle to determine it indirectly is



In an experiment to determine ΔH_1 , 0.050 mol of MCO_3 was placed in a 100 cm^3 beaker. 60 cm^3 of 2 mol dm^{-3} hydrochloric acid (an excess) was added and the mixture stirred. The maximum temperature rise measured was 6.0°C .

[Heat capacity of solution produced = $4.18 \text{ J g}^{-1} \text{ }^\circ\text{C}^{-1}$

Density of solution = 1.0 g cm^{-3}]

- (i) Calculate the enthalpy change, ΔH_1 , for the reaction between MCO_3 and hydrochloric acid in kJ mol^{-1} . Include a sign with your answer.

(2)

- (ii) Using your answer to (c)(i), calculate the enthalpy change, $\Delta_r H$, for the thermal decomposition of this Group 2 carbonate in kJ mol^{-1} . Include a sign with your answer.

(1)

(Total for Question 3 = 9 marks)



- 4 The halogenoalkane 2-chloro-2-methylpropane may be prepared from 2-methylpropan-2-ol.

Procedure

- Step 1** Add 35 cm³ of concentrated hydrochloric acid to 8.00 g of 2-methylpropan-2-ol in a conical flask. Swirl the mixture gently for 20 minutes.
- Step 2** Two distinct layers form. The upper (organic) layer contains the required product. The lower aqueous layer is removed using a separating funnel.
- Step 3** Add a solution of sodium hydrogencarbonate to the organic layer. Swirl gently. Stopper the separating funnel and shake it. Invert the separating funnel and open the tap.
- Step 4** Return the separating funnel to its upright position, remove the stopper and run off the aqueous layer. Transfer the organic layer into a clean conical flask.
- Step 5** Add some anhydrous sodium sulfate. Leave the flask to stand and decant off the liquid.
- Step 6** Distil the liquid, collecting the product between 50°C and 52°C.
- (a) (i) The concentrated hydrochloric acid used in Step 1 was labelled



Suggest **two** safety precautions, other than wearing safety spectacles and a laboratory coat, to minimise the risk when using this reagent in Step 1.

(2)

- (ii) Explain why the product in the organic layer in Step 2 does not mix with the aqueous layer.

(2)



(iii) State why the tap of the separating funnel must be opened in Step 3.

(1)

(iv) State why anhydrous sodium sulfate is added to the organic layer in Step 5.

(1)

(v) Draw the apparatus required to distil the product and collect the distillate between 50°C and 52°C in Step 6.

(4)


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



(b) The equation for the reaction is



The final product after distillation weighed 2.62 g.

Calculate the percentage yield.

(3)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 6 7 7 4 7 A 0 1 3 1 6

- (c) The haloalkane produced is used in an experiment to compare its rate of hydrolysis with two other haloalkanes.

A student dissolves separate 1.0 cm^3 samples of each haloalkane in ethanol and adds 2 cm^3 of silver nitrate solution.

The time taken for a precipitate to form is recorded. The results are shown.

Haloalkane	Time / s
2-chloro-2-methylpropane	5
1-chloro-2-methylpropane	320
1-bromo-2-methylpropane	140

The student concludes that both the structure of the haloalkane and the identity of the halogen affect the rate of hydrolysis.

Explain how the results support this conclusion.

(3)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(Total for Question 4 = 16 marks)

TOTAL FOR PAPER = 50 MARKS



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE



The Periodic Table of Elements

1 2 3 4 5 6 7 0 (8) (18)

1.0
H
hydrogen
1

Key

relative atomic mass
atomic symbol
name
atomic (proton) number

(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)																																																																																											
6.9 Li lithium 3	9.0 Be beryllium 4	23.0 Na sodium 11	24.3 Mg magnesium 12	39.1 K potassium 19	40.1 Ca calcium 20	85.5 Rb rubidium 37	87.6 Sr strontium 38	132.9 Cs caesium 55	[223] Fr francium 87	45.0 Sc scandium 21	47.9 Ti titanium 22	91.2 Zr zirconium 40	91.2 Y yttrium 39	138.9 Ba barium 56	[226] Ra radium 88	47.9 Ti titanium 22	91.2 Zr zirconium 40	91.2 Y yttrium 39	138.9 Ba barium 56	[226] Ra radium 88	50.9 V vanadium 23	50.9 Nb niobium 41	180.9 Ta tantalum 73	180.9 La* lanthanum 57	[227] Ac* actinium 89	50.9 V vanadium 23	50.9 Nb niobium 41	180.9 Ta tantalum 73	180.9 La* lanthanum 57	[227] Ac* actinium 89	54.9 Mn manganese 25	[98] Tc technetium 43	186.2 Re rhenium 75	186.2 W tungsten 74	54.9 Mn manganese 25	[98] Tc technetium 43	186.2 Re rhenium 75	186.2 W tungsten 74	55.8 Fe iron 26	101.1 Ru ruthenium 44	101.1 Rh rhodium 45	192.2 Os osmium 76	192.2 Ir iridium 77	192.2 Pt platinum 78	197.0 Au gold 79	197.0 Hg mercury 80	55.8 Fe iron 26	101.1 Ru ruthenium 44	101.1 Rh rhodium 45	192.2 Os osmium 76	192.2 Ir iridium 77	192.2 Pt platinum 78	197.0 Au gold 79	197.0 Hg mercury 80	58.7 Ni nickel 28	106.4 Pd palladium 46	195.1 Pt platinum 78	195.1 Ds darmstadtium 110	58.7 Ni nickel 28	106.4 Pd palladium 46	195.1 Pt platinum 78	195.1 Ds darmstadtium 110	63.5 Cu copper 29	107.9 Ag silver 47	197.0 Au gold 79	197.0 Rg roentgenium 111	63.5 Cu copper 29	107.9 Ag silver 47	197.0 Au gold 79	197.0 Rg roentgenium 111	65.4 Zn zinc 30	112.4 Cd cadmium 48	200.6 Hg mercury 80	65.4 Zn zinc 30	112.4 Cd cadmium 48	200.6 Hg mercury 80	69.7 Ga gallium 31	72.6 Ge germanium 32	118.7 Sn tin 50	118.7 Pb lead 82	69.7 Ga gallium 31	72.6 Ge germanium 32	118.7 Sn tin 50	118.7 Pb lead 82	74.9 As arsenic 33	74.9 Sb antimony 51	209.0 Bi bismuth 83	74.9 As arsenic 33	74.9 Sb antimony 51	209.0 Bi bismuth 83	79.0 Br bromine 35	79.0 Se selenium 34	127.6 Te tellurium 52	127.6 Po polonium 84	79.0 Br bromine 35	79.0 Se selenium 34	127.6 Te tellurium 52	127.6 Po polonium 84	83.8 Kr krypton 36	83.8 Kr krypton 36	131.3 Xe xenon 54	131.3 Xe xenon 54	83.8 Kr krypton 36	83.8 Kr krypton 36	131.3 Xe xenon 54	131.3 Xe xenon 54	131.3 Xe xenon 54	131.3 Xe xenon 54

Elements with atomic numbers 112-116 have been reported but not fully authenticated

140 Ce cerium 58	141 Pr praseodymium 59	144 Nd neodymium 60	147 Pm promethium 61	150 Sm samarium 62	152 Eu europium 63	157 Gd gadolinium 64	159 Tb terbium 65	163 Dy dysprosium 66	165 Ho holmium 67	167 Er erbium 68	169 Tm thulium 69	173 Yb ytterbium 70	175 Lu lutetium 71
232 Th thorium 90	231 Pa protactinium 91	238 U uranium 92	237 Np neptunium 93	242 Pu plutonium 94	243 Am americium 95	247 Cm curium 96	245 Bk berkelium 97	251 Cf californium 98	254 Es einsteinium 99	253 Fm fermium 100	256 Md mendelevium 101	254 No nobelium 102	257 Lr lawrencium 103

* Lanthanide series
* Actinide series

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

