Please check the examination deta	ails below	before enterir	g your candidate information
Candidate surname		(Other names
Pearson Edexcel International Advanced Level	Centre	Number	Candidate Number
Tuesday 5 May 2020			
Afternoon (Time: 1 hour 20 minu	ıtes)	Paper Ref	erence WCH13/01
Chemistry			
International Advanced Unit 3: Practical Skills in		•	dvanced Level
Candidates must have: Scienti Ruler	ific calc	ulator	Total Marks

Instructions

- Use **black** ink or **black** ball-point pen.
- Fill in the boxes at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
 - there may be more space than you need.

Information

- The total mark for this paper is 50.
- The marks for **each** question are shown in brackets
 - use this as a guide as to how much time to spend on each question.
- You will be assessed on your ability to organise and present information, ideas, descriptions and arguments clearly and logically, including your use of grammar, punctuation and spelling.
- There is a Periodic Table on the back cover of this paper.

Advice

- Read each question carefully before you start to answer it.
- Show all your working in calculations and include units where appropriate.
- Check your answers if you have time at the end.

Turn over ▶







	Answer ALL the questions.	
	Write your answers in the spaces provided.	
1	A white anhydrous crystalline solid A contains one cation and one anion.	
	Solid A was heated in a test tube and the following observations were made.	
	A brown gas was produced.	
	A glowing splint relit when placed in the mouth of the test tube.	
	A white solid remained in the test tube.	
	(a) Identify, by name or formula, the two gases formed.	(2)
	(b) Identify, by name or formula, the anion present in A .	(1)
	(c) A flame test was carried out on A and a green colour was observed.	
	Identify, by name or formula, the cation present in A .	(1)
	(d) Give the formula of solid A and the formula of the white solid formed on hear	ting.

(2)

Solid A

White solid



(e) About 5 cm³ of an aqueous solution of **A** was placed in each of two test tubes.

Five drops of aqueous sodium hydroxide were added to one of the test tubes and five drops of dilute sulfuric acid were added to the other.

In the table give the observations you would expect to make.

(2)

Addition of sodium hydroxide solution	Addition of dilute sulfuric acid solution

(Total for Question 1 = 8 marks)

2 (a) A student was provided with aqueous solutions of four compounds:

hydrochloric acid potassium carbonate silver nitrate sodium chloride

Four bottles, labelled **B**, **C**, **D** and **E**, each contained one of the solutions. The student mixed pairs of the solutions to determine which was in each bottle.

The results are shown.

Solutions mixed	Observations
B and C	A white precipitate formed which did not dissolve on the addition of dilute nitric acid
B and D	A precipitate formed which dissolved with effervescence on the addition of dilute nitric acid
B and E	A white precipitate formed which did not dissolve on the addition of dilute nitric acid
C and D	Effervescence with bubbles of a colourless gas given off
C and E	No change
D and E	No change

Using the observations in the table, deduce the identity of the compound in each bottle.

В		
L		
D)	
_		



- (b) To identify the cations in sodium chloride and potassium carbonate, a student carried out flame tests using the following method.
 - Step 1 A sample of solid sodium chloride was placed on a watch glass and a few drops of concentrated nitric acid were added. The solid and acid were mixed to form a paste.
 - Step **2** A length of copper wire was dipped into the paste.
 - Step **3** A Bunsen burner was set up with the air-hole closed. The copper wire containing the paste was placed into the Bunsen burner flame and the colour observed.
 - Step 4 The procedure was repeated using solid potassium carbonate.

For each of the Steps 1, 2 and 3 give an improvement in the procedure explaining why the change is necessary.

(6)

Step	Improvement	Explanation
1		
'		
2		
3		

(Total for Question 2 = 9 marks)



- This question is about three organic liquids, **F**, **G** and **H**.
 - (a) Tests were carried out on **F** and **G**.

Each liquid contained one functional group.

Test 1

A spatula measure of phosphorus(V) chloride, PCI₅, was added to about 1 cm³ of each liquid in separate test tubes.

Any gas evolved was tested with damp blue litmus paper.

F	G
Steamy fumes were given off. Damp blue litmus paper turned red	Steamy fumes were given off. Damp blue litmus paper turned red

(i) Identify, by name or formula, the steamy fumes produced in Test 1.

(1)

Test 2

About 1 cm³ of sodium hydrogencarbonate solution was added to 1 cm³ of each liquid in separate test tubes.

F	G
No reaction	A colourless gas was given off that turned limewater cloudy

(ii) Identify, by name or formula, the gas produced in Test 2.

(iii) Using the results from Tests 1 and 2 and the information at the start of the question, **name** the functional groups present in **F** and **G**.

(2)

Functional group in F	Functional group in G

(iv) **F** and **G** both have a molar mass of 46 g mol⁻¹.

Draw the **displayed** formula of **F** and **G**.

(2)

F	G

(v) State whether or not it is possible to distinguish between **F** and **G** using infrared spectra. Justify your answer.

Wavenumber values are not required.



(b) The organic liquid **H** is a pheromone thought to be involved in communication between rabbits.


State the initial and final appearance of each mixture when the tests described were carried out on liquid ${\bf H}$.

(4)

Tests	Observations
A few drops of H were shaken with bromine water.	
In a test tube, a few drops of H were added to 1 cm ³ of Benedict's or Fehling's solution.	
The mixture was warmed in a water bath.	

(Total for Question 3 = 11 marks)

4 The enthalpy change of neutralisation of hydrochloric acid may be determined using the apparatus shown.



The equation for the reaction is


$$HCI(aq) + NaOH(aq) \rightarrow NaCI(aq) + H_2O(I)$$

Procedure

- Step 1 Place 25.0 cm³ of 1.00 mol dm⁻³ hydrochloric acid in a polystyrene cup. Record the temperature of the hydrochloric acid.
- Step 2 Record the temperature of 30.0 cm³ of 1.00 mol dm⁻³ sodium hydroxide.
- Step **3** Add the sodium hydroxide to the hydrochloric acid in the polystyrene cup. Stir the mixture and record the maximum temperature reached.
- (a) (i) Give a reason why an excess of sodium hydroxide was used.



(ii) The diagram shows part of the thermometer when the temperature had reached its maximum.



Record the temperature in the table of results and then complete the table by giving the temperature change.

(1)

Results

Measurement	Temperature /°C
Temperature of 25 cm ³ hydrochloric acid	21.5
Temperature of 30 cm ³ sodium hydroxide	21.5
Mean starting temperature	21.5
Maximum temperature of the mixture	
Temperature change	

(b) Calculate the enthalpy change of neutralisation of hydrochloric acid.

Include a sign and units in your answer.

[Assume: the density of both solutions and the mixture = 1.0 g cm⁻³ the specific heat capacity of the mixture = $4.2 \text{ J g}^{-1} \,^{\circ}\text{C}^{-1}$]

(4)

(c) The experiment was repeated using a glass beaker instead of a polystyrene cup.

Explain how the value obtained for the enthalpy change of neutralisation would be different.

(2)

(Total for Question 4 = 8 marks)

5 A student carried out an experiment to identify the metal M in the hydrated carbonate M_2CO_3 .10 H_2O .

A solution was made by dissolving 3.56 g of the hydrated metal carbonate in distilled water and making the volume up to 250.0 cm³ in a volumetric flask.

25.0 cm³ of this solution was placed in a conical flask and titrated with 0.100 mol dm⁻³ of hydrochloric acid.

The equation for the reaction is

$$M_2CO_3(aq) + 2HCI(aq) \rightarrow 2MCI(aq) + H_2O(I) + CO_2(g)$$

(a) Name a suitable piece of apparatus to measure the 25.0 cm³ of solution.

(1)

(b) Methyl orange indicator was used in this titration.

Give the colour change in the conical flask at the end-point.

(2)

Colour change from to to

(c) The results of the titration are shown.

Number of titration	1	2	3
Burette reading (final) / cm ³	25.25	26.00	24.85
Burette reading (initial) / cm ³	0.00	1.00	0.05
Titre / cm ³			

(i) Complete the table.



(ii) Using appropriate titrations, calculate the mean titre.

(1)

(iii) Using your answer to (c)(ii), calculate the number of moles of HCl in the mean titre.

(1)

(iv) Calculate the number of moles of M_2CO_3 in 25.0 cm³ of the solution. Hence calculate the number of moles of M_2CO_3 in the 250.0 cm³ volumetric flask.

(2)

(v) Using your answer in (c)(iv) and the mass of M₂CO₃.10H₂O in the 250 cm³ of solution, calculate the molar mass of M₂CO₃.10H₂O.

(1)

(vi) Use your answer to (c)(v) to identify metal M.

(2)



(d) The titration was repeated without using an indicator.	
Describe how you would obtain large, dry crystals of the metal chloride, MCI, from this titration solution. Diagrams are not required.	
	(3)
(Total for Question 5 = 14 r	marks)

TOTAL FOR PAPER = 50 MARKS

BLANK PAGE



The Periodic Table of Elements

				4 7	
ted	[222] Rn radon 86	Xe xenon 54	83.8 Kr krypton 36	He hetium 2 2 20.2 Ne neon 10 39.9 Ar argon 18	0 (8)
een repor	[210] At astatine 85	126.9 I todine 53	79.9 Br bromine 35	(17) 19.0 F Fluorine 9 9 9 35.5 CL chlorine	1
16 have b	[209] Po potonium 84	127.6 Te tellurium 52	79.0 Se selenium 34	(16) 16.0 0 00 0 000 8 8 8 8 8 32.1 S sulfur 16	9
tomic numbers 112-116 hav but not fully authenticated	209.0 Bi bismuth 83	121.8 Sb antimony 51	74.9 As arsenic 33	(15) 14.0 N introgen 7 7 7 7 7 31.0 P phosphorus 15	2
stomic num but not fu	207.2 Pb tead 82	118.7 Sn tin 50	72.6 Ge germanium 32	(14) 12.0 C carbon 6 6 Si Si silicon p	4
Elements with atomic numbers 112-116 have been reported but not fully authenticated	204.4 Tl thallium 81	114.8 In indium 49	69.7 Ga gattium 31	(13) 10.8 B boron 5 27.0 Al atuminium 13	m
Elem	200.6 Hg mercury 80	112.4 Cd cadmium 48	65.4 Zn zinc 30	(12)	
[272] Rg roentgenium	197.0 Au gold 79	Ag silver 47	63.5 Cu copper 29	aa	
Ds damstadtum r	195.1 Pt platinum 78	106.4 Pd patladium 46	58.7 Ní nickel 28	(01)	
Mt meitnerium of	192.2 Ir iridium 77	Rh rhodium 45	Co cobalt 27	(6)	
Hs Hassium n	190.2 Os osmium 76	101.1 Ru ruthenium 44	55.8 Fe iron 26	1.0 T 1 1	
[264] Bh bohrium 107	Re rhenium 75		54.9 Mn manganese 25	6	
[266] Sg seaborgium 106	183.8 W tungsten 74	95.9 [98] Mo Tc molybdenum technetium 42 43	52.0 Cr chromium r 24	nass ool umber (6)	
[262] Db dubnium s	180.9 Ta tantalum 73	92.9 Nb niobium 41	50.9 V vanadium 23	relative atomic mass atomic symbol name atomic (proton) number (4) (5) (6)	
[261] Rf nutherfordium	178.5 Hf hafnium 72	91.2 Zr zirconium 40	47.9 Th titanium 22	relativ ator	
[227] Ac* actinium r	138.9 La* lanthanum 57	88.9 × yttrium 39	45.0 Sc scandium 21	(3)	
[226] Ra radium 88	137.3 Ba barium 56	87.6 Sr strontium 38	Ca calcium 20	(2) 9.0 Be beryllium 4 24.3 Mg magnesium	7
[223] Fr francium 87	132.9 Cs caesium 55	85.5 Rb rubidium 3	39.1 K potassium 19	(1) 6.9 Li 13 13 3 3 23.0 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	-

* Lanthanide series

* Actinide series

Cerium cerium 58	141 Pr praseodymium 59	144 Nd neodymium 60	[147] Pm promethium 61	150 Sm samarium 62	152 Eu europium 63	157 Gd gadotinium 64	159 Tb terbium 65	163 Dy dysprosium 66	165 Ho hotmium 67	167 Er erbium 68	169 Tm thuflum 69	173 Yb ytterbium 70	175 Lu Lutetiun 71
232 Th thorium p	[231] Pa protactinium 91	238 U uranium 92	[237] Np neptunium 93	[242] Pu plutonium 94	[243] Am americium 95	(247) Cm curlum 96	[245] BK berkelium 97	Cf Cf californium 98	[254] Es einsteinlum 99	[253] Fm fermium 100	Md Mamerdelevlum 101	No nobettum 102	[257] Lr lawrenctum