



# Mark Scheme (Results)

October 2021

Pearson Edexcel International Advanced Level  
In Chemistry (WCH15)  
Paper 01: Transition Metals and Organic  
Nitrogen Chemistry

## Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at [www.edexcel.com](http://www.edexcel.com) or [www.btec.co.uk](http://www.btec.co.uk). Alternatively, you can get in touch with us using the details on our contact us page at [www.edexcel.com/contactus](http://www.edexcel.com/contactus).

## Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

October 2021

Question Paper Log Number P67131A

Publications Code WCH15\_01\_2110\_MS

All the material in this publication is copyright

© Pearson Education Ltd 2021

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

## Section A (Multiple Choice)

Question number	Answer	Mark
1(a)	<p><b>The only correct answer is D</b> (Pt, Pt)</p> <p><i>A is incorrect because both electrodes should be made of platinum</i></p> <p><i>B is incorrect because both electrodes should be made of platinum</i></p> <p><i>C is incorrect because both electrodes should be made of platinum</i></p>	<b>1</b>

Question number	Answer	Mark
1(b)	<p><b>The only correct answer is C</b> (<math>1.00 \text{ mol dm}^{-3} \text{ HCl(aq)}</math>)</p> <p><i>A is incorrect because <math>\text{H}_3\text{PO}_4</math> is not completely ionised</i></p> <p><i>B is incorrect because <math>\text{H}_2\text{SO}_4</math> is not completely ionised</i></p> <p><i>D is incorrect because <math>\text{CH}_3\text{COOH}</math> is not completely ionised</i></p>	<b>1</b>

Question number	Answer	Mark
1(c)	<p><b>The only correct answer is B</b> (17.91 g)</p> <p><i>A is incorrect because there should be only one mol of chromium ions per mol of dichromate ions</i></p> <p><i>C is incorrect because there should be only one mol of chromium ions per mol of dichromate ions</i></p> <p><i>D is incorrect because there should be only one mol of chromium ions per mol of dichromate ions</i></p>	<b>1</b>

Question number	Answer	Mark
1(d)	<p><b>The only correct answer is A</b> (<math>\text{H}_2\text{SO}_4</math>)</p> <p><i>B is incorrect because chloride ions could be oxidised</i></p> <p><i>C is incorrect because bromide ions would be oxidised</i></p> <p><i>D is incorrect because this would introduce additional chromium species into the mixture</i></p>	1
2	<p><b>The only correct answer is C</b> (<math>\text{Pt} \mid \text{Fe}^{2+}, \text{Fe}^{3+} \parallel [\text{MnO}_4^- + 8\text{H}^+], [\text{Mn}^{2+} + 4\text{H}_2\text{O}] \mid \text{Pt}</math>)</p> <p><i>A is incorrect because both electrodes should be made of platinum</i></p> <p><i>B is incorrect because both electrodes should be made of platinum and the <math>\text{MnO}_4^- / \text{Mn}^{2+}</math> half-cell does not show reduction</i></p> <p><i>D is incorrect because the <math>\text{MnO}_4^- / \text{Mn}^{2+}</math> half-cell does not show reduction</i></p>	1
3	<p><b>The only correct answer is D</b> (<math>2\text{Ag}^{2+} \rightarrow \text{Ag}^+ + \text{Ag}^{3+}</math>)</p> <p><i>A is incorrect because the disproportionation is not thermodynamically feasible</i></p> <p><i>B is incorrect because the disproportionation is not thermodynamically feasible</i></p> <p><i>C is incorrect because the disproportionation is not thermodynamically feasible</i></p>	1

Question number	Answer	Mark
4	<p><b>The only correct answer is A</b> (the cathode has a more positive potential than the anode)</p> <p><i>B is incorrect because oxidation always occurs at the anode</i></p> <p><i>C is incorrect because oxygen is reduced at the positive electrode</i></p> <p><i>D is incorrect because the overall reaction is the same under both acidic and alkaline conditions</i></p>	1

Question number	Answer	Mark
5	<p><b>The only correct answer is B</b> (carbon monoxide forms stronger dative covalent bonds with haemoglobin than does oxygen)</p> <p><i>A is incorrect because carbon monoxide can be displaced from carboxyhaemoglobin</i></p> <p><i>C is incorrect because the formation of carboxyhaemoglobin does not lead to an increase in the entropy of the system</i></p> <p><i>D is incorrect because the difference in bond type does not fully explain the difference in dative covalent bond strength</i></p>	1


Question number	Answer	Mark
6	<p><b>The only correct answer is D</b> (pink solution → blue precipitate → yellow-brown solution)</p> <p><i>A is incorrect because <math>\text{CoCl}_2(\text{aq})</math> is a pink solution</i></p> <p><i>B is incorrect because the blue precipitate dissolves in excess aqueous ammonia to form a yellow-brown solution</i></p> <p><i>C is incorrect because <math>\text{CoCl}_2(\text{aq})</math> is a pink solution</i></p>	1

Question number	Answer	Mark
7	<p><b>The only correct answer is B</b> (<math>[\text{Ni}(\text{EDTA})]^{2-}</math>)</p> <p><i>A is incorrect because chloride ions act as monodentate ligands</i></p> <p><i>C is incorrect because ethanedioate ions act as bidentate ligands</i></p> <p><i>D is incorrect because 1,2-diaminoethane molecules act as bidentate ligands</i></p>	1

Question number	Answer	Mark
8	<p><b>The only correct answer is D</b> (none of the products are harmful to the environment)</p> <p><i>A is incorrect because the reactions occurring in catalytic converters involve heterogeneous catalysis</i></p> <p><i>B is incorrect because carbon monoxide is adsorbed onto the surface of the catalyst</i></p> <p><i>C is incorrect because nitrogen is desorbed from the surface of the catalyst</i></p>	1

Question number	Answer	Mark
9	<p><b>The only correct answer is C</b> (<math>\text{Mn}^{2+}</math>)</p> <p><i>A is incorrect because <math>\text{MnO}_4^-</math> ions are neither a product nor a catalyst in this reaction</i></p> <p><i>B is incorrect because <math>\text{H}^+</math> ions are neither a product nor a catalyst in this reaction</i></p> <p><i>D is incorrect because <math>\text{CO}_2</math> is not a catalyst in this reaction</i></p>	1


Question number	Answer	Mark
10	<p><b>The only correct answer is A</b> (both <math>\text{Fe}^{2+}(\text{aq})</math> and <math>\text{Fe}^{3+}(\text{aq})</math> catalyse the reaction)</p> <p><i>B is incorrect because both <math>\text{Fe}^{2+}(\text{aq})</math> and <math>\text{Fe}^{3+}(\text{aq})</math> catalyse the reaction</i></p> <p><i>C is incorrect because both <math>\text{Fe}^{2+}(\text{aq})</math> and <math>\text{Fe}^{3+}(\text{aq})</math> catalyse the reaction</i></p> <p><i>D is incorrect because both <math>\text{Fe}^{2+}(\text{aq})</math> and <math>\text{Fe}^{3+}(\text{aq})</math> catalyse the reaction</i></p>	1

Question number	Answer	Mark
11	<p><b>The only correct answer is D</b> (  )</p> <p><i>A is incorrect because this product is formed by the substitution of one chlorine atom in <math>\text{CHCl}_3</math></i></p> <p><i>B is incorrect because this product is formed by the substitution of two chlorine atoms in <math>\text{CHCl}_3</math></i></p> <p><i>C is incorrect because this product is formed by the substitution of all three chlorine atoms in <math>\text{CHCl}_3</math></i></p>	1



Question number	Answer	Mark
12	<p><b>The only correct answer is C (330.7)</b></p> <p><i>A is incorrect because this is the molar mass of bromobenzene</i></p> <p><i>B is incorrect because this is the molar mass of the monosubstituted product</i></p> <p><i>D is incorrect because this is the molar mass of the fully substituted product</i></p>	1

Question number	Answer	Mark
13	<p><b>The only correct answer is C (<math>\text{CH}_3\text{CH}_2\text{CH}_2\text{NH}_2 &gt; \text{NH}_3 &gt; \text{C}_6\text{H}_5\text{NH}_2</math>)</b></p> <p><i>A is incorrect because <math>\text{C}_6\text{H}_5\text{NH}_2</math> is the weakest base in the sequence</i></p> <p><i>B is incorrect because <math>\text{CH}_3\text{CH}_2\text{CH}_2\text{NH}_2</math> is a stronger base than <math>\text{NH}_3</math></i></p> <p><i>D is incorrect because this shows the order of increasing basicity</i></p>	1

Question number	Answer	Mark
14	<p><b>The only correct answer is B</b> ()</p> <p><i>A is incorrect because this amine could be prepared by the reduction of butanenitrile</i></p> <p><i>C is incorrect because this amine could be prepared by the reduction of 2-methylpropanenitrile</i></p> <p><i>D is incorrect because this amine could be prepared by the reduction of 2,2-dimethylpropanenitrile</i></p>	1

Question number	Answer	Mark
15	<p><b>The only correct answer is B (4)</b></p> <p><i>A is incorrect because the repeat unit of the polymer is formed from four different amino acids</i></p> <p><i>C is incorrect because the repeat unit of the polymer is formed from four different amino acids</i></p> <p><i>D is incorrect because the repeat unit of the polymer is formed from four different amino acids</i></p>	1


Question number	Answer	Mark
16	<p><b>The only correct answer is D (carbon dioxide giving carboxylic acids )</b></p> <p><i>A is incorrect because Grignard reagents react with water giving alkanes</i></p> <p><i>B is incorrect because Grignard reagents react with methanal giving primary alcohols</i></p> <p><i>C is incorrect because Grignard reagents react with ketones giving tertiary alcohols only</i></p>	1

Question number	Answer	Mark
17	<p><b>The only correct answer is C (will be lower than the true value)</b></p> <p><i>A is incorrect because using a sample that is impure would cause the value to be lower</i></p> <p><i>B is incorrect because using a sample that is impure would cause the value to be lower</i></p> <p><i>D is incorrect because using a sample that is impure would cause the value to be lower</i></p>	1

**Total for Section A = 20 marks**

## Section B

Question Number	Answer	Additional guidance	Mark
18(a)	<ul style="list-style-type: none"> <li>any indication that <b>A</b> contains FeCl<sub>2</sub>/iron(II) chloride (1)</li> <li>working to show that <b>A</b> is a tetrahydrate (1)</li> </ul>	<p>Example of calculation:</p> <p>Ignore (<b>A</b> contains) Fe<sup>2+</sup> Ignore [FeCl<sub>4</sub>]<sup>2-</sup></p> <p>mass of water = 198.8 – (55.8 + 2 × 35.5) = 72.0 (g) moles of water = 72.0 ÷ 18.0 = 4</p> <p><b>A</b> is FeCl<sub>2</sub>.4H<sub>2</sub>O/iron(II) chloride tetrahydrate</p> <p>Allow FeCl<sub>2</sub>(H<sub>2</sub>O)<sub>4</sub></p>	2
Question Number	Answer	Additional guidance	Mark
18(b)	<ul style="list-style-type: none"> <li>[Fe(H<sub>2</sub>O)<sub>6</sub>]<sup>2+</sup></li> </ul>	<p>Allow [Fe(OH)(H<sub>2</sub>O)<sub>5</sub>]<sup>+</sup> / [Fe(Cl)(H<sub>2</sub>O)<sub>5</sub>]<sup>+</sup></p> <p>Ignore omission of square brackets Ignore name even if incorrect</p>	1

Question Number	Answer	Additional guidance	Mark
18(c)	<ul style="list-style-type: none"> <li>A diagram showing the octahedral shape</li> </ul>	<p>Example of diagram:</p>  <p>Accept arrows for dative covalent bonds</p> <p>Allow CN for C≡N Do not award KCN/HCN for C≡N Do not award M for Fe</p> <p>Ignore connectivity of CN ligands Ignore lone pairs Ignore omission of square brackets Ignore all charges</p> <p>Do not award diagrams with no 3D shape</p>	1

Question Number	Answer	Additional guidance	Mark
18(d)	<ul style="list-style-type: none"> <li data-bbox="340 292 1077 323">• moles of K, Fe (1)</li>   <li data-bbox="340 627 1077 659">• moles of C and N (1)</li>   <li data-bbox="340 890 1077 994">• calculation of K:Fe:C:N mole ratio <b>and</b> empirical formula (1)</li> </ul>	<p data-bbox="1108 215 1413 247">Example of calculation:</p> <p data-bbox="1108 292 1559 355">mol K = <math>35.6 \div 39.1 = 0.91049</math> Allow 0.91282 from <math>A_r</math> value of 39</p> <p data-bbox="1108 403 1559 467">mol Fe = <math>17.0 \div 55.8 = 0.30466</math> Allow 0.30357 from <math>A_r</math> value of 56</p> <p data-bbox="1108 515 1238 547">Ignore SF</p> <p data-bbox="1108 627 1491 659">mol C = <math>21.9 \div 12.0 = 1.8250</math></p> <p data-bbox="1108 699 1491 730">mol N = <math>25.5 \div 14.0 = 1.8214</math></p> <p data-bbox="1108 770 1395 802">Ignore SF except 1 SF</p> <div style="text-align: center; margin: 10px 0;"> <p data-bbox="1193 882 1597 914">K : Fe : C : N</p> <p data-bbox="1149 922 1641 954">0.91049 : 0.30466 : 1.825 : 1.8214</p> <p data-bbox="1193 962 1597 994">3 : 1 : 6 : 6</p> </div> <p data-bbox="1108 1002 1507 1034">empirical formula is <math>K_3FeC_6N_6</math></p> <p data-bbox="1108 1042 1350 1074">Allow <math>K_3Fe(CN)_6</math></p> <p data-bbox="1108 1082 1473 1114">Allow elements in any order</p> <p data-bbox="1108 1121 1899 1185">TE on moles of K, Fe, C and N provided empirical formula is closest whole number ratio</p> <p data-bbox="1108 1217 1653 1249">Correct answer with no working scores (3)</p>	<b>3</b>

Question Number	Answer	Additional guidance	Mark
18(e)	<p>An equation including:</p> <ul style="list-style-type: none"> <li><math>[\text{Fe}(\text{CN})_6]^{4-}</math> reactant <b>and</b> <math>[\text{Fe}(\text{CN})_6]^{3-}</math> product (1)</li> <li>rest of equation and balancing (1)</li> </ul>	<p>Example of equation:</p> $2[\text{Fe}(\text{CN})_6]^{4-} + \text{Cl}_2 \rightarrow 2[\text{Fe}(\text{CN})_6]^{3-} + 2\text{Cl}^-$ <p>Allow <math>\text{K}_4[\text{Fe}(\text{CN})_6]</math> reactant <b>and</b> <math>\text{K}_3[\text{Fe}(\text{CN})_6]</math> product Ignore omission of square brackets</p> <p><b>M2 dependent on M1</b> Allow multiples Ignore state symbols even if incorrect Do not award uncanceled <math>\text{K}^+</math> spectator ions</p> <p>If no other mark awarded, award (1) for any multiple of the following equation: <math>2\text{Fe}^{2+} + \text{Cl}_2 \rightarrow 2\text{Fe}^{3+} + 2\text{Cl}^-</math></p>	2

Question Number	Answer	Additional guidance	Mark												
18(f)	<p>A completed table showing:</p> <ul style="list-style-type: none"> <li>correct identification for reaction 2 (1)</li> <li>correct identification for reaction 3 (1)</li> </ul>	<p>Example of completed table:</p> <table border="1"> <thead> <tr> <th></th> <th>Neutralisation</th> <th>Ligand exchange</th> <th>Redox</th> </tr> </thead> <tbody> <tr> <td>Reaction 2</td> <td></td> <td>✓</td> <td></td> </tr> <tr> <td>Reaction 3</td> <td></td> <td></td> <td>✓</td> </tr> </tbody> </table> <p>Allow any form of positive identification, including crosses</p> <p>Ignore any form of negative identification</p> <p>Do not award more than 1 box ticked in each row</p>		Neutralisation	Ligand exchange	Redox	Reaction 2		✓		Reaction 3			✓	2
	Neutralisation	Ligand exchange	Redox												
Reaction 2		✓													
Reaction 3			✓												


(Total for Question 18 = 11 marks)

Question Number	Answer	Additional guidance	Mark
19(a)	<ul style="list-style-type: none"> <li data-bbox="338 250 1043 288">• suitable test: bromine water/Br<sub>2</sub>(aq) (1)</li>   <li data-bbox="338 624 1043 694">• result of test: decolourises (from orange) with Dewar structure (and no change with benzene) (1)</li> </ul>	<p data-bbox="1070 250 1693 320">Allow bromine / Br<sub>2</sub>(l) / Br<sub>2</sub> in organic solvent Do not award Br/Br<sup>-</sup></p> <p data-bbox="1070 362 1832 432">Accept potassium manganate(VII)/KMnO<sub>4</sub> <b>and</b> acidified/ named acid/H<sup>+</sup></p> <p data-bbox="1070 474 1440 544">Do not award hydrogenation Do not award combustion</p> <p data-bbox="1070 624 1850 767">Allow does not decolourise with benzene Allow brown/orange/yellow for colour of bromine water Allow red/brown/orange for colour of bromine Allow pink/purple for colour of potassium manganate(VII)</p> <p data-bbox="1070 809 1585 879">Ignore colour fades for decolourises Ignore reference to addition/substitution</p> <p data-bbox="1070 920 1843 984">Do not award any reference to decolourisation/reaction with benzene</p>	2

Question Number	Answer	Additional guidance	Mark
19(b)	<ul style="list-style-type: none"> <li data-bbox="338 327 1120 359">• similarity: (both compounds have) one (NMR) peak (1)</li>   <li data-bbox="338 624 1120 655">• difference: expected chemical shift values (1)</li> </ul>	<p data-bbox="1144 248 1704 280">Ignore any reference to IR and/or <math>^{13}\text{C}</math> NMR</p> <p data-bbox="1144 327 1854 395">Allow (both compounds have) one proton environment Ignore just same number of peaks</p> <p data-bbox="1144 437 1899 469">Ignore references to relative peak area/integration/splitting</p> <p data-bbox="1144 510 1671 542">Do not award any other number of peaks</p> <p data-bbox="1144 620 1883 689">chemical shift for benzene within range of 6.4 to 8.4 ppm (actual value is 7.3 ppm)</p> <p data-bbox="1144 695 1845 802"><b>and</b> chemical shift for Ladenburg structure within range of 0 to 2.3 ppm (actual value is 2.3 ppm)</p> <p data-bbox="1144 844 1783 876">Allow any range or value within the above ranges</p> <p data-bbox="1144 917 1917 986">Ignore just benzene would have a higher chemical shift than Ladenburg structure or reverse argument</p> <p data-bbox="1144 1027 1778 1059">Do not award additional incorrect chemical shifts</p>	2




Question Number	Answer	Additional guidance	Mark
19(c)	<p>An explanation that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>showed that all C–C bonds are the same length in benzene (1)</li> <li>in Kekulé structure the C=C bonds would be shorter than the C–C bonds (or reverse argument) (1)</li> </ul>	<p>Ignore any reference to:            C–H bonds            bond strength/bond angle            delocalised electrons            Dewar/Ladenburg structures</p> <p>Allow showed benzene is a regular hexagon</p> <p>Allow showed benzene contains only one type of carbon-carbon bond</p> <p>Allow benzene bond lengths are in between C=C and C–C</p> <p>Ignore just benzene has no C=C bonds</p> <p>Do not award benzene bond lengths are longer than C–C/            shorter than C=C</p> <p>Allow Kekulé structure would have shown two different lengths/types of carbon-carbon bond</p> <p>Allow Kekulé structure would have alternating carbon-carbon bond lengths</p> <p>Ignore just Kekulé has C=C bonds</p> <p>Do not award C–C bonds would be shorter than the C=C bonds</p> <p>If no other mark awarded, just bond lengths equal in benzene but different in Kekulé scores (1)</p>	2


Question Number	Answer	Additional guidance	Mark
19(d)(i)	<p>A diagram showing:</p> <ul style="list-style-type: none"> <li>• correct relative stabilities (1)</li> <li>• two or three numerical differences in enthalpy with appropriate arrows (1)</li> </ul>	<p>Example of diagram:</p>  <p>Allow names for structures</p> <p>If three values and arrows are given they must all be correct to score M2</p> <p>Allow slight imprecision in start and end of arrows in M2</p> <p>Ignore any x-axis label</p> <p>Do not award double headed arrows in M2</p> <p>Do not award incorrect sign in M2</p>	2

Question Number	Answer	Additional guidance	Mark
19(d)(ii)	<p>An answer that makes reference to the following:</p> <ul style="list-style-type: none"> <li>• pi bonds are weaker/more reactive/require less energy to break (than sigma bonds)</li> </ul> <p><b>or</b></p> <p>fewer bonds must break to convert the Dewar structure to benzene</p>	<p>Ignore just Dewar structure has pi/double bonds/is unsaturated            Ignore just Dewar structure has weaker bonds            Do not award C=C/double bonds weaker/require less energy to break (than C–C/single bonds)</p> <p>Accept reverse argument            Allow any specified numbers to indicate fewer bonds must break</p> <p>Ignore fewer new bonds must form            Ignore Dewar structure is more similar to benzene            Ignore carbon atoms already in a ring/hexagon            Ignore any reference to intermolecular forces</p> <p>Do not award (higher) ring strain in Ladenburg            Do not award smaller difference in enthalpy            Do not award Ladenburg structure more stable</p>	<b>1</b>

Question Number	Answer	Additional guidance	Mark
19(e)	<p>An explanation that makes reference to the following points:</p> <p><b><i>E</i>-hexa-1,4-diene</b></p> <ul style="list-style-type: none"> <li>twice the hydrogenation enthalpy (of hex-3-ene) as two (isolated) C=C bonds (1)</li> </ul> <p><b><i>E</i>-hexa-1,3-diene</b></p> <ul style="list-style-type: none"> <li>less exothermic/more stable (by 22 kJ mol<sup>-1</sup> than <i>E</i>-hexa-1,4-diene <b>and</b> as some delocalisation of pi-bond(s) (1)</li> </ul>	<p>Allow double bond for C=C throughout</p> <p>Accept <math>-118 \times 2</math> (<math>=-236</math>) as two C=C bonds</p> <p>Allow twice the hydrogenation enthalpy as no delocalisation of pi-bond(s)</p> <p>Accept less negative</p> <p>Allow more positive</p> <p>Allow some delocalisation of double bond(s)</p> <p>Allow double bonds/p-orbitals are conjugated</p> <p>Allow double bonds/p-orbitals are close enough to overlap</p> <p>Ignore just C=C are close</p> <p>Ignore just delocalisation of electrons</p> <p>Ignore electron density more spread out</p> <p>Ignore resonance stabilised</p>	2

Question Number	Answer	Additional guidance	Mark
19(f)(i)	<ul style="list-style-type: none"> <li>skeletal formula of any one isomer (1)</li> <li>skeletal formulae of second and third isomers (1)</li> </ul>	<p>Example of correct skeletal formulae:</p>  <p>(1,2-isomer)      (1,3-isomer)      (1,4-isomer)</p> <p>Allow Kekulé benzene ring</p> <p>Allow structural/displayed CH<sub>3</sub> and CH<sub>3</sub>CO groups</p> <p>If no other mark awarded, 1,2-, 1,3- and 1,4-isomers with incorrect side chain/cyclohexane ring scores (1)</p>	2


Question Number	Answer	Additional guidance	Mark
19(f)(ii)	<ul style="list-style-type: none"> <li>(identification of <b>X</b> as) 1,4-isomer (1)</li> <li>(7 peaks consistent with) 7 carbon environments (1)</li> </ul>	<p>Allow any form of identification, including (f)(i) annotation Allow just '1,4' or 'para'</p> <p><b>M2 dependent on a structure containing 7 carbon environments</b></p> <p>Accept 1,2-isomer and/or 1,3-isomer have 9 carbon environments/would have 9 peaks</p> <p>Allow (4 arene peaks consistent with) 4 arene carbon environments</p> <p>Allow 1,2-isomer and/or 1,3-isomer have 6 arene carbon environments/would have 6 arene peaks</p>	2

Question Number	Answer	Additional guidance	Mark
19(f)(iii)	<p>A mechanism including:</p> <ul style="list-style-type: none"> <li>• curly arrow from on or within circle to C<sup>+</sup> of CH<sub>3</sub>CO<sup>+</sup> (1)</li> <li>• structure of intermediate ion (1)</li> <li>• curly arrow from C–H bond to within ring <b>and</b> correct product (1)</li> <li>• balanced equation for regeneration of catalyst (1)</li> </ul>	<p>Example of mechanism:</p>  <p>Allow Kekulé benzene ring</p> <p>‘Horseshoe’ facing tetrahedral carbon and covering at least three carbons with some part of positive sign within ‘horseshoe’</p> <p>Allow methyl at 1,2- or 1,3-positions Ignore missing methyl substituent</p> <p>Do not award dotted/dashed C–H/C–C bonds unless 3D structure</p> <p>Allow 1,2- or 1,3-product from corresponding intermediate Do not award missing methyl substituent</p> <p><math>\text{AlCl}_4^- + \text{H}^+ \rightarrow \text{AlCl}_3 + \text{HCl}</math></p>	4

(Total for Question 19 = 19 marks)

Question Number	Answer	Additional guidance	Mark
20(a)	<p>An answer that makes reference to one of the following points:</p> <ul style="list-style-type: none"> <li>• variable oxidation state/oxidation number <b>or</b></li> <li>• (easily) oxidised <b>and</b> reduced (back to original oxidation state) <b>or</b></li> <li>• (easily) donate <b>and</b> accept electrons (from other molecules/species)</li> </ul>	<p>Ignore references to ionisation energy Ignore partially full d orbital(s)/d subshell Ignore more than one stable ion Ignore references to heterogeneous catalysis/adsorption Ignore references to alternative reaction pathways/activation energy</p> <p>Allow can change oxidation state/oxidation number Allow have different oxidation state(s)/oxidation number(s) Ignore variable valency</p> <p>Allow just lose and gain electrons (easily)</p>	1

Question Number	Answer	Additional guidance	Mark
20(b)(i)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>• monodentate: forms a single/one dative (covalent) bond (1)</li> <li>• ligand: (a species with a) lone pair (of electrons) that can form a dative (covalent) bond to a (central transition) metal (ion) (1)</li> </ul>	<p>Accept coordinate for dative throughout</p> <p>Accept donates a single/one lone pair Allow occupies a single/one coordination site</p>	2

Question Number	Answer	Additional guidance	Mark
20(b)(ii)	<p>A completed diagram showing:</p> <ul style="list-style-type: none"> <li>three adjacent THF/Cl ligands</li> </ul>	<p>Expected diagram:</p>  <p>Ignore lone pairs Ignore 1- charge on Cl ligands</p>	1

Question Number	Answer	Additional guidance	Mark
20(c)(i)	<ul style="list-style-type: none"> <li>yellow to (permanent pale) green</li> </ul>	<p>Ignore qualifiers, eg pale Ignore precipitate</p> <p>Do not award green to yellow Do not award any combination of yellow and green, eg yellowy-green Do not award any other colour</p>	1



Question Number	Answer	Additional guidance	Mark
20(c)(ii)	<p>A calculation including:</p> <ul style="list-style-type: none"> <li>• moles of <math>\text{Ti}^{3+}</math> in titre (1)</li> <li>• moles of <math>\text{Mg}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}</math> in <math>100 \text{ cm}^3</math> (1)</li> <li>• moles of <math>\text{NO}_3^-</math> in <math>10.00 \text{ cm}^3</math> (1)</li> <li>• <math>\text{Ti}^{3+} : \text{NO}_3^-</math> mol ratio (1)</li> <li>• final oxidation state of nitrogen (1)</li> </ul>	<p>Example of calculation:</p> <p>moles of <math>\text{Ti}^{3+} = 0.085 \times \frac{20.70}{1000} = 0.0017595 / 1.7595 \times 10^{-3}</math></p> <p>moles of <math>\text{Mg}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O} = \frac{0.75}{256.3} = 0.0029263 / 2.9263 \times 10^{-3}</math></p> <p>moles of <math>\text{NO}_3^- = \frac{0.0029263}{10} \times 2 = 0.00058525 / 5.8525 \times 10^{-4}</math></p> <p>TE on moles <math>\text{Mg}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}</math></p> <p>Ignore SF except 1 SF in M1, M2 and M3 Ignore truncation of moles in M1, M2 and M3, eg 0.0005852</p> <p><math>\text{Ti}^{3+} : \text{NO}_3^-</math> mol ratio = <math>0.0017595 : 0.00058525</math> = 3 : 1</p> <p>TE on moles <math>\text{Ti}^{3+}</math> and moles <math>\text{NO}_3^-</math></p> <p>final oxidation state of nitrogen = (+)2</p> <p>TE on mol ratio provided final oxidation state of nitrogen is between -3 and +4, eg <math>\text{Ti}^{3+} : \text{NO}_3^-</math> mol ratio = <math>0.0017595 : 0.00029263</math> = 6 : 1</p> <p>final oxidation state of nitrogen = -1</p> <p>Do not award incorrect oxidation state of N in <math>\text{NO}_3^-</math></p> <p>Correct answer with no working scores (1)</p>	5

Question Number	Answer	Additional guidance	Mark
20(c)(iii)	<p>An equation including:</p> <ul style="list-style-type: none"> <li>• selection of correct nitrogen half-equation (1)</li> <li>• balanced ionic equation from chosen half-equations (1)</li> </ul>	<p>Example of equation:  <math>3\text{Ti}^{3+} + \text{H}_2\text{O} + \text{NO}_3^- \rightarrow 3\text{TiO}^{2+} + 2\text{H}^+ + \text{NO}</math></p> <p>TE on (c)(ii) provided +3 or +4 oxidation state</p> <p>Allow multiples            Allow reversible arrow            Ignore state symbols even if incorrect</p>	2

Question Number	Answer	Additional guidance	Mark
20(c)(iv)	<p>An answer that makes reference to the following point:</p> <ul style="list-style-type: none"> <li>• (calculation of) <math>E^\ominus_{\text{cell}}</math> value</li> </ul>	<p>Example of calculation:</p> <p><math>(E^\ominus_{\text{cell}} = 0.96 - 0.10 =) (+)0.86 \text{ (V)}</math></p> <p>TE on ionic equation from (c)(iii):            (+)0.7(0) (V) for <math>\text{Ti}^{3+} + \text{NO}_3^- \rightarrow \text{TiO}^{2+} + \text{NO}_2</math>            (+)0.84 (V) for <math>2\text{Ti}^{3+} + \text{H}_2\text{O} + \text{NO}_3^- \rightarrow 2\text{TiO}^{2+} + \text{H}^+ + \text{HNO}_2</math></p>	1

Question Number	Answer	Additional guidance	Mark
20(c)(v)	<p>An answer that makes reference to the following point:</p> <ul style="list-style-type: none"> <li>• (heat is to) speed up/increase rate of reaction</li> </ul>	<p>Allow to ensure fast oxidation of <math>\text{Ti}^{3+}</math>            Allow to provide activation energy/<math>E_a</math>            Allow (reaction has a) high activation energy/<math>E_a</math></p> <p>Ignore just to provide (more) energy            Ignore to increase collision frequency            Ignore to ensure complete reaction            Ignore any reference to thermodynamic feasibility</p>	1


Question Number	Answer	Additional guidance	Mark																				
20(c)(vi)	<p>This question assesses a student's ability to show a coherent and logically structured answer with linkages and fully-sustained reasoning.</p> <p>Marks are awarded for indicative content and for how the answer is structured and shows lines of reasoning.</p> <p>The following table shows how the marks should be awarded for indicative content.</p> <table border="1" data-bbox="338 579 1176 847"> <thead> <tr> <th data-bbox="338 579 779 655">Number of indicative marking points seen in answer</th> <th data-bbox="779 579 1176 655">Number of marks awarded for indicative marking points</th> </tr> </thead> <tbody> <tr> <td data-bbox="338 655 779 692">6</td> <td data-bbox="779 655 1176 692">4</td> </tr> <tr> <td data-bbox="338 692 779 729">5-4</td> <td data-bbox="779 692 1176 729">3</td> </tr> <tr> <td data-bbox="338 729 779 766">3-2</td> <td data-bbox="779 729 1176 766">2</td> </tr> <tr> <td data-bbox="338 766 779 802">1</td> <td data-bbox="779 766 1176 802">1</td> </tr> <tr> <td data-bbox="338 802 779 839">0</td> <td data-bbox="779 802 1176 839">0</td> </tr> </tbody> </table> <p>The following table shows how the marks should be awarded for structure and lines of reasoning.</p> <table border="1" data-bbox="338 959 1176 1406"> <thead> <tr> <th data-bbox="338 959 813 1070"></th> <th data-bbox="813 959 1176 1070">Number of marks awarded for structure and sustained lines of reasoning</th> </tr> </thead> <tbody> <tr> <td data-bbox="338 1070 813 1222">Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning demonstrated throughout.</td> <td data-bbox="813 1070 1176 1222">2</td> </tr> <tr> <td data-bbox="338 1222 813 1334">Answer is partially structured with some linkages and lines of reasoning.</td> <td data-bbox="813 1222 1176 1334">1</td> </tr> <tr> <td data-bbox="338 1334 813 1406">Answer has no linkages between points and is unstructured.</td> <td data-bbox="813 1334 1176 1406">0</td> </tr> </tbody> </table>	Number of indicative marking points seen in answer	Number of marks awarded for indicative marking points	6	4	5-4	3	3-2	2	1	1	0	0		Number of marks awarded for structure and sustained lines of reasoning	Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning demonstrated throughout.	2	Answer is partially structured with some linkages and lines of reasoning.	1	Answer has no linkages between points and is unstructured.	0	<p>The mark for indicative content should be added to the mark for lines of reasoning. For example, an answer with five indicative marking points that is partially structured with some linkages and lines of reasoning scores 4 marks (3 marks for indicative content and 1 mark for partial structure and some linkages and lines of reasoning).</p> <p>If there are no linkages between points, the same five indicative marking points would yield an overall score of 3 marks (3 marks for indicative content and no marks for linkages).</p> <p>If there is any incorrect chemistry, deduct mark(s) from the reasoning. If no reasoning mark(s) awarded, do not deduct mark(s).</p> <p>Comment: Look for the indicative marking points first, then consider the mark for the structure of the answer and sustained line of reasoning.</p>	6
Number of indicative marking points seen in answer	Number of marks awarded for indicative marking points																						
6	4																						
5-4	3																						
3-2	2																						
1	1																						
0	0																						
	Number of marks awarded for structure and sustained lines of reasoning																						
Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning demonstrated throughout.	2																						
Answer is partially structured with some linkages and lines of reasoning.	1																						
Answer has no linkages between points and is unstructured.	0																						


	<p>Indicative points:</p> <ul style="list-style-type: none"> <li>• IP1: identification of <math>[\text{Ti}(\text{H}_2\text{O})_6]^{3+}</math> complex ion</li> <li>• IP2: partially filled d-subshell/d-orbital(s) (in <math>\text{Ti}^{3+}</math>)</li> <li>• IP3: splitting in energy of d-subshell/d-orbitals by water/ligands</li> <li>• IP4: absorption of light/photon/(electromagnetic) radiation <b>and</b> electronic transition</li> <li>• IP5: origin of observed colour of complex ion</li> <li>• IP6: clearer colour change at end-point with indicator</li> </ul>	<p>Allow <math>[\text{Ti}(\text{H}_2\text{O})_5\text{Cl}]^{2+}</math> / <math>[\text{Ti}(\text{H}_2\text{O})_4\text{Cl}_2]^+</math></p> <p>Accept incomplete for partially filled Accept (<math>\text{Ti}^{3+}</math> is) <math>1s^22s^22p^63s^23p^63d^1(4s^0)</math> Allow (<math>\text{Ti}^{3+}</math> is) <math>(3)d^1</math></p> <p>water/ligands <b>and</b> split the energy of the d-subshell/d-orbitals Allow ligands cause d-d splitting Do not award d-orbital (singular)</p> <p>(visible) light/photon/(electromagnetic) radiation is absorbed <b>and</b> promoting electrons from lower to higher energy Allow light etc causes d-d transitions</p> <p>colour due to reflected/transmitted light Allow colour due to wavelengths/frequencies of light that are not absorbed Allow complementary colour observed Do not award any reference to emission of light</p> <p>Accept reverse argument Allow colours are more intense/distinct/sharp/strong Allow concentration (of <math>[\text{Ti}(\text{H}_2\text{O})_6]^{3+}/\text{TiCl}_3</math>) too low to accurately determine end-point in absence of indicator Ignore just easier to determine end-point Ignore just more accurate/precise Ignore mention of specific colours, even if incorrect Do not award reference to acid-base colour change</p>	
--	---	---	--

(Total for Question 20 = 20 marks)

**TOTAL FOR SECTION B = 50 MARKS**


## Section C

Question Number	Answer	Additional guidance	Mark
21(a)	<p>A completed mechanism showing:</p> <ul style="list-style-type: none"> <li>curly half-arrows to show homolytic fission of O–H bond</li> </ul>	<p>Example of completed mechanism:</p>  <p>Accept curly half-arrows originating from opposite sides of the O–H bond</p> <p>Left-hand curly half-arrow must terminate between T• and H</p> <p>Right hand curly half-arrow must terminate on or near to O atom of H–O</p>	1

Question Number	Answer	Additional guidance	Mark
21(b)	<p>A completed mechanism showing:</p> <ul style="list-style-type: none"> <li>• curly arrow from lone pair on <math>\text{Se}^-</math> to correct C of <math>\text{C}=\text{C}</math> (1)</li> <li>• curly arrow from <math>\text{C}=\text{C}</math> bond to <math>\text{C}-\text{C}</math> bond</li> </ul> <p><b>and</b></p> <ul style="list-style-type: none"> <li>curly arrow from <math>\text{C}=\text{O}</math> bond to O (1)</li> </ul>	<p>Example of completed mechanism:</p>  <p>Penalise curly half-arrows once only</p> <p>Do not award curly arrow from negative charge on <math>\text{Se}^-</math></p> <p>Ignore <math>(\delta^+)\text{C}=\text{C}(\delta^-)</math> dipole  Do not award <math>(\delta^-)\text{C}=\text{C}(\delta^+)</math> dipole  Do not award full charge on either carbon of <math>\text{C}=\text{C}</math> bond</p> <p>Do not award incorrect <math>(\delta^-)\text{C}=\text{O}(\delta^+)</math> dipole</p>	2


Question Number	Answer	Additional guidance	Mark															
21(c)	A completed table showing: <ul style="list-style-type: none"> <li>two or three correct answers (1)</li> <li>four correct answers (2)</li> </ul>	Example of completed table: <table border="1" style="margin-left: 20px;"> <thead> <tr> <th></th> <th>[Au(Curc)<sub>2</sub>]<sup>+</sup></th> <th>[Al(Curc)(C<sub>2</sub>H<sub>5</sub>OH)<sub>2</sub>(NO<sub>3</sub>)<sub>2</sub>]</th> </tr> </thead> <tbody> <tr> <td>Coordination number</td> <td>4</td> <td><u>6</u></td> </tr> <tr> <td>O–M–O bond angle</td> <td>90°</td> <td><u>90°</u> Ignore 180°</td> </tr> <tr> <td>Shape</td> <td><u>square planar</u></td> <td>octahedral</td> </tr> <tr> <td>Charge on metal ion</td> <td><u>+3</u></td> <td>+3</td> </tr> </tbody> </table>		[Au(Curc) <sub>2</sub> ] <sup>+</sup>	[Al(Curc)(C <sub>2</sub> H <sub>5</sub> OH) <sub>2</sub> (NO <sub>3</sub> ) <sub>2</sub> ]	Coordination number	4	<u>6</u>	O–M–O bond angle	90°	<u>90°</u> Ignore 180°	Shape	<u>square planar</u>	octahedral	Charge on metal ion	<u>+3</u>	+3	2
	[Au(Curc) <sub>2</sub> ] <sup>+</sup>	[Al(Curc)(C <sub>2</sub> H <sub>5</sub> OH) <sub>2</sub> (NO <sub>3</sub> ) <sub>2</sub> ]																
Coordination number	4	<u>6</u>																
O–M–O bond angle	90°	<u>90°</u> Ignore 180°																
Shape	<u>square planar</u>	octahedral																
Charge on metal ion	<u>+3</u>	+3																

Question Number	Answer	Additional guidance	Mark
21(d)(i)	<ul style="list-style-type: none"> <li>K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> <b>and</b> H<sub>2</sub>SO<sub>4</sub> (1)</li> <li>heat/reflux (1)</li> </ul>	<p><b>If name and formula given, both must be correct to score M1</b></p> <p>Accept names (eg sodium dichromate(VI) and sulfuric acid)            Allow Cr<sub>2</sub>O<sub>7</sub><sup>2-</sup> <b>and</b> H<sup>+</sup> / acidified dichromate</p> <p>Ignore concentration of acid</p> <p>Do not award KMnO<sub>4</sub> for K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub></p> <p>Do not award HCl for H<sub>2</sub>SO<sub>4</sub></p> <p><b>M2 dependent on some mention of dichromate (or manganate) oxidising agent</b>            Ignore distillation</p>	2


Question Number	Answer	Additional guidance	Mark
21(d)(ii)	<ul style="list-style-type: none"> <li>correct structure for 2-aminobenzoic acid</li> </ul>	<p>Example of correct structure:</p>  <p>Accept hydrochloride salt (<math>-\text{NH}_3\text{Cl}</math>)  Allow protonated amine group (<math>-\text{NH}_3^+</math>)</p> <p>Allow any correct combination of skeletal, structural or displayed formulae</p> <p>Allow Kekulé benzene</p> <p>Ignore connectivity  Ignore name, even if incorrect</p>	<b>1</b>

Question Number	Answer	Additional guidance	Mark
21(d)(iii)	<ul style="list-style-type: none"> <li><math>\text{NaNO}_2</math>/sodium nitrite/sodium nitrate(III)  <b>and</b>  HCl/hydrochloric acid</li> </ul>	<p>Allow <math>\text{HNO}_2</math>/nitrous acid  Allow <math>\text{H}^+</math> <b>and</b> <math>\text{NO}_2^-</math></p> <p>Ignore conditions, including concentration of HCl  Ignore <math>\text{H}_2\text{O}</math></p> <p>Do not award <math>\text{NaNO}_3</math>/sodium nitrate</p>	<b>1</b>




Question Number	Answer	Additional guidance	Mark
21(d)(iv)	<ul style="list-style-type: none"> <li>correct structure for <i>N,N</i>-dimethylphenylamine</li> </ul>	<p>Example of correct structure:</p>  <p>Allow any correct combination of skeletal, structural or displayed formulae</p> <p>Allow Kekulé benzene</p> <p>Ignore quaternary salt (<math>-\text{N}(\text{CH}_3)_3^+</math>)</p> <p>Ignore name, even if incorrect</p>	1



Question Number	Answer	Additional guidance	Mark
21(d)(v)	<p>An explanation including:</p> <ul style="list-style-type: none"> <li>effect of temperature higher than 5°C (1)</li> <li>effect of temperature lower than 5°C (1)</li> </ul>	<p>(diazonium/it) decomposes / reacts with water / forms a phenol / undergoes nucleophilic substitution (above 5°C)</p> <p>Ignore byproducts form / side reactions occur / yield too low (above 5°C)</p> <p>(rate of reaction) too slow (below 5°C) Allow just slows down (below 5°C)</p> <p>Ignore insufficient energy for reaction to occur (below 5°C) Ignore any reference to activation energy/collision frequency Ignore freezes (at 0°C and below)</p>	2

Question Number	Answer	Additional guidance	Mark
21(e)(i)	<p>A completed equation showing:</p> <ul style="list-style-type: none"> <li>correct balancing of propanone and sodium ethanoate (1)</li> <li>correct balancing of sodium hydroxide and water (1)</li> </ul>	<p>Example of equation:</p>  <p><b>M2 dependent on M1</b></p>	2

Question Number	Answer	Additional guidance	Mark
21(e)(ii)	<p><b>M1: molar masses</b></p> <ul style="list-style-type: none"> <li><math>M(2\text{-nitrobenzaldehyde})</math> <b>and</b> <math>M(\text{indigotin})</math></li> </ul> <p>(1)</p> <p>Then, for M2 and M3, either:</p> <p><b>Method 1 (M2 and M3)</b></p> <ul style="list-style-type: none"> <li>moles indigotin in 10.0 g</li> </ul> <p><b>and</b> indigotin:2-nitrobenzaldehyde mol ratio</p> <p>(1)</p> <ul style="list-style-type: none"> <li>moles 2-nitrobenzaldehyde required</li> </ul> <p><b>and</b> mass 2-nitrobenzaldehyde required</p> <p>(1)</p>	<p>Correct answer with no working scores (3)</p> <p>Example of calculation:</p> $M(2\text{-nitrobenzaldehyde}) = 7 \times 12.0 + 5 \times 1.0 + 1 \times 14.0 + 3 \times 16.0 = 151.0 \text{ g mol}^{-1}$ $M(\text{indigotin}) = 16 \times 12.0 + 10 \times 1.0 + 2 \times 14.0 + 2 \times 16.0 = 262.0 \text{ g mol}^{-1}$ <p>Allow truncation of mass/moles throughout, eg 0.03816 for 0.038168 Ignore SF except 1 SF in M2 and M3</p> <p>moles indigotin in 10.0 g = <math>\frac{10.0}{262} = 0.038168</math></p> <p>TE on <math>M(\text{indigotin})</math></p> <p>moles 2-nitrobenzaldehyde = <math>2 \times 0.038168 (= 0.076336)</math> TE on moles indigotin</p> <p>moles 2-nitrobenzaldehyde required = <math>\frac{100}{85} \times 0.076336 = 0.089807</math></p> <p>mass 2-nitrobenzaldehyde required = <math>0.089807 \times 151 = 13.561 = 14 \text{ (g)}</math></p> <p>TE on moles 2-nitrobenzaldehyde TE on <math>M(2\text{-nitrobenzaldehyde})</math></p>	3

	<p>OR</p> <p><b>Method 2 (M2 and M3)</b></p> <ul style="list-style-type: none"> <li>mass indigotin if 100% yield</li> </ul> <p><b>and</b></p> <p>moles indigotin if 100% yield (1)</p> <ul style="list-style-type: none"> <li>indigotin:2-nitrobenzaldehyde mol ratio</li> </ul> <p><b>and</b></p> <p>mass 2-nitrobenzaldehyde (1)</p>	<p>mass indigotin if 100% yield = <math>\frac{100}{85} \times 10.0</math> = 11.765 (g)</p> <p>moles indigotin if 100% yield = <math>\frac{11.765}{262} = 0.044903</math></p> <p>TE on <math>M(\text{indigotin})</math></p> <p>moles 2-nitrobenzaldehyde = <math>2 \times 0.044903 (= 0.089807)</math></p> <p>TE on moles indigotin</p> <p>mass 2-nitrobenzaldehyde = <math>0.089807 \times 151</math> = 13.561 = 14 (g)</p> <p>TE on moles 2-nitrobenzaldehyde</p> <p>TE on <math>M(2\text{-nitrobenzaldehyde})</math></p>	
--	---	--	--

Question Number	Answer	Additional guidance	Mark
21(f)(i)	<p>A drawing showing:</p> <ul style="list-style-type: none"> <li>hydrolysed ester linkage <b>and</b> correct carbon frame and amine group</li> <li>deprotonated carboxylic acid <b>and</b> phenol groups</li> </ul>	<p>Example of drawing:</p> <p>(1) </p> <p>(1)</p> <p>Allow -ONa Do not award -O-Na</p> <p>Allow <i>E</i> isomer Allow Kekulé benzene</p> <p>Allow any correct combination of skeletal, structural or displayed formulae</p>	2

Question Number	Answer	Additional guidance	Mark
21(f)(ii)	A drawing showing: <ul style="list-style-type: none"> <li>• correct condensation product</li> </ul>	Example of drawing:    Allow Kekulé benzene  Allow any correct combination of skeletal, structural or displayed formulae  Allow diacylated product, ie  	1

Total for Section C = 20 marks

Total for Paper = 90 marks

