

Mark Scheme (SAM)

Pearson Edexcel International Advanced Subsidiary in Chemistry

Unit 3: Chemistry Laboratory Skills I

All the material in this publication is copyright
© Pearson Education Ltd 2013

General marking guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed-out work should be marked UNLESS the candidate has replaced it with an alternative response.
- Mark schemes will indicate within the table where, and which strands of Quality of Written Communication, are being assessed. The strands are as follows:
 - i. ensure that text is legible and that spelling, punctuation and grammar are accurate so that meaning is clear
 - ii. select and use a form and style of writing appropriate to purpose and to complex subject matter
 - iii. organise information clearly and coherently, using specialist vocabulary when appropriate.

Using the Mark Scheme

Examiners should NOT give credit for incorrect or inadequate answers, but allow candidates to be rewarded for answers showing correct application of principles and knowledge. Examiners should therefore read carefully and consider every response: even if it is not what is expected, it may still be creditworthy.

The mark scheme gives examiners:

- an idea of the types of response expected
- how individual marks are to be awarded
- the total mark for each question
- examples of responses that should NOT receive credit.

| | |
|-------------|---|
| / | Means that the responses are alternatives and either answer should receive full credit. |
| () | Means that a phrase/word is not essential for the award of the mark, but helps the examiner to get the sense of the expected answer. |
| Bold | Phrases/words in bold indicate that the meaning of the phrase or the actual word is essential to the answer. |
| ecf/TE/cq | (error carried forward)(transfer error)(consequential) means that a wrong answer given in an earlier part of a question is used correctly in answer to a later part of the same question. |

Candidates must make their meaning clear to the examiner to gain the mark. Do not give credit for correct words/phrases which are put together in a meaningless manner. Answers must be in the correct context.

| Question Number | Acceptable Answer | Reject | Mark |
|-----------------|---------------------------------|---|------|
| 1(a)(i) | Potassium (ions)/K ⁺ | K/incorrect formula Name with incorrect formula, e.g. 'Potassium, K' | (1) |

| Question Number | Acceptable Answer | Reject | Mark |
|-----------------|--|--|------|
| 1(a)(ii) | No precipitate forms/no change/no reaction/colourless solution . ALLOW clear for colourless. | White precipitate dissolves. Just 'dissolves' | (1) |

| Question Number | Acceptable Answer | Reject | Mark |
|-----------------|--|-------------------------|------|
| 1(a)(iii) | Silver nitrate (solution)/AgNO ₃ ALLOW acidified silver nitrate. (1) Yellow precipitate/solid ALLOW yellow suspension. (1) Second mark depends on first mark (use of silver nitrate). | Pale yellow precipitate | (2) |

| Question Number | Acceptable Answer | Reject | Mark |
|-----------------|--|------------|------|
| 1(a)(iv) | (Precipitate) does not dissolve/(precipitate) is insoluble/(precipitate) becomes paler in colour. ALLOW 'no change/no reaction'. ALLOW mark for insoluble even if wrong reagent is used in (a)(iii) to form a precipitate regardless of colour. Mark can only be given if there is a precipitate in (a)(iii). | Grey solid | (1) |

| Question Number | Acceptable Answer | Reject | Mark |
|-----------------|---|---|------|
| 1(a)(v) | KI Consequential on cation other than K ⁺ in (a)(i). ALLOW K ⁺ I ⁻ | Just potassium iodide Formula based on cation with incorrect charge or anion other than iodide | (1) |

| Question Number | Acceptable Answer | Reject | Mark |
|-----------------|---|---|------|
| 1(b)(i) | Calcium (ions)/Ca ²⁺ ALLOW +2 for 2+. | Ca/incorrect formula Name with incorrect formula, e.g. 'Calcium, Ca' | (1) |

| Question Number | Acceptable Answer | Reject | Mark |
|-----------------|---|--------|------|
| 1(b)(ii) | Dissolved/disappeared (1) Limewater/calcium hydroxide (solution) /Ca(OH) ₂ ((aq)) (1) | Melted | (2) |

| Question Number | Acceptable Answer | Reject | Mark |
|-----------------|---|---|------|
| 1(b)(iii) | CaCO ₃ ALLOW Ca(HCO ₃) ₂ . TE on incorrect metal ion in b(i) if correct formula given e.g SrCO ₃ , Na ₂ CO ₃ . | Name Formula based on cation with incorrect charge, e.g Ca ₂ CO ₃ or anion other than carbonate or hydrogencarbonate | (1) |

| Question Number | Acceptable Answer | Reject | Mark |
|-----------------|--|--|------|
| 1(c)(i) | No double bonds between C atoms /C=C absent/not an alkene/ Z is saturated/only single bonds between C atoms. ALLOW not an alkene or alkyne. | Just 'no double bonds' Just 'single bond(s)' alkane or any other functional group stated even if alcohol | (1) |

| Question Number | Acceptable Answer | Reject | Mark |
|-----------------|--|---|------|
| 1(c)(ii) | Alcohol/(-)OH/ROH/hydroxyl group present. ALLOW (-)COH. | OH ⁻ /hydroxide for hydroxyl CHO Carboxylic acid Phenol | (1) |

| Question Number | Acceptable Answer | Reject | Mark |
|------------------|---|-----------------------------|------------|
| 1(c)(iii) | Primary/1° alcohol (1) | Just 'alcohol' | (2) |
| | Secondary/2° alcohol (1) | Tertiary alcohols | |
| | ALLOW Not a tertiary alcohol for 1 mark. | Other specific examples | |
| | ALLOW propan-1-ol (1) propan-2-ol (1) | Alcohol and carboxylic acid | |

| Question Number | Acceptable Answer | Reject | Mark | | | | | | | | | |
|--|--|-------------|-------|---|------------|--------|---|-------------|--------|---|----------------------------------|------------|
| 1(d)(i) | <table border="1"> <tr> <td>C 60.0 ÷ 12</td> <td>= 5.0</td> <td>3</td> </tr> <tr> <td>H 13.3 ÷ 1</td> <td>= 13.3</td> <td>8</td> </tr> <tr> <td>O 26.7 ÷ 16</td> <td>= 1.67</td> <td>1</td> </tr> </table> | C 60.0 ÷ 12 | = 5.0 | 3 | H 13.3 ÷ 1 | = 13.3 | 8 | O 26.7 ÷ 16 | = 1.67 | 1 | C ₃ H ₇ OH | (2) |
| | C 60.0 ÷ 12 | = 5.0 | 3 | | | | | | | | | |
| | H 13.3 ÷ 1 | = 13.3 | 8 | | | | | | | | | |
| | O 26.7 ÷ 16 | = 1.67 | 1 | | | | | | | | | |
| | Ratio 5.0: 13.3: 1.67 (1) | | | | | | | | | | | |
| C ₃ H ₈ O (1) | | | | | | | | | | | | |
| Correct answer without working scores 2 marks. | | | | | | | | | | | | |
| Correct answer with incorrect working (e.g. mole calculations inverted) scores 1 mark. No TE on incorrect ratios. | | | | | | | | | | | | |

| Question Number | Acceptable Answer | Reject | Mark |
|-----------------|--|--------|------|
| 1(d)(ii) | $\begin{array}{ccccccc} & \text{H} & & \text{H} & & \text{H} & \\ & & & & & & \\ \text{H} & - \text{C} & - & \text{C} & - & \text{C} & - \text{O} - \text{H} \\ & & & & & & \\ & \text{H} & & \text{H} & & \text{H} & \end{array}$ <p style="text-align: right;">(1)</p> | C-H-O | |
| | $\begin{array}{ccccccc} & \text{H} & & \text{H} & & \text{H} & \\ & & & & & & \\ \text{H} & - \text{C} & - & \text{C} & - & \text{C} & - \text{H} \\ & & & & & & \\ & \text{H} & & \text{O} & & \text{H} & \\ & & & & & & \\ & & & \text{H} & & & \end{array}$ <p style="text-align: right;">(1)</p> <p>ALLOW – OH for the hydroxyl group, but bond should go from C to O in propan-1-ol.</p> <p>IGNORE poorly placed OH in propan-2-ol.</p> <p>ALLOW skeletal formulae/structural formulae both correct – 1 mark.</p> <p>ALLOW TE from 1(d)(i), e.g. if a different number of carbon atoms is given in 1di then allow 2 different isomers drawn displayed correctly.</p> | | |

Total for Question 1 = 18 Marks

| Question Number | Acceptable Answer | Reject | Mark |
|-----------------|---|---|------------|
| 2(a)(i) | $\text{Zn(s)} + \text{Cu}^{2+}(\text{aq}) \rightarrow \text{Zn}^{2+}(\text{aq}) + \text{Cu(s)}$ <p>First mark for correct species in a balanced equation.</p> <p>ALLOW hexaqua ions providing the equation is balanced.</p> <p>IGNORE reversible arrows. (1)</p> <p>Second mark for states. Consequential on a reasonable attempt at the equation, including for a full equation or unbalanced equation.</p> <p>For example, ALLOW as reasonable Zn with 1+ instead of 2+ Sulfate ions shown correctly but not cancelled out. (1)</p> | <p>Sulfate ions not cancelled out</p> <p>Reversed equation Zn and Cu metal both shown as ions</p> | (2) |

| Question Number | Acceptable Answer | Reject | Mark |
|-----------------|--|--------|------------|
| 2(a)(ii) | $q = 50.0 \times 4.18 \times 46.5$ $= 9718.5 \text{ (J)}/9.7185 \text{ kJ}$ <p>(1) (1)</p> <p>Correct answer with no working scores (2)</p> <p>IGNORE sf except 1 (i.e. allow 9719/9720 /9700 or 9.719/9.72/9.7 kJ)</p> <p>IGNORE sign of q if given.</p> <p>If mass used 55.0g and $q = 55.0 \times 4.18 \times 46.5$ $= 10690.35 \text{ (J)}$ (1)</p> <p>If mass used is 5g and $q = 5 \times 4.18 \times 46.5 = 971.85 \text{ (J)}$ (1)</p> | 9718 | (2) |

| Question Number | Acceptable Answer | Mark |
|-----------------|--|------------|
| 2(a)(iii) | $\frac{50.0 \times 1}{1000}$ $= 0.05 \text{ (mol)}$ <p>Mark is for final answer.</p> | (1) |

| Question Number | Acceptable Answer | Mark |
|-----------------|--|------------|
| 2(a)(iv) | $\Delta H = -9718.5 \div (0.050 \times 1000)$ $= -194.370 \text{ (kJ mol}^{-1}\text{)}$ $= \mathbf{-194} \text{ (kJ mol}^{-1}\text{)}$ <p>ALLOW = -194000 J mol⁻¹</p> <p>First mark:</p> <p>Value, ignore sign and sf. Only penalise units if value is in J (mol⁻¹) without stating this</p> <p style="text-align: right;">(1)</p> <p>TE (a)(ii) ÷ ((a)(iii) × 1000)</p> <p>Using 10690.5 gives - 2138810J = -214 kJ mol⁻¹.</p> <p>Second mark:</p> <p>Sign and 3 sf.</p> <p style="text-align: right;">(1)</p> <p>This mark depends on a correct calculation method.</p> | (2) |

| Question Number | Acceptable Answer | Mark |
|-----------------|-------------------|------------|
| 2(b)(i) | 47.5 (°C) | (1) |

| Question Number | Acceptable Answer | Reject | Mark |
|-----------------|--|---------------------------------------|------------|
| 2(b)(ii) | $(1.0 \times 100) \div 46.5 = 2.1505376$ $= (\pm) \mathbf{2.15/2.2/2} \text{ (\%)}$ <p>IGNORE sf.</p> <p>ALLOW answer with 47.5 in the denominator which gives 2.1052631 = (±) 2.11/2.1/2 (%).</p> <p>ALLOW TE on value (b)(i).</p> | <p>2.0 and 2.1</p> <p>2.0 and 2.2</p> | (1) |

| Question Number | Acceptable Answer | Reject | Mark |
|-----------------|---|-------------------------------------|------------|
| 2(c) | <p>First mark: Measure the temperature (of copper(II) sulfate) every minute/at realistic time interval (15 seconds to 1 minute) for, e.g., 2-4 minutes before adding zinc.</p> <p>OR measure temperature (of copper(II) sulfate) before adding zinc. (1)</p> <p>Second mark: Measure temperature each minute/at realistic time intervals (after adding zinc) for several minutes. (1)</p> <p>N.B. these readings may be started after the maximum temperature is reached/after reaction has stopped and taken until the mixture has cooled to room temperature. Intervals should be chosen to allow at least 4 readings on cooling section of curve.</p> <p>Third mark: Plot a temperature – time graph/plot a graph using measurements (of temp and time) obtained. (1)</p> <p>This mark can be awarded if first two marks are insufficient for credit.</p> <p>Fourth mark: Extrapolate to find ΔT/ maximum temperature (at the time of mixing).</p> <p>OR use properly described intersecting lines to find maximum temperature. (1)</p> <p>ALLOW third and fourth marking points to be shown on annotated diagrams/graph.</p> <p>If zinc is added in small portions or over a period of time only first and third marks can be awarded.</p> <p>(since measurements of cooling will be incorrect and there is no definite time when reaction starts.)</p> | Readings more often than every 15 s | (4) |

Total for Question 2 = 13 Marks

| Question Number | Acceptable Answer | Reject | Mark |
|-----------------|--|--|------------|
| 3(a)(i) | <p>Any two from: Misty/steamy fumes (1)</p> <p>Purple/violet fumes (1)</p> <p>ALLOW purple gas/vapour.</p> <p>Brown or black solution/liquid/solid OR grey/grey-black solid (1)</p> <p>Yellow solid/deposit (1)</p> <p>ALLOW yellow precipitate.</p> <p>IGNORE effervescence, bubbles, colour change, coloured fumes, solid disappears, description of smells, identification of products even if incorrect, follow-on tests, e.g. effect on potassium dichromate paper.</p> | <p>White fumes Steamy white</p> <p>Smoke</p> <p>Yellow fumes Yellow liquid</p> | (2) |

| Question Number | Acceptable Answer | Reject | Mark |
|-----------------|--|--|------------|
| 3(a)(ii) | <p>There is little/no HI formed (which is the reagent needed). (1)</p> <p>Because HI is oxidized (to iodine)/because iodide ions are oxidized (to iodine)/sulfuric acid is oxidizing/HI reduces sulfuric acid/iodide ions reduce sulfuric acid. (1)</p> <p>Must mention oxidation or reduction correctly for second mark.</p> <p>IGNORE 'an elimination reaction would occur'.</p> <p>ALLOW 'HI is oxidized to iodine' for both marks.</p> | <p>Iodide ions react with sulfuric acid</p> <p>Sulfuric acid oxidizes iodine</p> <p>HI is reduced to iodine</p> | (2) |

| Question Number | Acceptable Answer | Reject | Mark |
|-----------------|---|------------------------------------|------------|
| 3(b) | $2P + 3I_2 \rightarrow 2PI_3$ OR $P + 3/2I_2 \rightarrow PI_3$ OR $P_4 + 6I_2 \rightarrow 4PI_3$ ALLOW reversible sign. IGNORE state symbols even if incorrect. | Equations with ions I for I_2 | (1) |

| Question Number | Acceptable Answer | Reject | Mark |
|-----------------|--|--|------------|
| 3(c)(i) | Exothermic ALLOW fast/vigorous/violent. | Dangerous Reactive (In)flammable Volatile | (1) |

| Question Number | Acceptable Answer | Reject | Mark |
|-----------------|--|---|------------|
| 3(c)(ii) | (Very) pale purple/yellow/straw coloured OR Colourless mixture/is decolourised OR Co purple colour | Clear for colourless No (grey) solid remains Add starch | (1) |

| Question Number | Acceptable Answer | Reject | Mark |
|-----------------|---|---------------|------------|
| 3(c)(iii) | Diagram to show: Distillation flask and still-head and heat. (1) (no need for a thermometer) ALLOW appropriate tubing as alternative to still head. ALLOW heating with electrical, water bath, Bunsen or just arrow. IGNORE thermometer and position, tap funnel in still head, absence of reagents in flask Condenser sloping downwards (1) With water entering at the bottom and suitable receiver (e.g. flask or beaker) (1) | Conical flask | (3) |

| Question Number | Acceptable Answer | Reject | Mark |
|-----------------|---|---|------------|
| 3(c)(iv) | This removes/reacts with (any residual) iodine. OR Removes excess iodine/ I_3^- . | Removes acid Removes impurities Removes iodide Removes ions (other than I_3^-) Just reduces iodine to iodide | (1) |

| Question Number | Acceptable Answer | Reject | Mark |
|-----------------|---|---|------------|
| 3(c)(v) | Anhydrous calcium chloride/it is drying agent OR anhydrous salt needed to remove water/hydrated salt will not remove water Allow moisture for water and absorb for remove. | Just 'calcium chloride is a drying agent' | (1) |

| Question Number | Acceptable Answer | Reject | Mark |
|-----------------|---|--|------------|
| 3(c)(vi) | Distillation/re-distillation (over a narrow range of temperature) (either side of the boiling temperature of 1-iodobutane). ALLOW fractional distillation. IGNORE filtering before distillation and any temperatures given. | Recrystallisation Just 'purification' | (1) |

| Question Number | Acceptable Answer | Mark |
|-----------------|--|--|
| 3(d)(i) | $(95.0 \div 74.0) \times 92.5 \text{ g}$ $= 118.75/118.8/119 \text{ g}$ ALLOW 118.77 (from use of 1.284) 3, 4 or 5 sf in final answer Correct final answer scores 2 marks. OR Rounding errors by dividing $95.0 \div 74.0$ as a first step: e.g. $(95.0 \div 74.0) = 1.28$, followed by $1.28 \times 92.5 = 118.4/118$ e.g. $(95.0 \div 74.0) = 1.3$ followed by $1.3 \times 92.5 = 120.25/120.3/120$ | (1) (1) (2) (1) (1) |

| Question Number | Acceptable Answer | Reject | Mark |
|-----------------|--|--------|------------|
| 3d(ii) | $95.3 \div 3(d)(i)$ $(95.3 \div 118.75) \times 100 = 80.2563$ $= \mathbf{80.25/80.3 \%}$ Many candidates give the answer to 3 d(i) to 3sf, e.g. 119 but keep the full answer in their calculator, resulting in an answer of 80.25 which is correct and should be allowed. TE from 3(d)(i). | 80.2 | (1) |

| Question Number | Acceptable Answer | Reject | Mark |
|-----------------|---|---|------------|
| 3d(iii) | <p>One of:</p> <p>Handling/transfer losses</p> <p>Competing reactions/(unwanted) side reaction/by-products form</p> <p>Incomplete reaction</p> | <p>Just 'losses'/spillage</p> <p>Impure reagents</p> <p>Loss by evaporation</p> <p>Other products form</p> <p>Not enough PCl_5 to react</p> | (1) |

| Question Number | Acceptable Answer | Reject | Mark |
|-----------------|--|---|------|
| 3d(iv) | <p>Two of:</p> <p>Low atom economy IGNORE 'low percentage/80% yield'.</p> <p>Phosphorus(V) chloride expensive.</p> <p>Disposal of unwanted materials expensive or difficult.</p> <p>No (large scale) use for POCl_3.</p> <p>Difficult/expensive to separate required product.</p> <p>No credit for:</p> <p>Slow/time consuming</p> <p>Exothermic</p> <p>Not efficient</p> <p>High energy use</p> <p>Competing reactions</p> <p>Non-renewable reactants</p> <p>HCl toxic/acidic</p> <p>Unwanted products</p> | <p>Just Atom economy not 100%</p> <p>Just 'It' would be expensive</p> <p>Anything to do with environmental friendliness or the ozone layer or the end of life on Earth.</p> | (2) |

Total for Question 3 = 19 Marks

Total for Paper = 50 Marks