

Please check the examination details below before entering your candidate information

Candidate surname					Other names			
Pearson Edexcel		Centre Number			Candidate Number			
Level 3 GCE		<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>			<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>			
Thursday 21 May 2020								
Morning (Time: 1 hour 30 minutes)					Paper Reference 8CH0/02			
Chemistry								
Advanced Subsidiary								
Paper 2: Core Organic and Physical Chemistry								
Candidates must have: Scientific calculator							Total Marks	
Data Booklet								
Ruler								

Instructions

- Use **black** ink or **black** ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
– *there may be more space than you need.*

Information

- The total mark for this paper is 80.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*
- For the question marked with an **asterisk** (*), marks will be awarded for your ability to structure your answer logically showing the points that you make are related or follow on from each other where appropriate.
- A Periodic Table is printed on the back cover of this paper.

Advice

- Read each question carefully before you start to answer it.
- Show all your working in calculations and include units where appropriate.
- Check your answers if you have time at the end.

Turn over ►

P62307A

©2020 Pearson Education Ltd.

1/1/1/1/1/



Pearson

Answer ALL questions.

**Some questions must be answered with a cross \boxtimes .
If you change your mind about an answer, put a line through the box \boxtimes
and then mark your new answer with a cross \boxtimes .**

- 1** This question is about organic compounds containing fluorine and chlorine.
- (a) The use of chlorofluorocarbons as refrigerants has ceased due to concerns about their effects on the ozone layer. One such compound is dichlorodifluoromethane.

Give the molecular formula of dichlorodifluoromethane.

(1)

-
- (b) (i) A different refrigerant contains 34.0% chlorine and 54.5% fluorine by mass, with the remainder carbon.

Calculate the empirical formula of this compound.

(3)

DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



(ii) Use the mass spectrum to show that the empirical and the molecular formulae of this compound are the same.

(1)



(iii) Suggest the species responsible for the peak at $m/z = 69$.

(1)



(c) Compounds containing carbon and fluorine but no chlorine can be used as refrigerants as they are not harmful to the ozone layer. These can be made by the reaction of fluorine with alkanes or fluoroalkanes. A refrigerant currently in use contains the compound trifluoromethane, CHF_3 .

(i) Write the equation for the formation of trifluoromethane by the reaction of difluoromethane with fluorine. State symbols are not required. (1)

(ii) The mechanism for this reaction is similar to that of the reaction between chlorine and methane.

Give the equations for the following steps in the mechanism for the reaction between fluorine and difluoromethane. Curly arrows are not required. (3)

Initiation step

First propagation step

Second propagation step

(Total for Question 1 = 10 marks)






2 This question is about alcohols and their reactions.

The table gives some of the names and skeletal formulae of isomers having the formula C_4H_9OH .

(a) Complete the table.

(2)

Name	Skeletal formula
	
butan-2-ol	
2-methylpropan-1-ol	
2-methylpropan-2-ol	

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



(b) (i) Some alcohols react with concentrated phosphoric acid to form alkenes.

What is the type of this reaction?

(1)

- A addition
- B elimination
- C oxidation
- D substitution

(ii) When butan-2-ol reacts with concentrated phosphoric acid, two stereoisomers are formed.

Explain what is meant by the term stereoisomers.

(2)

.....

.....

.....

.....

(iii) Draw the structures and give the names of the two stereoisomers.

(2)

Stereoisomer 1	Stereoisomer 2
Name:	Name:

(iv) Name this type of stereoisomerism.

(1)



- (c) 2-methylpropan-2-ol may be formed by the reaction between 2-bromo-2-methylpropane and aqueous potassium hydroxide.

What is the role of the hydroxide ions in this reaction?

(1)

- A alkali
 B catalyst
 C electrophile
 D nucleophile

- (d) (i) $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$ reacts with the oxidising agent potassium dichromate(VI) in dilute sulfuric acid.

Two organic products can be formed, depending on the conditions.

Write a balanced equation for the formation of **one** of these products, giving its name and the condition required to achieve this product in high yield.

Use [O] in the equation to represent each oxygen atom from the oxidising agent.

(3)

Equation

Name

Condition

- (ii) The colour of the solution at the end of the reaction in (d)(i) will be

(1)

- A brown
 B green
 C orange
 D yellow

(Total for Question 2 = 13 marks)



3 This question is about reaction kinetics.

(a) The best way to describe the activation energy of a reaction is

(1)





- A the average energy of the particles when they react
- B the difference in energy between the reactants and the products
- C the minimum energy required to make the particles collide
- D the minimum energy required for a reaction to occur

(b) The diagrams show two reaction profiles for the same reversible reaction involving gaseous reactants.

Shown on each diagram are the reaction profiles for the pathway without a catalyst and the pathway catalysed by a heterogeneous catalyst.


(i) In which diagram does the arrow represent the activation energy for the backward reaction when a catalyst is present?

(1)



(ii) Estimate, using the diagram, the **decrease** in the activation energy for the forward reaction when a catalyst is added.

(1)



- A 75 kJ mol⁻¹
- B 100 kJ mol⁻¹
- C 175 kJ mol⁻¹
- D 200 kJ mol⁻¹

(c) State why a **solid** (heterogeneous) catalyst is suitable for a reaction in the **gas** phase.

(1)


.....

.....

.....



- (d) The diagram shows a Maxwell-Boltzmann distribution of molecular energies for gaseous molecules.




- (i) Which is the area of the graph corresponding to the number of molecules with sufficient energy to react when a catalyst is present? (1)
- A Y
- B Y – Z
- C Y + Z
- D Z
- (ii) Which would always result in a **decrease** in the number of molecules contained within area Y? (1)
- A decreasing the temperature of the gas
- B increasing the pressure of the gas
- C putting the gas in a smaller container
- D removing a quarter of the catalyst

(Total for Question 3 = 6 marks)



4 Methanol, CH_3OH , is a liquid fuel.

An experiment was carried out to determine the enthalpy change of combustion of liquid methanol.



The energy obtained from burning 2.08 g of methanol was used to heat 75.0 g of water.

The temperature of the water rose from 25.0°C to 91.0°C .

[Specific heat capacity of water = $4.18\text{ J g}^{-1}\text{ }^\circ\text{C}^{-1}$]

- (a) Use the data to calculate a value for the enthalpy change of combustion of one mole of methanol.

Give your answer to an appropriate number of significant figures and include a sign and units.

(4)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



(b) Methanol can be synthesised from methane and steam by a process that occurs in two steps.



(i) Explain the effects of increasing the pressure on the yield of the products and on the rate of the reaction in Step 1.

(4)

(ii) Step 2 is carried out at a compromise temperature of 500 K.

Explain why 500 K is considered to be a compromise for Step 2 by considering what would happen at higher and lower temperatures.

(3)



- (c) Calculate a value for the standard enthalpy change of combustion of gaseous methanol using the enthalpy change for Step 2 and the standard enthalpy change of combustion of gaseous carbon monoxide and of hydrogen.

Substance	Standard enthalpy change of combustion / kJ mol^{-1}
CO	-283
H ₂	-286

(3)


(Total for Question 4 = 14 marks)



- 5 This question concerns iodine monochloride, ICl, a red-brown solid which melts at 27°C to form a red-brown liquid.

Iodine monochloride is used in measuring unsaturation in organic compounds.

- (a) Iodine monochloride gas can be produced by the reaction between iodine vapour and chlorine gas. The reaction is exothermic.



The table shows bond energy values for the bonds in iodine and chlorine.

Calculate the value of the bond energy of the I—Cl bond using these data and the equation.

Bond	Energy / kJ mol^{-1}
I—I	151
Cl—Cl	243

(2)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



(b) Iodine monochloride is a polar molecule which adds rapidly to double bonds in a similar way to hydrogen chloride. This reaction can be used to determine the degree of unsaturation in oils.

(i) Add the dipole to a molecule of iodine monochloride.

(1)



(ii) Draw the mechanism for the addition of iodine monochloride to propene. You should include all curly arrows and relevant lone pairs and dipoles.

(3)


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



- (c) (i) To determine the extent of unsaturation of an oil, 0.250 g of the oil was treated with 25.00 cm³ of a 0.100 mol dm⁻³ ICl solution. Unreacted ICl reacted with excess potassium iodide solution, forming iodine according to the equation:



The amount of iodine produced was measured by reacting the mixture with a solution of sodium thiosulfate, Na₂S₂O₃.

The iodine released reacted with 32.65 cm³ of 0.100 mol dm⁻³ sodium thiosulfate solution in the mole ratio of 1 mol I₂ : 2 mol Na₂S₂O₃.

Calculate the number of moles of iodine monochloride which reacted with 0.250 g of the oil.

(3)



- (ii) Unsaturation in oils is measured using a scale called 'Iodine number'. This is the mass of iodine which will react with 100g of the oil. Because iodine adds very slowly to double bonds, the reaction of iodine monochloride is used instead.

Given that 1 mol of I_2 is equivalent to 1 mol of ICl , use your answer in (c)(i) to calculate the mass of iodine that would react with 100g of oil and hence identify the unsaturated oil from the list of possible oils and their iodine numbers.

Oil	Iodine number
cocoa butter	35–40
coconut oil	7–10
cod liver oil	145–180
palm oil	44–51
peanut oil	84–106

(2)

- (iii) Give a reason why the reaction of iodine monochloride is significantly faster than the reaction of iodine.

(1)

.....

.....

.....

(Total for Question 5 = 12 marks)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE



- 6 Aqueous hydrogen peroxide decomposes according to the following equation.




The decomposition is catalysed by manganese(IV) oxide.

This can be investigated by measuring the volume of oxygen produced at various times as the reaction proceeds. Part of the apparatus used in the experiment is shown. The manganese(IV) oxide is placed in a small glass container, which is then tipped over to start the reaction. A stop clock is started at the same time.

- (a) Complete the diagram to show how the gas can be collected **and** its volume measured, labelling the apparatus used.

(2)



DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



(b) An experiment was carried out using 0.25 g of manganese(IV) oxide granules and 50 cm³ of aqueous hydrogen peroxide of concentration 0.16 mol dm⁻³. The results are shown in the table and plotted on a graph.

Time/s	0.0	20.0	30.0	50.0	60.0	80.0	100	120	150
Volume of O ₂ /cm ³	0	51	68	85	88	91	92	92	92



- (i) The rate of reaction may be assumed to be approximately constant up to the first volume measurement (20.0 s in this experiment).

Use this approximation to calculate the initial rate of this reaction, giving the **units** with your answer.

(1)




- (ii) Draw a tangent at 40 s on the graph on Page 20 and use it to calculate the rate of reaction at this time.

(2)

- (iii) The experiment was repeated on a different day when the laboratory was 20°C warmer. The volume of oxygen was recorded for the same total time of 150 s.

Draw the line that you would expect to obtain in this experiment. Assume the pressure in the laboratory is the same. No calculation is required.

(2)



(iv) Explain, using collision theory, any differences between the line you have drawn and the original line of best fit.

(2)

DO NOT WRITE IN THIS AREA

(c) Catalysts are not used up during a reaction. Manganese(IV) oxide acts as a heterogeneous catalyst.

Describe in outline a method to show that the manganese(IV) oxide is not used up in the decomposition of hydrogen peroxide **and** that it still functions as a catalyst.

(4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 6 = 13 marks)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE



7 Halogenoalkanes react with water to produce alcohols and halide ions.



(a) Test tube experiments can be carried out to investigate the relative rates of these substitution reactions.

The halogenoalkanes 1-chlorobutane, 1-bromobutane and 1-iodobutane can be used.

Some of the steps in these experiments are

- each halogenoalkane is added to a different tube containing 1 cm^3 of ethanol
- the test tubes are placed in the same beaker of hot water
- aqueous silver nitrate is added to each tube and the tubes are shaken
- a precipitate forms in each tube.

(i) State the purpose of adding ethanol to each of the test tubes.

(1)

(ii) Give **one** reason why the test tubes were put in the same beaker of hot water.

(1)

(iii) Give **one** reason why the test tubes were shaken after the addition of aqueous silver nitrate.

(1)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



(b) (i) State how the halogen atom present in each halogenoalkane can be identified using observations from this experiment in (a).

(1)

.....

.....

.....

.....

.....

.....

(ii) Identify further reagents that can be added, including relevant observations, to confirm the identity of the halogen atom present in each halogenoalkane.

(2)

.....

.....

.....

.....

.....

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 7 = 12 marks)

TOTAL FOR PAPER = 80 MARKS



The Periodic Table of Elements

1 2 3 4 5 6 7 0 (8) (18)

1.0	H
	hydrogen
	1

Key

relative atomic mass
atomic symbol
name
atomic (proton) number

(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)
6.9	9.0	45.0	47.9	50.9	52.0	54.9	55.8	58.9	58.7	63.5	65.4	10.8	12.0	14.0	16.0	19.0	4.0
Li	Be	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	B	C	N	O	F	He
lithium	beryllium	scandium	titanium	vanadium	chromium	manganese	iron	cobalt	nickel	copper	zinc	boron	carbon	nitrogen	oxygen	fluorine	helium
3	4	21	22	23	24	25	26	27	28	29	30	5	6	7	8	9	2
23.0	24.3	88.9	91.2	92.9	95.9	[98]	101.1	102.9	106.4	107.9	112.4	27.0	28.1	31.0	32.1	35.5	39.9
Na	Mg	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	Al	Si	P	S	Cl	Ar
sodium	magnesium	yttrium	zirconium	niobium	molybdenum	technetium	ruthenium	rhodium	palladium	silver	cadmium	aluminium	silicon	phosphorus	sulfur	chlorine	argon
11	12	39	40	41	42	43	44	45	46	47	48	13	14	15	16	17	18
39.1	40.1	88.9	91.2	92.9	95.9	[98]	101.1	102.9	106.4	107.9	112.4	27.0	28.1	31.0	32.1	35.5	39.9
K	Ca	La*	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	Ga	Ge	As	Se	Br	Kr
potassium	calcium	lanthanum	hafnium	tantalum	tungsten	rhenium	osmium	iridium	platinum	gold	mercury	gallium	germanium	arsenic	selenium	bromine	krypton
19	20	57	72	73	74	75	76	77	78	79	80	31	32	33	34	35	36
85.5	87.6	138.9	178.5	180.9	183.8	186.2	190.2	192.2	195.1	197.0	200.6	69.7	72.6	74.9	79.0	79.9	83.8
Rb	Sr	Ba	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	In	Sn	Sb	Te	I	Xe
rubidium	strontium	barium	hafnium	tantalum	tungsten	rhenium	osmium	iridium	platinum	gold	mercury	indium	tin	antimony	tellurium	iodine	xenon
37	38	56	72	73	74	75	76	77	78	79	80	49	50	51	52	53	54
132.9	137.3	138.9	178.5	180.9	183.8	186.2	190.2	192.2	195.1	197.0	200.6	114.8	118.7	121.8	127.6	126.9	131.3
Cs	Ba	La*	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	Pb	Bi	Po	At	Rn	Rn
caesium	barium	lanthanum	hafnium	tantalum	tungsten	rhenium	osmium	iridium	platinum	gold	mercury	lead	bismuth	polonium	astatine	radon	radon
55	56	57	72	73	74	75	76	77	78	79	80	82	83	84	85	86	86
[223]	[226]	[227]	[261]	[262]	[266]	[264]	[277]	[268]	[271]	[272]	[272]	204.4	207.2	209.0	[210]	[222]	[222]
Fr	Ra	Ac*	Rf	Db	Sg	Bh	Hs	Mt	Ds	Rg	Rg	Tl	Pb	Bi	Po	At	Rn
francium	radium	actinium	rutherfordium	dubnium	seaborgium	bohrium	hassium	meitnerium	darmstadtium	roentgenium	roentgenium	thallium	lead	bismuth	polonium	astatine	radon
87	88	89	104	105	106	107	108	109	110	111	111	81	82	83	84	85	86

Elements with atomic numbers 112-116 have been reported but not fully authenticated

140	141	144	150	152	157	163	165	167	169	173	175
Ce	Pr	Nd	Sm	Eu	Gd	Dy	Ho	Er	Tm	Yb	Lu
cerium	praseodymium	neodymium	samarium	europium	gadolinium	dysprosium	holmium	erbium	thulium	ytterbium	lutetium
58	59	60	62	63	64	66	67	68	69	70	71
232	[231]	238	[242]	[243]	[247]	[251]	[254]	[253]	[256]	[254]	[257]
Th	Pa	U	Pu	Am	Cm	Cf	Es	Fm	Md	No	Lr
thorium	protactinium	uranium	plutonium	americium	curium	californium	einsteinium	fermium	mendeleevium	nobelium	lawrencium
90	91	92	94	95	96	98	99	100	101	102	103

* Lanthanide series

* Actinide series

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

