

Please check the examination details below before entering your candidate information

Candidate surname					Other names				
Centre Number					Candidate Number				

Pearson Edexcel Level 3 GCE

Monday 12 June 2023

Morning (Time: 1 hour 45 minutes)

Paper reference **9CH0/01**

Chemistry

Advanced

PAPER 1: Advanced Inorganic and Physical Chemistry

You must have:
Scientific calculator, Data Booklet, ruler

Total Marks

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
– *there may be more space than you need.*

Information

- The total mark for this paper is 90.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*
- For the question marked with an **asterisk** (*), marks will be awarded for your ability to structure your answer logically, showing the points that you make are related or follow on from each other where appropriate.
- A Periodic Table is printed on the back cover of this paper.

Advice

- Read each question carefully before you start to answer it.
- Show all your working in calculations and include units where appropriate.
- Check your answers if you have time at the end.

Turn over ►

P71912A

©2023 Pearson Education Ltd.

N:1/1/1/1/1/1/1/





Pearson

Answer ALL questions.

Some questions must be answered with a cross in a box ☐. If you change your mind about an answer, put a line through the box ☒ and then mark your new answer with a cross ☐.

1 This question is about d-block elements.

(a) Which of the labels 1–4 identifies a d-block element in Period 4?



- ☒ **A** 1
- ☒ **B** 2
- ☒ **C** 3
- ☒ **D** 4

(1)

(b) State what is meant by the term d-block element.

(1)

.....

.....

.....

.....

(Total for Question 1 = 2 marks)



2 Chemists often use the term 'orbital' when considering atomic structure.

(a) State what is meant by the term orbital.

(2)

.....

.....

.....

.....

.....

(b) Draw diagrams to show the shape of an s and a p orbital.

(2)

s orbital	p orbital

(c) What is the electronic configuration of a copper atom?

(1)

- ☐ A $[\text{Ar}]4s^13d^{10}$
- ☐ B $[\text{Ar}]4s^23d^9$
- ☐ C $[\text{Ar}]4s^24p^13d^8$
- ☐ D $[\text{Ar}]4s^24p^23d^7$

(Total for Question 2 = 5 marks)



3 This question is about compounds containing elements from Group 7.

(a) Which change occurs when concentrated sulfuric acid is added to potassium bromide?

(1)

- ☐ **A** bromide ions oxidise sulfuric acid forming sulfur
- ☐ **B** bromide ions oxidise sulfuric acid forming sulfur dioxide
- ☐ **C** bromide ions reduce sulfuric acid forming sulfur
- ☐ **D** bromide ions reduce sulfuric acid forming sulfur dioxide

(b) Chemists can test for the presence of bromide ions in solution by adding a small amount of acidified silver nitrate solution.
The solubility of the precipitate in aqueous ammonia is then tested.

(i) Which statement is correct for bromide ions?

(1)

- ☐ **A** a white precipitate forms that dissolves in concentrated ammonia only
- ☐ **B** a white precipitate forms that dissolves in both dilute and concentrated ammonia
- ☐ **C** a cream precipitate forms that dissolves in concentrated ammonia only
- ☐ **D** a cream precipitate forms that dissolves in both dilute and concentrated ammonia

(ii) Give a reason why the silver nitrate must be acidified.

(1)

(iii) Explain which acid needs to be used to acidify the silver nitrate solution and why other acids are unsuitable.

(2)



- (c) Iodine trichloride forms a dimer, I_2Cl_6 , in the solid state.
When molten, it is suggested that it breaks down as shown.



- (i) Draw a labelled diagram of a simple experiment to confirm this dissociation has occurred, stating the positive result.

(2)

Result

- (ii) What is the shape of the ICl_4^- ion?

(1)

- ☐ **A** octahedral
☐ **B** square planar
☐ **C** tetrahedral
☐ **D** trigonal bipyramidal

- (iii) The equilibrium position for the dissociation of molten I_2Cl_6 lies to the left.



What is the most likely numerical value of K_c for this equilibrium?


(1)

- ☐ **A** 1.0×10^6
☐ **B** 5.0×10^3
☐ **C** 1.0
☐ **D** 5.0×10^{-3}

(Total for Question 3 = 9 marks)



4 The graph shows the melting temperatures of some elements in Period 3.



Explain the variations in melting temperature across the period in terms of the structure and bonding in these elements.

(6)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 4 = 6 marks)



- 5 This question is about the decomposition of dinitrogen tetroxide.
The reaction eventually reaches equilibrium.




- (a) The table shows values of ΔG at different temperatures for this reaction.

Temperature / K	$\Delta G / \text{kJ mol}^{-1}$
350	-4.0
400	-13
450	-22
500	-31
550	-40

- (i) Plot a graph of ΔG against temperature.

(2)

$\Delta G / \text{kJ mol}^{-1}$



Temperature / K



- (ii) Calculate the entropy change of the system, ΔS_{system} , in $\text{JK}^{-1} \text{mol}^{-1}$, using your straight line from the graph in (a)(i) and the equation shown.

$$\Delta G = -T\Delta S_{\text{system}} + \Delta H \quad (3)$$


- (iii) What feature of the graph in (a)(i) gives the enthalpy change of the reaction? (1)

- ☐ **A** intercept of the x-axis
- ☐ **B** (intercept of the x-axis) $\times -1$
- ☐ **C** intercept of the y-axis
- ☐ **D** (intercept of the y-axis) $\times -1$



- (b) What happens to the position of the equilibrium and the colour of the mixture when the pressure is increased?

The volume of the system remains constant.




(1)

		Position of equilibrium	Change in colour
☐	A	moves to the right	mixture gets lighter
☐	B	moves to the right	mixture gets darker
☐	C	moves to the left	mixture gets lighter
☐	D	moves to the left	mixture gets darker

(Total for Question 5 = 7 marks)



6 Benzoic acid is a weak acid found in cranberries.




$\text{C}_6\text{H}_5\text{COOH}$ – benzoic acid

- (a) Which of these answers identifies the types of species present when benzoic acid is mixed with nitric acid?

$[K_a \text{ of benzoic acid} = 6.3 \times 10^{-5} \text{ mol dm}^{-3}; \quad K_a \text{ of nitric acid} = 40 \text{ mol dm}^{-3}]$

(1)



- | | | | | | |
|--------------------------|----------|------|------|------|------|
| ☐ | A | acid | base | acid | base |
| ☐ | B | acid | base | base | acid |
| ☐ | C | base | acid | base | acid |
| ☐ | D | base | acid | acid | base |

- (b) The ionic salts sodium benzoate and potassium benzoate are both used as food preservatives.

Explain why the melting temperature of sodium benzoate is higher than the melting temperature of potassium benzoate.

(2)

.....

.....

.....

.....

.....

.....

.....

.....

(c) The value of K_a for benzoic acid = $6.28 \times 10^{-5} \text{ mol dm}^{-3}$.

(i) Write the expression for the acid dissociation constant, K_a , of benzoic acid.


(1)

(ii) Calculate the mass of benzoic acid needed to prepare 250 cm^3 of a solution with a $\text{pH} = 3.51$

(4)



(d) Weak acids such as benzoic acid can be neutralised by sodium hydroxide solution.



- (i) Which of these could be used to show the end-point of a titration of benzoic acid with sodium hydroxide solution?

(1)

- ☐ **A** bromothymol blue
- ☐ **B** litmus
- ☐ **C** methyl orange
- ☐ **D** phenolphthalein

- (ii) Another weak acid found in cranberries is quinic acid, $\text{C}_6\text{H}_7(\text{OH})_4\text{COOH}$. It is neutralised by sodium hydroxide solution in a similar way to benzoic acid.

A 25.0 cm^3 sample of 0.500 mol dm^{-3} quinic acid solution was neutralised under standard conditions in a polystyrene cup using 25.0 cm^3 of 0.800 mol dm^{-3} of sodium hydroxide solution.

This resulted in a temperature rise of 2.9°C .

Calculate the standard enthalpy change of neutralisation, $\Delta_{\text{neut}}H^\ominus$, of quinic acid in kJ mol^{-1} .

[Assume the density of both solutions is 1.0 g cm^{-3} .

specific heat capacity of solution formed = $4.18\text{ J g}^{-1}\text{ }^\circ\text{C}^{-1}$]

(3)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 7 1 9 1 2 A 0 1 3 2 8

- (iii) The standard enthalpy change of neutralisation of the weak acid HCN by sodium hydroxide is $-11.7 \text{ kJ mol}^{-1}$ while that of the strong acid HCl is $-57.9 \text{ kJ mol}^{-1}$.

Explain the difference between these values.

(2)

(Total for Question 6 = 14 marks)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



7 This question is about chromium and chromium compounds.

(a) Naturally occurring chromium has four isotopes, ^{50}Cr , ^{52}Cr , ^{53}Cr and ^{54}Cr .

State what is meant by the term isotopes.

(2)

(b) Both chromium and calcium can form ions with a +2 charge.

(i) Complete the electronic configuration of a Cr^{2+} ion.

(1)

$1s^2$

(ii) Explain which of chromium or calcium most easily forms a +2 ion using all of the data in the table.

Element	Atomic number	1st ionisation energy / kJ mol^{-1}	2nd ionisation energy / kJ mol^{-1}	Metallic radius / nm
Chromium	24	653	1592	0.129
Calcium	20	590	1145	0.197

(3)



P 7 1 9 1 2 A 0 1 5 2 8

(c) Chromium(III) sulfate, $\text{Cr}_2(\text{SO}_4)_3$, dissolves in water to form the complex ion $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}(\text{aq})$.

(i) State the colour of this complex ion.

(1)

(ii) Explain why the aqueous solution of this complex ion has an acidic pH by considering the interaction between the metal ion and the ligands.

(2)



- (d) A student researching the role of dichromate(VI) ions, $\text{Cr}_2\text{O}_7^{2-}$, as an oxidising agent made the statement shown.

'Standard electrode potential data shows that it is never feasible for a 1.00mol dm^{-3} solution of potassium dichromate(VI) to oxidise the chloride ions in hydrochloric acid.'

Comment on this statement using the data and equilibria shown.

Equilibrium 1



Equilibrium 2



(4)

(Total for Question 7 = 13 marks)



***8** Transition metals and their compounds can act as catalysts in many reactions such as the ones shown:

- platinum, Pt, in the catalytic converters of vehicles
- manganese(II) ions, $\text{Mn}^{2+}(\text{aq})$, in the oxidation of ethanedioate ions, $\text{C}_2\text{O}_4^{2-}(\text{aq})$, by manganate(VII) ions, $\text{MnO}_4^{-}(\text{aq})$.

Compare and contrast the role of the catalysts in these reactions.

(6)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA


(Total for Question 8 = 6 marks)



P 7 1 9 1 2 A 0 1 9 2 8

9 This question is about silver compounds.

- (a) The diagram shows a Born–Haber cycle for the formation of silver(I) oxide, Ag_2O . All quantities are measured in kJ mol^{-1} .



- (i) Complete the diagram by adding appropriate species and state symbols to the empty boxes. (2)
- (ii) Explain why the value for the first electron affinity of oxygen is negative and the value for the second electron affinity is positive. (3)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



- (iii) Calculate a value for the standard enthalpy change of atomisation of silver, $\Delta_{\text{at}}H^{\ominus}$, using the Born–Haber cycle.

(3)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA






(c) The amount of dissolved oxygen in a sample of river water was found using the process outlined.


- excess alkaline manganese(II) sulfate, MnSO_4 , was added to a 150 cm^3 sample of river water
- the Mn^{2+} ions reacted with the dissolved oxygen forming a precipitate of manganese(IV) oxide hydroxide




- the precipitate was then dissolved using excess sulfuric acid, forming $\text{Mn}^{4+}(\text{aq})$ ions



- excess potassium iodide solution was then added, forming iodine



- the liberated iodine was then titrated with sodium thiosulfate solution, $\text{Na}_2\text{S}_2\text{O}_3(\text{aq})$, of concentration $0.00518\text{ mol dm}^{-3}$



- the mean volume of the titre of $\text{Na}_2\text{S}_2\text{O}_3(\text{aq})$ was 34.20 cm^3 .

- (i) Calculate the concentration of dissolved oxygen in the sample of river water, in g dm^{-3} .

(5)



You may use this space to continue your answer to 10(c)(i).

- (ii) The concentration of oxygen in water is often expressed in parts per million (ppm), where 1 ppm equals 1 g of solute dissolved in 1×10^6 g of solvent.

Calculate the concentration of the oxygen in the sample of river water in ppm.
Assume the density of the river water is 1.00 g cm^{-3} .

(1)



(d) Some data is shown for electrode systems involving the $\text{Mn}^{3+}(\text{aq})$ ion.

Half-cell	Electrode system	$E^\ominus / \text{V}$
A	$\text{MnO}_2(\text{s}) + 4\text{H}^+(\text{aq}) + \text{e}^- \rightleftharpoons \text{Mn}^{3+}(\text{aq}) + 2\text{H}_2\text{O}(\text{l})$	+0.95
B	$\text{Mn}^{3+}(\text{aq}) + \text{e}^- \rightleftharpoons \text{Mn}^{2+}(\text{aq})$	+1.51

Explain why Mn^{3+} ions are unstable in aqueous solution.
Include an equation and the type of reaction that occurs.

(4)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(Total for Question 10 = 15 marks)

TOTAL FOR PAPER = 90 MARKS



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE



