


OCR

Oxford Cambridge and RSA

A Level Chemistry A**H432/01** Periodic table, elements and physical chemistry**Tuesday 5 June 2018 – Afternoon****Time allowed: 2 hours 15 minutes****You must have:**

- the Data Sheet for Chemistry A (sent with general stationery)

You may use:

- a scientific or graphical calculator



First name

Last name

Centre
numberCandidate
number**INSTRUCTIONS**

- Use black ink. You may use an HB pencil for graphs and diagrams.
- Complete the boxes above with your name, centre number and candidate number.
- Answer **all** the questions.
- Write your answer to each question in the space provided. If additional space is required, use the lined page(s) at the end of this booklet. The question number(s) must be clearly shown.
- Do **not** write in the barcodes.

INFORMATION

- The total mark for this paper is **100**.
- The marks for each question are shown in brackets [].
- Quality of extended responses will be assessed in questions marked with an asterisk (*).
- This document consists of **32** pages.

SECTION A

You should spend a maximum of 20 minutes on this section.

Write your answer to each question in the box provided.

Answer **all** the questions.

- 1 A sample of boron contains the isotopes ^{10}B and ^{11}B .
The relative atomic mass of the boron sample is 10.8.

What is the percentage of ^{11}B atoms in the sample of boron?

- A 8.0%
- B 20%
- C 80%
- D 92%



$$\left(\frac{10 \times 20}{100}\right) + \left(\frac{11 \times 80}{100}\right) = 10.8$$

Your answer


C

relative atomic mass of isotopes $\left(\frac{\text{mass number} \times \%}{100}\right) + (\dots)$ [1]

- 2 In the compound $[\text{ICl}_2]^+ [\text{SbCl}_6]^-$, the oxidation number of chlorine is -1 .

What are the oxidation numbers of I and Sb in the compound?

	I	Sb
A	+1	+5
B	+1	+7
C	+3	+5
D	+3	+7



Your answer

C

[1]

- 3 What is the number of hydrogen atoms in 0.125 mol of $\text{C}_2\text{H}_5\text{OH}$?

- A 7.525×10^{22}
- B 4.515×10^{23}
- C 3.7625×10^{23}
- D 3.612×10^{24}

$0.125 \times 6 = 0.75 \text{ mol}$ in compound
 $0.75 \times 6.02 \times 10^{23} = 4.515 \times 10^{23}$ H atoms

Your answer

B

[1]

3

- 4 A student titrates a standard solution of barium hydroxide, $\text{Ba}(\text{OH})_2$, with nitric acid, HNO_3 .
 25.00 cm^3 of $0.0450 \text{ mol dm}^{-3}$ $\text{Ba}(\text{OH})_2$ are needed to neutralise 23.35 cm^3 of $\text{HNO}_3(\text{aq})$.

What is the concentration, in mol dm^{-3} , of the nitric acid?

- A 0.0241
- B 0.0482
- C 0.0900
- D 0.0964



$$25 \times 10^{-3} \times 0.045 = 1.125 \times 10^{-3} \text{ mol of Ba}(\text{OH})_2$$

$$1.125 \times 10^{-3} \times 2 = 2.25 \times 10^{-3} \text{ mol of HNO}_3$$

Your answer

D

$$\frac{2.25 \times 10^{-3}}{23.35 \times 10^{-3}} = 0.0964 \text{ mol dm}^{-3}$$

[1]

- 5 Which statement best explains why nitrogen has a larger first ionisation energy than oxygen?

- A N atoms have less repulsion between p-orbital electrons than O atoms.
generally a high nuclear charge means higher 1st ionisation energy due to more nuclear attraction
- B N atoms have a smaller nuclear charge than O atoms.
more repulsion
- C N atoms lose an electron from the 2s subshell, while O atoms lose an electron from the 2p subshell.
both N and O lose electrons from 2p subshell
- D N atoms have an odd number of electrons, while O atoms have an even number.
trend: ionisation energies across periodic table

Your answer

A


so odd/even number of electrons has no effect

[1]

- 6 In the Periodic Table, element X is in Group 2 and element Y is in Group 15 (5).

What is the likely formula of an ionic compound of X and Y?

- A X_2Y_5
- B X_2Y_3
- C X_3Y_2
- D X_5Y_2



Your answer

C

[1]

4

7 Which statement about ammonium carbonate is **not** correct?

- ~~A~~ It reacts with $\text{Ba}(\text{NO}_3)_2(\text{aq})$ to form a white precipitate. *forms BaCO_3 a white ppt*
- ~~B~~ It effervesces with dilute nitric acid. *CO_2 produced which forms gas bubbles known as effervescing*
- ~~C~~ It release an alkaline gas with warm $\text{NaOH}(\text{aq})$. *NH_3 produced (weak alkali)*
- D** It has the formula NH_4CO_3 . *correct formula: $(\text{NH}_4)_2\text{CO}_3$*


Your answer

 A
 B
 C
 D

[1]

8 A reaction is first order with respect to a reactant X.

Which rate–concentration graph for reactant X is the correct shape?




Your answer

 A
 B
 C
 D

[1]

5

- 9 The reversible reaction of sulfur dioxide and oxygen to form sulfur trioxide is shown below.



An equilibrium mixture contains 2.4 mol SO_2 , 1.2 mol O_2 and 0.4 mol SO_3 .
The total pressure is 250 atm.

What is the **partial pressure** of SO_3 ?

A 15 atm

B 25 atm

C 100 atm

D 200 atm

\rightarrow mole fraction \times total pressure

$$\frac{0.4}{2.4 + 1.2 + 0.4} = 0.1 = \text{mole fraction}$$

$$0.1 \times 250 = 25 \text{ atm}$$

Your answer

B

[1]

- 10 A buffer solution is prepared by mixing 200 cm^3 of 2.00 mol dm^{-3} propanoic acid, $\text{CH}_3\text{CH}_2\text{COOH}$, with 600 cm^3 of 1.00 mol dm^{-3} sodium propanoate, $\text{CH}_3\text{CH}_2\text{COONa}$.

K_a for $\text{CH}_3\text{CH}_2\text{COOH} = 1.32 \times 10^{-5} \text{ mol dm}^{-3}$

What is the **pH** of the buffer solution?

A 4.58

B 4.70

C 5.06

D 5.18

$$K_a = \frac{[\text{A}^-][\text{H}^+]}{[\text{HA}]}$$

$$\text{pH} = -\log_{10}[\text{H}^+]$$

$$[\text{A}^-] = \frac{600 \times 10^{-3} \times 1}{800 \times 10^{-3}} = 7.5 \times 10^{-3} \text{ mol dm}^{-3}$$

\uparrow mol = vol (dm^3) \times concentration (mol dm^{-3})

$$[\text{HA}] = \frac{200 \times 10^{-3} \times 2}{800 \times 10^{-3}} = 5 \times 10^{-3} \text{ mol dm}^{-3}$$

\uparrow total volume

$$1.32 \times 10^{-5} = \frac{[7.5 \times 10^{-3}][\text{H}^+]}{[5 \times 10^{-3}]}$$

$$[\text{H}^+] = 8.8 \times 10^{-6} \text{ mol dm}^{-3}$$

$$\text{pH} = -\log_{10}[8.8 \times 10^{-6}] = 5.06 \text{ (3sf.)}$$

Your answer

C


[1]

6

11 The table below shows standard entropies, S^\ominus .

Substance	CO(g)	H ₂ (g)	CH ₃ OH(l)
$S^\ominus/\text{J mol}^{-1}\text{K}^{-1}$	197.6	130.6	239.7

What is the entropy change, ΔS^\ominus , in $\text{J mol}^{-1}\text{K}^{-1}$, for the following reaction?



A -219.1

B -88.5

C +88.5

D +219.1

P-r
products entropy
change - reactants
entropy
change

$$239.7 - (197.6 + (2 \times 130.6))$$

$$= -219.1 \text{ J mol}^{-1}\text{K}^{-1}$$

Your answer






A

[1]

12 The redox equilibria for a hydrogen–oxygen fuel cell in alkaline solution are shown below.



What is the equation for the overall cell reaction?



Your answer

D

[1]

7

13 Which enthalpy change(s) is/are **endothermic?** *+ve*

- 1 The bond enthalpy of the C–H bond ✓ *always +ve*
- 2 The **second** electron affinity of oxygen ✓ *2nd = +ve*
1st = -ve
 $O^-(g) + e^- \rightarrow O^{2-}(g)$
- ~~3~~ The standard enthalpy change of formation of magnesium = 0

- A 1, 2 and 3
- B Only 1 and 2
- C Only 2 and 3
- D Only 1

Your answer

B

[1]

14 Which statement(s) explain(s) why reaction rates increase as temperature increases?

- ~~1~~ The activation energy is less. *activation energy only changes with the addition of a catalyst*
- 2 Collisions between molecules are more frequent.
- 3 A greater proportion of molecules have energy greater than the activation energy.

- A 1, 2 and 3
- B Only 1 and 2
- C Only 2 and 3
- D Only 1

Your answer

C

[1]

8

15 Which statement(s) is/are correct for the complex $\text{Pt}(\text{NH}_3)_2\text{Cl}_2$

1 One of its stereoisomers is used as an anti-cancer drug. *recall*

~~2~~ It has bond angles of 109.5° . *1 lone pair on N*

~~3~~ It has optical isomers.

4 bonded pairs + 0 lone pairs

A 1, 2 and 3

B Only 1 and 2

C Only 2 and 3

D Only 1

*4 different groups
2 groups and 2 of each
so not an optical isomer*

Your answer

D

[1]

SECTION B

Answer **all** the questions.

16 This question is about enthalpy changes.

(a) **Table 16.1** shows enthalpy changes that can be used to determine the enthalpy change of hydration of fluoride ions, F^- .

aqueous / hydrated ions

Enthalpy change	Energy / kJ mol^{-1}
Hydration of Ca^{2+}	-1609
Solution of CaF_2	+13
Lattice enthalpy of CaF_2	-2630

Table 16.1


(i) Explain what is meant by the term *enthalpy change of hydration*.

1 mole of gaseous ions react to form 1 mole of aqueous / hydrated ions

[2]

(ii) The enthalpy change of hydration of F^- can be determined using the enthalpy changes in **Table 16.1** and the incomplete energy cycle below.

On the dotted lines, add the species present, including state symbols.



[4]

11

- (iii) Calculate the enthalpy change of hydration of fluoride ions, F^- .

$$2x = +1609 - 2630 + 13$$

$$2x = -1008$$

$$x = -504$$

enthalpy change of hydration = -504 kJ mol^{-1} [2]


- (iv) Predict how the enthalpy changes of hydration of F^- and Cl^- would differ.

Explain your answer.

$\Delta_{\text{hyd}}H$ of F^- is more exothermic
because of F^- 's smaller size
meaning greater attraction to
 H_2O . [2]

12

(b) Fluorine reacts with steam as shown in the equation below.



Average bond enthalpies are shown in the table. ↳ reactants - products = ΔH
bond enthalpies of ↑

Bond	Average bond enthalpy / kJ mol ⁻¹
O-H	+464
O=O	+498
H-F	+568

(i) Explain what is meant by the term average bond enthalpy. recall - learn this definition

The breaking of 1 mole of
bonds in gaseous molecules

[2]

(ii) Calculate the bond enthalpy of the F-F bond.

$$(2x + 2(2 \times 464)) - (498 + (4 \times 568)) = -598$$

$$(2x + 1856) - 2770 = -598$$

$$2x + 1856 = 2172$$


$$2x = 316$$

$$x = +158$$

bond enthalpy = + 158 kJ mol⁻¹ [3]

17 This question is about reaction rates.

Aqueous iron(III) ions, $\text{Fe}^{3+}(\text{aq})$, react with aqueous iodide ions, $\text{I}^{-}(\text{aq})$, as shown below.



A student carries out three experiments to investigate how different concentrations of $\text{Fe}^{3+}(\text{aq})$ and $\text{I}^{-}(\text{aq})$ affect the initial rate of this reaction. The results are shown below.

Experiment	$[\text{Fe}^{3+}(\text{aq})]$ / mol dm^{-3}	$[\text{I}^{-}(\text{aq})]$ / mol dm^{-3}	Initial rate / $\text{mol dm}^{-3} \text{s}^{-1}$
1	4.00×10^{-2}	3.00×10^{-2}	8.10×10^{-4}
2 $\times 1$	8.00×10^{-2}	3.00×10^{-2}	1.62×10^{-3}
3	4.00×10^{-2}	6.00×10^{-2}	3.24×10^{-3}


(a)* Determine the rate constant and a possible two-step mechanism for this reaction that are consistent with these results. [6]

$$\text{rate} = k[\text{Fe}^{3+}][\text{I}^{-}]^2$$

$$8.10 \times 10^{-4}$$

$$k = \frac{8.10 \times 10^{-4}}{(4 \times 10^{-2})(3 \times 10^{-2})^2} = 22.5 \text{ mol}^{-2} \text{ dm}^6 \text{ s}^{-1}$$

$$\frac{\text{mol dm}^{-3} \text{ s}^{-1}}{(\text{mol dm}^{-3})^3} = \frac{\text{mol dm}^{-3} \text{ s}^{-1}}{\text{mol}^2 \text{ dm}^{-9} \text{ s}^{-1}}$$



overall equation:



Additional answer space if required

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

- (b) A student carries out an investigation to find the activation energy, E_a , and the pre-exponential factor, A , of a reaction.

$$\ln k = -\frac{E_a}{RT} + \ln A$$

The student determines the rate constant, k , at different temperatures, T .
The student then plots a graph of $\ln k$ against $1/T$ as shown below.



17

- (i) Draw a best-fit straight line and calculate the activation energy, in J mol^{-1} .
Give your answer to three significant figures.

Show your working.

$$\text{gradient: } \frac{29 - 28}{2.9 \times 10^{-3} - 4 \times 10^{-3}} = -909 = \frac{-E_a}{R}$$

$$E_a = +909 \times 8.314 = 7557.4 \\ = 7560 \text{ (3sf.)}$$

activation energy, $E_a = + \dots\dots\dots 7560 \dots\dots\dots \text{J mol}^{-1}$ [3]

- (ii) Use the graph to calculate the value of the pre-exponential factor, A .

y intercept = $\ln A$


Show your working.

$$\ln A = 31.4$$

$$A = e^{31.4} = 4.33 \times 10^{13}$$

pre-exponential factor, $A = \dots\dots\dots 4.33 \times 10^{13} \dots\dots\dots$ [2]

- 18 Nitrogen monoxide, NO, and oxygen, O₂, react to form nitrogen dioxide, NO₂, in the reversible reaction shown in **equilibrium 18.1**.



Equilibrium 18.1

- (a) Write an expression for K_c for this equilibrium and state the units. *all parts in same state included*

$$K_c = \frac{[\text{NO}_2]^2}{[\text{NO}]^2 [\text{O}_2]}$$

molar ratio in question

[] = concentration (mol dm⁻³)²

$$\text{Units} = \frac{\text{mol}^{-1} \text{dm}^3}{(\text{mol dm}^{-3})^3} = \frac{1}{\text{mol dm}^{-3}} \quad [2]$$

- (b) A chemist mixes together nitrogen and oxygen and pressurises the gases so that their total gas volume is **4.0 dm³**.

- The mixture is allowed to reach equilibrium at constant temperature and volume.
- The equilibrium mixture contains **0.40 mol NO** and **0.80 mol O₂**.
- Under these conditions, the numerical value of K_c is **45**.

Calculate the amount, in mol, of NO₂ in the equilibrium mixture.



$$\frac{0.4}{4} = 0.1 \text{ mol dm}^{-3} = [\text{NO}]$$

$$\frac{0.8}{4} = 0.2 \text{ mol dm}^{-3} = [\text{O}_2]$$

$$45 = \frac{[\text{NO}_2]^2}{[0.1]^2 [0.2]}$$

$$[\text{NO}_2]^2 = 45 \times 0.1^2 \times 0.2 = 0.09$$


$$[\text{NO}_2] = \sqrt{0.09} = 0.3 \text{ mol dm}^{-3}$$

$$0.3 \times 4 = 1.2 \text{ mol}$$

opposite of first step

amount of NO₂ = 1.2 mol [4]

- (c) The values of K_p for equilibrium 18.1 at 298 K and 1000 K are shown below.



Temperature / K	K_p / atm^{-1}
298	$K_p = 2.19 \times 10^{12}$
1000	$K_p = 2.03 \times 10^{-1}$

- (i) Predict, with a reason, whether the forward reaction is exothermic or endothermic.

K_p decreases as temperature increases so exothermic [1]

- (ii) The chemist **increases the pressure** of the equilibrium mixture at the same temperature. State, and explain in terms of K_p , how you would expect the equilibrium position to change.

Equilibrium position will shift to the right so ratio in K_p expression decreases so ratio increases to restore K_p .

Handwritten notes: equilibrium shifts to side with fewer moles of gas; denominator increases more than numerator

[3]

19 This question is about acids and bases found in the home.

(a) Ethanoic acid, CH₃COOH, is the acid present in vinegar.

A student carries out an experiment to determine the pK_a value of CH₃COOH.

- The concentration of CH₃COOH in the vinegar is 0.870 mol dm⁻³.
- The pH of the vinegar is 2.41.

(i) Write the expression for the acid dissociation constant, K_a, of CH₃COOH.

$$K_a = \frac{[H^+][CH_3COO^-]}{[CH_3COOH]} \quad \rightarrow \quad \frac{[H^+][A^-]}{[HA]} \quad [1]$$

(ii) Calculate the pK_a value of CH₃COOH.

Give your answer to two decimal places.

$$[H^+] = 10^{-pH} = 10^{-2.41} = 3.89 \times 10^{-3} \text{ mol dm}^{-3}$$

assume $[H^+] = [A^-]$ because all H⁺ ions must come from ethanoic acid

$$K_a = \frac{[3.89 \times 10^{-3}]^2}{[0.870]} = 1.74 \times 10^{-5}$$

$$pK_a = -\log_{10} 1.74 \times 10^{-5} = 4.76$$

pK_a = 4.76 [3]

(iii) Determine the percentage dissociation of ethanoic acid in the vinegar.

Give your answer to three significant figures.

$$\% \text{ dissociation} = \frac{[H^+]}{[HA]} \times 100$$

$$\frac{[3.89 \times 10^{-3}]}{[0.870]} \times 100 = 0.447\%$$

percentage dissociation = 0.447 % [1]

(b) Many solid drain cleaners are based on sodium hydroxide, NaOH.

- A student dissolves 1.26 g of a drain cleaner in water and makes up the solution to 100.0 cm³.
- The student measures the pH of this solution as 13.48.

Determine the percentage, by mass, of NaOH in the drain cleaner. *Strong base*

Give your answer to three significant figures.

$$[H^+] = 10^{-pH} = 10^{-13.48} = 3.31 \times 10^{-14} \text{ mol dm}^{-3}$$

$$[OH^-] = \frac{K_w}{[H^+]} = \frac{1 \times 10^{-14}}{3.31 \times 10^{-14}} = 0.302 \text{ mol dm}^{-3}$$

cm³ → dm³

$$0.302 \times 100 \times 10^{-3} = 0.0302 \text{ mol}$$

n *mass*
v x c *RFM x mol*

$$0.302 \times 40 = 1.21 \text{ g}$$


RFM of NaOH

$$\frac{1.21}{1.26} \times 100 = 95.9\% \text{ (3sf.)}$$

percentage = 95.9 % [4]

(c) Sodium carbonate, Na₂CO₃, is a base used in washing soda.


Na₂CO₃ contains the carbonate ion, CO₃²⁻, shown below.



Structure to base dot and cross diagram from

Draw the 'dot-and-cross' diagram for the carbonate ion.

Show outer electrons only and use different symbols for electrons from C and O, and any 'extra' electrons.



[2]

20 This question is about the halogen group of elements and some of their compounds.

(a) The halogens show trends in their properties down the group.

The boiling points of three halogens are shown below.

Halogen	Boiling point/°C
Chlorine	-35
Bromine	59
Iodine	184

Explain why the halogens show this trend in boiling points.

Down the group: London forces increase because the number of electrons increases so more energy needed to break London forces (hence why boiling points increase down the group).

[3]


(b) Hydrogen iodide, HI, is decomposed by heat into its elements:



The decomposition is much faster in the presence of a platinum catalyst.

Complete the enthalpy profile diagram for this reaction using formulae for the reactants and products.

- Use E_a to label the activation energy **without** a catalyst.
- Use E_c to label the activation energy **with** a catalyst.
- Use ΔH to label the **enthalpy change** of reaction.



[3]

23

- (c) Compound A is an oxide of chlorine that is a liquid at room temperature and pressure and has a boiling point of 83°C . \rightarrow consists only of Cl and O

When 0.4485g of A is heated to 100°C at $1.00 \times 10^5\text{Pa}$, 76.0cm^3 of gas is produced.

Determine the molecular formula of compound A.

$$PV = nRT$$

$$n = \frac{PV}{RT}$$

Show all your working.

$$P = 1 \times 10^5 \text{ Pa}$$

$$V = 76 \times 10^{-6} \text{ cm}^3 \rightarrow \text{m}^3$$

$$n = ?$$

$$R = 8.314$$

$$T = 100 + 273 = 373\text{K}$$

$$n = \frac{1 \times 10^5 \times 76 \times 10^{-6}}{8.314 \times 373} = 2.45 \times 10^{-3} \text{ mol}$$

$$\frac{\text{mass}}{\text{mol}} = \text{RFM}$$

$$\frac{0.4485}{2.45 \times 10^{-3}} = 183$$

$$\underbrace{(35.5 \times 2)}_{\text{RFM of Cl}} + \underbrace{(16 \times 7)}_{\text{RFM of O}} = 183$$

molecular formula of A = Cl_2O_7 [4]


- (d) Compound **B** is an iodate(V) salt of a Group 1 metal. The iodate(V) ion has the formula IO_3^- .

A student carries out a titration to find the formula of compound **B**.


Step 1: The student dissolves 1.55 g of **B** in water and makes up the solution to 250.0 cm³ in a volumetric flask.

Step 2: The student pipettes 25.00 cm³ of the solution of **B** into a conical flask, followed by 10 cm³ of dilute sulfuric acid and an excess of KI(aq).

The iodate(V) ions are reduced to iodine, as shown below.



Step 3: The resulting mixture is titrated with 0.150 mol dm⁻³ Na₂S₂O₃(aq).



The student repeats **step 2** and **step 3** until concordant titres are obtained.

Titration readings

Titration	Trial	1	2	3
Final burette reading / cm ³	24.00	47.40	23.75	47.05
Initial burette reading / cm ³	0.00	24.00	0.00	23.20
Titre / cm ³	24.00	23.40	23.75	23.85

all titre readings written to 2 dp.
Table 20.1

- (i) Complete **Table 20.1** and calculate the mean titre that the student should use for analysing the results.

$$\frac{23.75 + 23.85}{2} = 23.80$$

mean titre = 23.80 cm³ [2]

- (ii) The uncertainty in each burette reading is ±0.05 cm³.

Calculate the percentage uncertainty in the titre obtained from **titration 1**.

Give your answer to **two decimal places**.

number of times equipment was used to find titre

$$\frac{0.05 \times 2}{23.40} \times 100 = 0.43\%$$

titre from titration 1

percentage uncertainty = 0.43 % [1]

25

- (iii) Describe and explain how the student should determine the end point of this titration accurately.

Add Starch

colour change: blue to colourless

[2]

- (iv) Determine the relative formula mass and formula of the Group 1 iodate(V), B.

Show your working.

$$\text{mol of } \text{S}_2\text{O}_3^{2-} : 0.15 \times 23.80 \times 10^{-3} = 3.57 \times 10^{-3} \text{ mol}$$

$$\text{mol of } \text{IO}_3^- : \frac{3.57 \times 10^{-3}}{6} = 5.95 \times 10^{-4} \text{ mol}$$

$$5.95 \times 10^{-4} \times 10 = 5.95 \times 10^{-3} \text{ mol} \quad \begin{matrix} (25 \text{ cm}^3) \\ (250 \text{ cm}^3) \end{matrix}$$

$$\frac{\text{mass}}{\text{mol}} = \text{RFM}$$

$$\frac{1.55}{5.95 \times 10^{-3}} = 260.5$$

$$260.5 - \underbrace{\text{RFM of } \text{IO}_3^-}$$

$$(126.9 + (16 \times 3)) = 174.9$$

$$260.5 - 174.9 = 85.6$$

closest group 1 metal with RFM near to 85.6 is Rb

so formula of B: RbIO_3

relative formula mass of B = 260.5

formula of B = RbIO_3 [5]

21 This question is about some reactions of d block elements and their ions.

Table 21.1 shows standard electrode potentials which will be needed within this question.

<i>stage 2</i>	$Zn^{2+}(aq) + 2e^{-}$	\rightleftharpoons	$Zn(s)$	$E^{\ominus} = -0.76V$	<i>Cell with more negative E^{\ominus} undergoes oxidation</i>
	$Cr^{3+}(aq) + e^{-}$	\rightleftharpoons	$Cr^{2+}(aq)$	$E^{\ominus} = -0.42V$	
	$Ni^{2+}(aq) + 2e^{-}$	\rightleftharpoons	$Ni(s)$	$E^{\ominus} = -0.25V$	
<i>stage 1</i>	$I_2(aq) + 2e^{-}$	\rightleftharpoons	$2I^{-}(aq)$	$E^{\ominus} = +0.54V$	
	$Fe^{3+}(aq) + e^{-}$	\rightleftharpoons	$Fe^{2+}(aq)$	$E^{\ominus} = +0.77V$	
	$Cr_2O_7^{2-}(aq) + 14H^{+}(aq) + 6e^{-}$	\rightleftharpoons	$2Cr^{3+}(aq) + 7H_2O(l)$	$E^{\ominus} = +1.33V$	
	$H_2O_2(aq) + 2H^{+}(aq) + 2e^{-}$	\rightleftharpoons	$2H_2O(l)$	$E^{\ominus} = +1.78V$	<i>↻</i>


Table 21.1

(a) Complete the electron configuration of

a Ni atom: $1s^2 \dots 2s^2 2p^6 3s^2 3p^6$ **$4s^2$** $3d^8$ *filled first and lost first*

a Ni^{2+} ion: $1s^2 \dots 2s^2 2p^6 3s^2 3p^6 3d^8$ [2]


(b) A standard cell is set up in the laboratory with the cell reaction shown below.



needs a pt electrode
can be an electrode

(i) Draw a labelled diagram to show how this cell could be set up to measure its standard cell potential.

Include details of apparatus, solutions and the standard conditions required.



Standard conditions 1 mol dm^{-3} , 298 K

[4]

27

- (ii) Predict the standard cell potential of this cell.

$$0.54 - (-0.25) = 0.79 \text{ V}$$

most positive - most negative

standard cell potential = 0.79 V [1]

- (c) Use the information in Table 21.1 to help you answer both parts of this question.


- (i) Write the overall equation for the oxidation of
- Fe^{2+}
- by acidified
- H_2O_2
- .



- (ii) Zinc reacts with acidified
- $\text{Cr}_2\text{O}_7^{2-}$
- ions to form
- Cr^{2+}
- ions in two stages.

Explain why this happens in terms of electrode potentials and equilibria.

Include overall equations for the reactions which occur.



E^\ominus of Zn is more negative than E^\ominus of $\text{Cr}_2\text{O}_7^{2-}$ so Zn system shifts left. [4]

(d)* Three different reactions of copper compounds are described below.

Reaction 1: Aqueous copper(II) sulfate reacts with excess aqueous ammonia in a ligand substitution reaction. A deep-blue solution is formed, containing an octahedral complex ion, C, which is a trans isomer.

Reaction 2: Copper(I) oxide reacts with hot dilute sulfuric acid in a disproportionation reaction. A blue solution, D, and a brown solid, E are formed.

Reaction 3: Copper(II) oxide reacts with warm dilute nitric acid in a neutralisation reaction, to form a blue solution. Unreacted copper(II) oxide is filtered off, and the solution is left overnight in an evaporating basin.




A hydrated salt, F, crystallises, with the percentage composition by mass:

Cu, 26.29%; H, 2.48%; N, 11.59%; O, 59.63%.


Identify C–F by formulae or structures, as appropriate.

Include equations, any changes in oxidation number, and working.


[6]



Cu oxidation states: $+1 \rightarrow +2 + 0$




Additional answer space if required.



29

$$\begin{array}{cccc} = 0.41 & = 2.49 & = 0.82 & = 3.72 \\ \text{divide each of these numbers by the smallest (0.41)} & & & \\ \frac{0.41}{0.41} = 1 & \frac{2.49}{0.41} = 6 & \frac{0.82}{0.41} = 2 & \frac{3.72}{0.41} = 9 \end{array}$$



END OF QUESTION PAPER

ADDITIONAL ANSWER SPACE

If additional space is required, you should use the following lined page(s). The question number(s) must be clearly shown in the margin(s).

A large area of lined paper for writing answers. It consists of a vertical solid line on the left side, creating a margin. To the right of this line, there are 25 horizontal dotted lines spaced evenly down the page, providing a grid for writing.