

Surname	Centre Number	Candidate Number
First name(s)		2



GCE AS

B410U20-1



S23-B410U20-1



TUESDAY, 23 MAY 2023 – MORNING

## CHEMISTRY – AS component 2

## Energy, Rate and Chemistry of Carbon Compounds

1 hour 30 minutes

## ADDITIONAL MATERIALS

In addition to this examination paper, you will need a:

- calculator;
- **Data Booklet** supplied by WJEC.

## INSTRUCTIONS TO CANDIDATES

Use black ink or black ball-point pen. Do not use gel pen or correction fluid.

You may use a pencil for graphs and diagrams only.

Write your name, centre number and candidate number in the spaces at the top of this page.

**Section A** Answer **all** questions.

**Section B** Answer **all** questions.

Write your answers in the spaces provided in this booklet. If you run out of space, use the additional page(s) at the back of the booklet, taking care to number the question(s) correctly.

## INFORMATION FOR CANDIDATES

The number of marks is given in brackets at the end of each question or part-question.

The maximum mark for this paper is 80.

Your answers must be relevant and must make full use of the information given to be awarded full marks for a question.

The assessment of the quality of extended response (QER) will take place in **Q.9(a)**.

Section A

Section B

For Examiner's use only		
Question	Maximum Mark	Mark Awarded
1. to 6.	10	
7.	19	
8.	12	
9.	9	
10.	16	
11.	14	
Total	80	

B410U201  
01

JUN23B410U20101

**BLANK PAGE**


**PLEASE DO NOT WRITE  
ON THIS PAGE**



**SECTION A**Answer **all** questions.


1. Name the compound whose formula is shown below.

[1]



.....

2. Geraniol is an alcohol which occurs in the essential oils of many aromatic plants. Its skeletal formula is shown below.

Give the **molecular** formula of geraniol.

[1]

.....

3. State the meaning of the term homolytic bond fission.

[1]

.....

.....



4. Hydrofluorocarbons, HFCs, are synthetic compounds which have replaced chlorofluorocarbons, CFCs, as refrigerants in cooling systems because they do not deplete the ozone layer.

Explain why HFCs do not deplete the ozone layer.

[2]

.....

.....

.....


.....

5. Draw the structure of (*E*)-2-iodobut-2-ene.

[2]



6. The diagram below shows the distribution of energies of gas molecules in a reaction at 20 °C.



- (a) On the same axes, draw a curve to show the distribution of energies of the same gas molecules with the reaction at 10 °C. [1]
- (b) Use the diagram to explain the difference in reaction rate between the two reactions. [2]

.....

.....

.....

.....



**SECTION B**Answer **all** questions.

7. (a) Propan-2-ol is the simplest example of a secondary alcohol. It is a common ingredient in many antiseptics.

A student carried out an experiment to determine the enthalpy change of combustion of propan-2-ol.

A spirit burner containing propan-2-ol was placed under a beaker containing  $100\text{ cm}^3$  of water. The burner was ignited and the alcohol allowed to burn until the temperature of the water rose by  $25.4^\circ\text{C}$ . After the burner had cooled it was found that  $0.362\text{ g}$  of propan-2-ol had been burned.

- (i) State what is meant by a secondary alcohol. [1]

.....

.....

- (ii) Write an equation for the complete combustion of propan-2-ol. [1]

.....

- (iii) Calculate a value for the enthalpy change of combustion of propan-2-ol in  $\text{kJ mol}^{-1}$ . [4]

$\Delta_c H = \dots\dots\dots \text{kJ mol}^{-1}$



- (iv) The teacher said that the actual value for the enthalpy change of combustion of propan-2-ol is  $-2006 \text{ kJ mol}^{-1}$ .

Calculate the percentage error in the student's experimental value. [1]


Percentage error = ..... %

- (v) Suggest **one** reason, other than heat loss, why the value obtained for the enthalpy change of combustion is smaller than the theoretical value. [1]

.....



(b) The equation for the combustion of propene is as shown.



- (i) Use this and the average bond enthalpy values given in the table to calculate the average bond enthalpy for the  $\text{O}=\text{O}$  bond. [4]

Bond	Average bond enthalpy / $\text{kJ mol}^{-1}$
$\text{C}=\text{C}$	612
$\text{C}-\text{C}$	348
$\text{C}-\text{H}$	412
$\text{C}=\text{O}$	805
$\text{O}-\text{H}$	463

Average bond enthalpy = .....  $\text{kJ mol}^{-1}$

- (ii) Suggest why the student could not use the method in (a) to determine the enthalpy change of combustion of propene. [1]

.....





- (c) Propan-2-ol can react with ethanoic acid in the presence of concentrated sulfuric acid to form an ester and water. This reaction is reversible so an equilibrium mixture is produced and the ester is removed by distillation.

- (i) Write the equation for the reaction between propan-2-ol and ethanoic acid.

Clearly show the structure of the ester formed.

[2]

- (ii) Explain fully why the ester can be separated from the equilibrium mixture by distillation.

[2]

.....

.....

.....

.....

- (iii) I. Explain why the yield of ester in the equilibrium mixture is increased by removing the ester.

[1]

.....

.....

- II. Concentrated sulfuric acid is a dehydrating agent.

Suggest why the yield of ester in the equilibrium mixture is increased by adding concentrated sulfuric acid.

[1]

.....

.....



**BLANK PAGE**

**PLEASE DO NOT WRITE  
ON THIS PAGE**



8. (a) Alkanes and alkenes are hydrocarbons. Alkanes can be obtained from fossil fuels such as petroleum.

Alkenes can be obtained by cracking long-chain hydrocarbons. They are more reactive than alkanes.

- (i) The use of fossil fuels to meet our energy needs has many disadvantages, but there are some benefits.

Give one advantage of the use of fossil fuels. [1]

.....

.....

- (ii) Name a **solid** pollutant that may form if a fossil fuel is burned incompletely in air. [1]

.....

- (iii) Explain why alkenes are more reactive than alkanes. [2]

.....

.....

.....

.....

.....

.....

- (iv) One molecule of decane,  $C_{10}H_{22}$ , can be cracked to give one molecule of pentane and two other products.

Write an equation for this reaction. [1]

.....




- (b) The information given below relates to liquid **X**.

Quantitative analysis shows that it contains 29.2% carbon and 5.8% hydrogen by mass. The remainder is bromine.

The mass spectrum shows two molecular ion signals at  $m/z$  122 and  $m/z$  124 in the ratio of 1:1.

Its  $^{13}\text{C}$  NMR spectrum is shown below.



- (i) Use **all** the information to identify liquid **X**.

[5]

.....

.....

.....

.....


.....



(ii) Sketch the low resolution  $^1\text{H}$ NMR spectrum of liquid **X**.

Identify which protons are responsible for each peak giving the approximate chemical shift (ppm) and the relative area of each peak.

[2]



9. (a) A student is provided with unlabelled samples of the three liquids listed below.

1-chloropropane

hex-1-ene

propanoic acid

Describe chemical tests, apart from the use of an indicator, that the student should carry out to positively identify each of the three liquids.

You should include balanced equations for any reactions that occur.

[6 QER]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



- (b) 1-Chloropropane,  $\text{C}_3\text{H}_7\text{Cl}$ , can be converted into propylamine,  $\text{C}_3\text{H}_7\text{NH}_2$ .

If the percentage yield of the reaction is 34.5%, calculate the mass of propylamine made from 8.93 g of 1-chloropropane.

Give your answer to an **appropriate** number of significant figures.

[3]

$$M_r(\text{C}_3\text{H}_7\text{Cl}) = 78.6$$

$$M_r(\text{C}_3\text{H}_7\text{NH}_2) = 59.1$$

Mass of propylamine = ..... g

9



10. A student carried out an experiment to study the rate of the reaction between barium carbonate and hydrochloric acid.



$M_r$  197

He started with 1.50 g of barium carbonate and 30.0 cm<sup>3</sup> of 0.400 mol dm<sup>-3</sup> hydrochloric acid. The experiment was carried out at 25 °C and 1 atm.

He determined the rate by following the loss in mass over 5 minutes.


- (a) State which reactant is in excess. Use the data given to justify your answer. [2]

.....  
.....





The student plotted his results as follows.



- (b) (i) Use the graph to calculate the **mean** rate of the reaction, in  $\text{g s}^{-1}$ . [1]

Mean rate = .....  $\text{g s}^{-1}$

- (ii) Use the graph to calculate the **initial** rate of the reaction, in  $\text{g s}^{-1}$ . [2]

Initial rate = .....  $\text{g s}^{-1}$



- (c) (i) Use the graph to describe how the rate of the reaction changes over the 5 minutes. [1]

.....

.....

- (ii) Use collision theory to explain why the rate of this reaction changes as the reaction proceeds. [2]

.....

.....

.....

.....

- (d) Outline the method the student used to carry out this experiment.

A diagram may be used in support of your answer. [3]

.....

.....

.....

.....

.....

.....



- (e) Another method to follow this reaction is by measuring the volume of carbon dioxide produced over time.

Name the apparatus you would use to collect the carbon dioxide. [1]

.....

- (f) Use the graph to calculate the volume (in  $\text{cm}^3$ ) of carbon dioxide formed during this reaction at  $25^\circ\text{C}$ .

You **must** show your working. [2]

Volume = .....  $\text{cm}^3$

- (g) Another student said that if you used the same mass of calcium carbonate instead of barium carbonate, the volume of carbon dioxide formed would be less because the relative formula mass of calcium carbonate is less.

Is she correct? Justify your answer. [2]

.....

.....

.....

.....




**BLANK PAGE**

**PLEASE DO NOT WRITE  
ON THIS PAGE**




11. Five non-cyclic structural isomers have the molecular formula  $C_5H_{10}$ .


Three of these are shown below.



pent-1-ene



pent-2-ene



3-methylbut-1-ene

(a) Draw the structures of the other **two** isomers.


Do **not** include both the *E* and the *Z* isomer of the same structure.

[2]



(b) Compound **A** is one of the five isomers on the previous page.

Study the reaction scheme and the information below.



Compound **D** does **not** react with aqueous sodium carbonate.

Compound **E** does **not** show *E-Z* isomerism.

- (i) Use the information to name the homologous series to which compounds **B–E** belong. Give your reasoning. [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



- (ii) State what you would observe when compound **C** is oxidised to compound **D**. [1]

.....

- (iii) Compound **B** can also undergo an elimination reaction.

State the reagent(s) and conditions needed for this reaction. [1]

.....

- (iv) Compound **A** can **only** be 3-methylbut-1-ene.

It cannot be pent-2-ene because its reaction with HBr would give two products in significant quantities.

- I. Explain why compound **A** cannot be pent-1-ene. [2]

.....

.....

.....

.....

.....

.....

- II. Explain why compound **A** cannot be either of the two isomers drawn in part (a). [2]

.....

.....

.....

.....

.....

.....

END OF PAPER









**BLANK PAGE**

**PLEASE DO NOT WRITE  
ON THIS PAGE**



**BLANK PAGE**

**PLEASE DO NOT WRITE  
ON THIS PAGE**



**BLANK PAGE**

**PLEASE DO NOT WRITE  
ON THIS PAGE**





GCE AS

B410U20-1A



S23-B410U20-1A



TUESDAY, 23 MAY 2023 – MORNING

## CHEMISTRY – AS component 2

### Data Booklet

Avogadro constant  
 molar gas constant  
 molar gas volume at 273 K and 1 atm  
 molar gas volume at 298 K and 1 atm  
 Planck constant  
 speed of light  
 density of water  
 specific heat capacity of water  
 ionic product of water at 298 K  
 fundamental electronic charge

$$\begin{aligned}
 N_A &= 6.02 \times 10^{23} \text{ mol}^{-1} \\
 R &= 8.31 \text{ J mol}^{-1} \text{ K}^{-1} \\
 V_m &= 22.4 \text{ dm}^3 \text{ mol}^{-1} \\
 V_m &= 24.5 \text{ dm}^3 \text{ mol}^{-1} \\
 h &= 6.63 \times 10^{-34} \text{ J s} \\
 c &= 3.00 \times 10^8 \text{ m s}^{-1} \\
 d &= 1.00 \text{ g cm}^{-3} \\
 c &= 4.18 \text{ J g}^{-1} \text{ K}^{-1} \\
 K_w &= 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6} \\
 e &= 1.60 \times 10^{-19} \text{ C}
 \end{aligned}$$

temperature (K) = temperature (°C) + 273

$1 \text{ dm}^3 = 1000 \text{ cm}^3$   
 $1 \text{ m}^3 = 1000 \text{ dm}^3$   
 1 tonne = 1000 kg  
 $1 \text{ atm} = 1.01 \times 10^5 \text{ Pa}$

Multiple	Prefix	Symbol
$10^{-9}$	nano	n
$10^{-6}$	micro	$\mu$
$10^{-3}$	milli	m




Multiple	Prefix	Symbol
$10^3$	kilo	k
$10^6$	mega	M
$10^9$	giga	G

## Infrared absorption values





Bond	Wavenumber / $\text{cm}^{-1}$
C — Br	500 to 600
C — Cl	650 to 800
C — O	1000 to 1300
C = C	1620 to 1670
C = O	1650 to 1750
C $\equiv$ N	2100 to 2250
C — H	2800 to 3100
O — H (carboxylic acid)	2500 to 3200 (very broad)
O — H (alcohol / phenol)	3200 to 3550 (broad)
N — H	3300 to 3500

 $^{13}\text{C}$  NMR chemical shifts relative to TMS = 0

Type of carbon	Chemical shift, $\delta$ (ppm)
----------------	--------------------------------



**$^1\text{H}$  NMR chemical shifts relative to TMS = 0**

Type of proton	Chemical shift, $\delta$ (ppm)
$-\text{CH}_3$	0.1 to 2.0
$\text{R}-\text{CH}_3$	0.9
$\text{R}-\text{CH}_2-\text{R}$	1.3
$\text{CH}_3-\text{C}\equiv\text{N}$	2.0
$\text{CH}_3-\text{C}(=\text{O})$	2.0 to 2.5
$-\text{CH}_2-\text{C}(=\text{O})$	2.0 to 3.0
 - $\text{CH}_3$	2.2 to 2.3
$\text{HC}-\text{Cl}$ or $\text{HC}-\text{Br}$	3.1 to 4.3
$\text{HC}-\text{O}$	3.3 to 4.3
$\text{R}-\text{OH}$	4.5 *
$-\text{C}=\text{CH}$	4.5 to 6.3
$-\text{C}=\text{CH}-\text{CO}$	5.8 to 6.5
 - $\text{CH}=\text{C}$	6.5 to 7.5
 - $\text{H}$	6.5 to 8.0
 - $\text{OH}$	7.0 *
$\text{R}-\text{C}(=\text{O})\text{H}$	9.8 *
$\text{R}-\text{C}(=\text{O})\text{OH}$	11.0 *

\*variable figure dependent on concentration and solvent

# THE PERIODIC TABLE

Group

1 2 3 4 5 6 7 0

Period

1 2


1	1.01 H Hydrogen 1						4.00 He Helium 2
2	6.94 Li Lithium 3	9.01 Be Beryllium 4					
3	23.0 Na Sodium 11	24.3 Mg Magnesium 12					
4	39.1 K Potassium 19	40.1 Ca Calcium 20					
5	85.5 Rb Rubidium 37	87.6 Sr Strontium 38					
6	133 Cs Caesium 55	137 Ba Barium 56					
7	(223) Fr Francium 87	(226) Ra Radium 88					

	10.8 B Boron 5	12.0 C Carbon 6	14.0 N Nitrogen 7	16.0 O Oxygen 8	19.0 F Fluorine 9	20.2 Ne Neon 10
	27.0 Al Aluminium 13	28.1 Si Silicon 14	31.0 P Phosphorus 15	32.1 S Sulfur 16	35.5 Cl Chlorine 17	40.0 Ar Argon 18
	69.7 Ga Gallium 31	72.6 Ge Germanium 32	74.9 As Arsenic 33	79.0 Se Selenium 34	79.9 Br Bromine 35	83.8 Kr Krypton 36
	115 In Indium 49	119 Sn Tin 50	122 Sb Antimony 51	128 Te Tellurium 52	127 I Iodine 53	131 Xe Xenon 54
	204 Tl Thallium 81	207 Pb Lead 82	209 Bi Bismuth 83	(210) Po Polonium 84	(210) At Astatine 85	(222) Rn Radon 86

	65.4 Zn Zinc 30	63.5 Cu Copper 29	58.7 Ni Nickel 28	58.9 Co Cobalt 27	55.8 Fe Iron 26	54.9 Mn Manganese 25	52.0 Cr Chromium 24	50.9 V Vanadium 23	47.9 Ti Titanium 22	45.0 Sc Scandium 21
	112 Cd Cadmium 48	108 Ag Silver 47	106 Pd Palladium 46	103 Rh Rhodium 45	101 Ru Ruthenium 44	98.9 Tc Technetium 43	95.9 Mo Molybdenum 42	92.9 Nb Niobium 41	91.2 Zr Zirconium 40	88.9 Y Yttrium 39
	201 Hg Mercury 80	197 Au Gold 79	195 Pt Platinum 78	192 Ir Iridium 77	190 Os Osmium 76	186 Re Rhenium 75	184 W Tungsten 74	181 Ta Tantalum 73	179 Hf Hafnium 72	139 La Lanthanum 57
	(227) Ac Actinium 89	(227) Ac Actinium 89	(227) Ac Actinium 89	(227) Ac Actinium 89	(227) Ac Actinium 89	(227) Ac Actinium 89	(227) Ac Actinium 89	(227) Ac Actinium 89	(227) Ac Actinium 89	(227) Ac Actinium 89



f block

► Lanthanoid elements

140 Ce Cerium 58	141 Pr Praseodymium 59	144 Nd Neodymium 60	(147) Pm Promethium 61	150 Sm Samarium 62	(153) Eu Europium 63	157 Gd Gadolinium 64	159 Tb Terbium 65	163 Dy Dysprosium 66	165 Ho Holmium 67	167 Er Erbium 68	169 Tm Thulium 69	173 Yb Ytterbium 70	175 Lu Lutetium 71
---------------------------	---------------------------------	------------------------------	---------------------------------	-----------------------------	-------------------------------	-------------------------------	----------------------------	-------------------------------	----------------------------	---------------------------	----------------------------	------------------------------	-----------------------------

►► Actinoid elements

232 Th Thorium 90	(231) Pa Protactinium 91	238 U Uranium 92	(237) Np Neptunium 93	(242) Pu Plutonium 94	(243) Am Americium 95	(247) Cm Curium 96	(245) Bk Berkelium 97	(251) Cf Californium 98	(254) Es Einsteinium 99	(253) Fm Fermium 100	(256) Md Mendelevium 101	(254) No Nobelium 102	(257) Lr Lawrencium 103
----------------------------	-----------------------------------	---------------------------	--------------------------------	--------------------------------	--------------------------------	-----------------------------	--------------------------------	----------------------------------	----------------------------------	-------------------------------	-----------------------------------	--------------------------------	----------------------------------