



---

# **GCE A LEVEL MARKING SCHEME**

---

**AUTUMN 2020**

**A LEVEL  
CHEMISTRY – COMPONENT 3  
A410U30-1**

## **INTRODUCTION**

This marking scheme was used by WJEC for the 2020 examination. It was finalised after detailed discussion at examiners' conferences by all the examiners involved in the assessment. The conference was held shortly after the paper was taken so that reference could be made to the full range of candidates' responses, with photocopied scripts forming the basis of discussion. The aim of the conference was to ensure that the marking scheme was interpreted and applied in the same way by all examiners.

It is hoped that this information will be of assistance to centres but it is recognised at the same time that, without the benefit of participation in the examiners' conference, teachers may have different views on certain matters of detail or interpretation.

WJEC regrets that it cannot enter into any discussion or correspondence about this marking scheme.

**GCE A LEVEL CHEMISTRY COMPONENT 3****CHEMISTRY IN PRACTICE****AUTUMN 2020 MARK SCHEME****GENERAL INSTRUCTIONS**Recording of marks

Examiners must mark in red ink.

One tick must equate to one mark, apart from extended response questions where a level of response mark scheme is applied.

Question totals should be written in the box at the end of the question.

Question totals should be entered onto the grid on the front cover and these should be added to give the script total for each candidate.

Extended response questions

A level of response mark scheme is applied. The complete response should be read in order to establish the most appropriate band. Award the higher mark if there is a good match with content and communication criteria. Award the lower mark if either content or communication barely meets the criteria.

Marking rules

All work should be seen to have been marked.

Marking schemes will indicate when explicit working is deemed to be a necessary part of a correct answer.


Crossed out responses not replaced should be marked.

### Marking abbreviations

The following may be used in marking schemes or in the marking of scripts to indicate reasons for the marks awarded.

cao = correct answer only  
ecf = error carried forward  
bod = benefit of doubt

Credit should be awarded for correct and relevant alternative responses which are not recorded in the mark scheme.

Question			Marking details	Marks available					
				AO1	AO2	AO3	Total	Maths	Prac
1	(a)		<b>D</b> methylpropan-2-ol (1) <b>H</b> pentanenitrile (1)	2			2		
	(b)	(i)	award (1) for correctly drawn structures of both optical isomers of $\text{CH}_3\text{CH}(\text{CN})\text{CH}_2\text{CH}_3$ 		1		1		
		(ii)	they rotate the plane of plane polarised light in opposite directions	1			1		
		(iii)	award (1) for reagent and (1) for product e.g. (reduction with) $\text{LiAlH}_4$ (1) to form $\text{CH}_3\text{CH}(\text{CH}_2\text{NH}_2)\text{CH}_2\text{CH}_3$ (1) (hydrolysis with) $\text{HCl}$ (1) to form $\text{CH}_3\text{CH}(\text{COOH})\text{CH}_2\text{CH}_3$ (1) (hydrolysis with) $\text{NaOH}$ (1) to form $\text{CH}_3\text{CH}(\text{COONa})\text{CH}_2\text{CH}_3$ (1)		2		2		2

Question		Marking details				Marks available					
						AO1	AO2	AO3	Total	Maths	Prac
(c)		Compounds	Reagent(s) and condition(s)	Observation(s)	Organic compound(s) formed	4	4	8			
		CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> OH and CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> Br	acidified dichromate(VI) / heat (reflux)	orange to green solution	CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CHO or CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> COOH						
		CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CHO and (CH <sub>3</sub> ) <sub>3</sub> COH	Tollens' reagent (alkaline solution of ammoniacal silver nitrate)  warm gently in hot water bath	silver mirror	CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> COOH						
		CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> COOH and CH <sub>3</sub> CH <sub>2</sub> CH(OH)CH <sub>3</sub>	I <sub>2</sub> / NaOH(aq)  or KI / NaClO(aq)	yellow solid	CHI <sub>3</sub> and CH <sub>3</sub> CH <sub>2</sub> COONa						
		CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> NH <sub>2</sub> and CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CN	nitric(III) acid (HNO <sub>2</sub> )  room temperature	bubbles of gas	CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> OH						
		award (1) for each reagent with conditions, for each observation and for each compound formed									

Question				Marking details	Marks available					
					AO1	AO2	AO3	Total	Maths	Prac
	(d)	(i)		reflux in the presence of concentrated sulfuric acid	1			1		1
		(ii)		$\text{CH}_3\text{CH}_2\text{CH}_2\text{COOH} + \text{CH}_3\text{CH}_2\text{CH}(\text{OH})\text{CH}_3 \rightleftharpoons \text{CH}_3\text{CH}_2\text{CH}_2\text{COOCH}(\text{CH}_3)\text{CH}_2\text{CH}_3 + \text{H}_2\text{O}$			1	1		
		(iii)		(addition of sodium carbonate to neutralise acid and) distillation		1		1		1
				<b>Question 1 total</b>	<b>8</b>	<b>8</b>	<b>1</b>	<b>17</b>	<b>0</b>	<b>8</b>

Question			Marking details				Marks available																									
							AO1	AO2	AO3	Total	Maths	Prac																				
2	(a)		add further sodium carbonate solution to the filtrate and no precipitate will form						1	1		1																				
	(b)		$2\text{H}^+ + \text{CO}_3^{2-} \rightarrow \text{H}_2\text{O} + \text{CO}_2$				1			1																						
	(c)	(i)	<table border="1" style="width: 100%; border-collapse: collapse; text-align: center;"> <thead> <tr> <th></th> <th>Titration 1</th> <th>Titration 2</th> <th>Titration 3</th> <th>Titration 4</th> </tr> </thead> <tbody> <tr> <td>Initial burette reading / cm<sup>3</sup></td> <td>0.50</td> <td>18.45</td> <td>2.10</td> <td>19.70</td> </tr> <tr> <td>Final burette reading / cm<sup>3</sup></td> <td>18.45</td> <td>35.95</td> <td>19.70</td> <td>37.25</td> </tr> <tr> <td>Titre / cm<sup>3</sup></td> <td>17.95</td> <td>17.50</td> <td>17.60</td> <td>17.55</td> </tr> </tbody> </table> <p>all titres calculated and given to <b>2dp</b> (1)</p> <p>mean titre 17.55 cm<sup>3</sup> (1)</p>					Titration 1	Titration 2	Titration 3	Titration 4	Initial burette reading / cm <sup>3</sup>	0.50	18.45	2.10	19.70	Final burette reading / cm <sup>3</sup>	18.45	35.95	19.70	37.25	Titre / cm <sup>3</sup>	17.95	17.50	17.60	17.55			2	2		2
	Titration 1	Titration 2	Titration 3	Titration 4																												
Initial burette reading / cm <sup>3</sup>	0.50	18.45	2.10	19.70																												
Final burette reading / cm <sup>3</sup>	18.45	35.95	19.70	37.25																												
Titre / cm <sup>3</sup>	17.95	17.50	17.60	17.55																												
		(ii)	titration 2 <u>smallest volume</u> (of HCl) therefore <u>largest percentage error</u>						1	1		1																				



Question		Marking details		Marks available						
				AO1	AO2	AO3	Total	Maths	Prac	
(d)	(i)		Correct order							
		Calculate the number of moles of HCl used in the titration of 25.0 cm <sup>3</sup> of solution Y	2							
		Calculate the number of moles of CO <sub>3</sub> <sup>2-</sup> that reacted with 200 cm <sup>3</sup> of solution Y	4							
		Use the balanced equation to calculate the number of moles of unreacted CO <sub>3</sub> <sup>2-</sup> in 200 cm <sup>3</sup> of solution Y	3							
		Calculate the concentration of the barium chloride solution in g dm <sup>-3</sup>	5							
		Calculate the total number of moles of CO <sub>3</sub> <sup>2-</sup> added to the 50.0 cm <sup>3</sup> of barium chloride solution	1			1	1			1
		award (1) for correct sequence								
		accept alternative order		1						
				4						
				2						
				5						
				3						

Question			Marking details	Marks available					
				AO1	AO2	AO3	Total	Maths	Prac
		(ii)	<p>total mol <math>\text{CO}_3^{2-}</math> added = <math>\frac{50 \times 0.506}{1000} = 0.0253</math> (1)</p> <p>mol HCl in mean titre = <math>\frac{17.55 \times 0.200}{1000} = 0.00351</math> (1)</p> <p>mol unreacted <math>\text{CO}_3^{2-}</math> in 200 <math>\text{cm}^3</math> of solution Y</p> <p><math>\frac{0.00351 \times 8}{2} = 0.0140</math> (1)</p> <p>mol <math>\text{CO}_3^{2-}</math> reacted with 200 <math>\text{cm}^3</math> of solution Y</p> <p><math>0.0253 - 0.0140 = 0.0113</math> (1)</p> <p><math>[\text{BaCl}_2] = \frac{0.0113}{\frac{50}{1000}} = 0.226 \text{ mol dm}^{-3}</math></p> <p><math>0.226 \times 208 = 47.0 \text{ g dm}^{-3}</math> (1)</p>		5		5	4	
		(iii)	<p>mass <math>\text{BaCO}_3 = 0.226 \times \frac{50}{1000} \times 197 = 2.23</math></p>		1		1	1	
			<b>Question 2 total</b>	<b>1</b>	<b>8</b>	<b>3</b>	<b>12</b>	<b>5</b>	<b>5</b>



Question		Marking details		Marks available						
				AO1	AO2	AO3	Total	Maths	Prac	
3	(a)		<p>award (2) for any <b>three</b> of following conclusions leading to identification of <b>X</b> as magnesium award (1) for any <b>two</b> conclusions</p> <ul style="list-style-type: none"> <li>• high melting temperature of oxide and chloride <math>\Rightarrow</math> ionic compounds</li> <li>• basic oxide (not amphoteric) <math>\Rightarrow</math> oxide of metal</li> <li>• insoluble carbonate <math>\Rightarrow</math> not Group 1 metal</li> <li>• hydroxide and carbonate are both white <math>\Rightarrow</math> not transition metal</li> <li>• white precipitate with hydroxide which is insoluble in excess <math>\Rightarrow</math> Mg</li> <li>• no precipitate with sulfate <math>\Rightarrow</math> Mg</li> </ul> <p>award (1) each for <b>ionic</b> equations</p> $\text{Mg}^{2+}(\text{aq}) + 2\text{OH}^{-}(\text{aq}) \rightarrow \text{Mg}(\text{OH})_2(\text{s})$ $\text{Mg}^{2+}(\text{aq}) + \text{CO}_3^{2-}(\text{aq}) \rightarrow \text{MgCO}_3(\text{s})$ <p>accept without state symbols</p>			2				2
				2			4			

Question		Marking details		Marks available					
				AO1	AO2	AO3	Total	Maths	Prac
	(b)		<p>award (2) for any <b>three</b> of following conclusions award (1) for any <b>two</b> conclusions</p> <ul style="list-style-type: none"> <li>no reaction with acids/alkalis <math>\Rightarrow</math> not a metal oxide</li> <li>high melting temperature of oxide <math>\Rightarrow</math> oxide has giant molecular structure</li> <li>low boiling temperature of chloride <math>\Rightarrow</math> chloride has simple covalent structure</li> <li>oxide has giant molecular structure and chloride has simple covalent structure <math>\Rightarrow</math> metalloid / Group 4 element</li> <li>misty fumes and white precipitate with water <math>\Rightarrow</math> Si</li> </ul> <p>chloride of Y</p> <p>333 K and <math>1.01 \times 10^5</math> Pa (1)</p> <p>from <math>pV = nRT</math></p> $n = \frac{1.01 \times 10^5 \times 805.5 \times 10^{-6}}{8.31 \times 333} = 0.0294 \quad (1)$ $M_r = \frac{5.000}{0.0294} = 170.1 \quad \Rightarrow \text{chloride is SiCl}_4 / \text{Y is silicon (1)}$ <p>award (1) for either equation</p> $\text{SiCl}_4 + 2\text{H}_2\text{O} \rightarrow \text{SiO}_2 + 4\text{HCl}$ $\text{SiCl}_4 + 4\text{H}_2\text{O} \rightarrow \text{Si(OH)}_4 + 4\text{HCl}$			2			2
				1	3		6	3	
<b>Question 3 total</b>				<b>3</b>	<b>3</b>	<b>4</b>	<b>10</b>	<b>3</b>	<b>4</b>

Question			Marking details	Marks available					
				AO1	AO2	AO3	Total	Maths	Prac
4	(a)	(i)	<p>award (1) for either of following</p> <p>use gas syringe to measure increase in gas volume over time measure loss in mass over time</p> <p>allow sample at intervals, quench and titrate [credit only in (i) or (ii)]</p>		1		1		1
		(ii)	<p>award (1) for basic method e.g.</p> <p>colorimetry / follow change in colour over time sample at intervals, quench and titrate</p> <p>award (2) for more detailed method e.g.</p> <p>follow change in colour <u>due to change in iodine concentration</u> over time sample at intervals, quench and titrate (iodine) <u>against sodium thiosulfate solution</u></p>		2		2		2

Question				Marking details	Marks available																
					AO1	AO2	AO3	Total	Maths	Prac											
	(b)	(i)	I	award (1) for all four rates calculated  <table border="1" style="margin-left: 20px;"> <thead> <tr> <th>Expt</th> <th>Rate / s<sup>-1</sup></th> </tr> </thead> <tbody> <tr> <td>1</td> <td>2.98 × 10<sup>-3</sup></td> </tr> <tr> <td>2</td> <td>1.19 × 10<sup>-2</sup></td> </tr> <tr> <td>3</td> <td>2.38 × 10<sup>-2</sup></td> </tr> <tr> <td>4</td> <td>5.95 × 10<sup>-3</sup></td> </tr> </tbody> </table>	Expt	Rate / s <sup>-1</sup>	1	2.98 × 10 <sup>-3</sup>	2	1.19 × 10 <sup>-2</sup>	3	2.38 × 10 <sup>-2</sup>	4	5.95 × 10 <sup>-3</sup>							
Expt	Rate / s <sup>-1</sup>																				
1	2.98 × 10 <sup>-3</sup>																				
2	1.19 × 10 <sup>-2</sup>																				
3	2.38 × 10 <sup>-2</sup>																				
4	5.95 × 10 <sup>-3</sup>																				
			II	1 <sup>st</sup> order with respect to BrO <sub>3</sub> <sup>-</sup> (aq) when [BrO <sub>3</sub> <sup>-</sup> ] doubles (expt 2 ⇌ 3), time halves / rate doubles (1)  1 <sup>st</sup> order with respect to Br <sup>-</sup> (aq) when [Br <sup>-</sup> ] halves (expt 2 ⇌ 4), time doubles / rate halves (1)  2 <sup>nd</sup> order with respect to H <sup>+</sup> (aq) when [H <sup>+</sup> ] doubles (expt 1 ⇌ 2), time reduces by a factor of 4 / rate increases by a factor of 4 (1)																	
			III	rate = $k [\text{BrO}_3^-][\text{Br}^-][\text{H}^+]^2$  ecf possible from part II																	

Question				Marking details		Marks available																													
						AO1	AO2	AO3	Total	Maths	Prac																								
			IV	the sum of the powers of the reactant concentrations in the rate equation - this reaction is overall fourth order  ecf possible from parts II and III				1																											
			V	[BrO <sub>3</sub> <sup>-</sup> ], [Br <sup>-</sup> ] and [H <sup>+</sup> ] calculated for any of experiments 1-4 (1)  <table border="1" data-bbox="495 555 1263 807"> <thead> <tr> <th>Expt</th> <th>[BrO<sub>3</sub><sup>-</sup>]</th> <th>[Br<sup>-</sup>]</th> <th>[H<sup>+</sup>]</th> <th>Rate / s<sup>-1</sup></th> </tr> </thead> <tbody> <tr> <td>1</td> <td>0.05</td> <td>0.25</td> <td>0.30</td> <td>2.98 × 10<sup>-3</sup></td> </tr> <tr> <td>2</td> <td>0.05</td> <td>0.25</td> <td>0.60</td> <td>1.19 × 10<sup>-2</sup></td> </tr> <tr> <td>3</td> <td>0.10</td> <td>0.25</td> <td>0.60</td> <td>2.38 × 10<sup>-2</sup></td> </tr> <tr> <td>4</td> <td>0.05</td> <td>0.125</td> <td>0.60</td> <td>5.95 × 10<sup>-3</sup></td> </tr> </tbody> </table> $k = \frac{\text{rate}}{[\text{BrO}_3^-][\text{Br}^-][\text{H}^+]^2} \quad (1)$ $k = 2.64$ award (2) for answer to 3 sig figs award (1) for correct answer not to 3 sig figs  ecf possible from part III	Expt	[BrO <sub>3</sub> <sup>-</sup> ]	[Br <sup>-</sup> ]	[H <sup>+</sup> ]	Rate / s <sup>-1</sup>	1	0.05	0.25	0.30	2.98 × 10 <sup>-3</sup>	2	0.05	0.25	0.60	1.19 × 10 <sup>-2</sup>	3	0.10	0.25	0.60	2.38 × 10 <sup>-2</sup>	4	0.05	0.125	0.60	5.95 × 10 <sup>-3</sup>				4	4	4
Expt	[BrO <sub>3</sub> <sup>-</sup> ]	[Br <sup>-</sup> ]	[H <sup>+</sup> ]	Rate / s <sup>-1</sup>																															
1	0.05	0.25	0.30	2.98 × 10 <sup>-3</sup>																															
2	0.05	0.25	0.60	1.19 × 10 <sup>-2</sup>																															
3	0.10	0.25	0.60	2.38 × 10 <sup>-2</sup>																															
4	0.05	0.125	0.60	5.95 × 10 <sup>-3</sup>																															

Question				Marking details		Marks available						
						AO1	AO2	AO3	Total	Maths	Prac	
			VI	$[\text{BrO}_3^-]$  $[\text{H}^+]$ 	(1)							
					(1)			2	2	2		



Question			Marking details	Marks available					
				AO1	AO2	AO3	Total	Maths	Prac
		(ii)	<p><b>Indicative content</b></p> <ol style="list-style-type: none"> <li>Find time taken for methyl orange indicator to be bleached at a number of different temperatures</li> <li>Initial reactant concentrations remain unchanged across all experiments</li> <li>Calculate the rate at each temperature</li> <li>Calculate the rate constant, <math>k</math>, at each temperature</li> <li>Plot <math>\ln k</math> against <math>\frac{1}{T}</math> where <math>T</math> is temperature in Kelvin</li> <li>Arrhenius equation <math>k = Ae^{\frac{-E_a}{RT}}</math></li> <li>Arrhenius equation rearranged <math>\Rightarrow \ln k = \ln A - \frac{E_a}{RT}</math></li> <li>Gradient of straight line <math>\Rightarrow m = -\frac{E_a}{R}</math></li> <li>Calculate <math>E_a</math></li> </ol>		3	3	6	3	3

Question				Marking details	Marks available						
					AO1	AO2	AO3	Total	Maths	Prac	
				<p><b>5-6 marks</b> Clear outline of how data is processed; good understanding of relationship between graph and Arrhenius equation <i>The candidate constructs a relevant, coherent and logically structured account including all key elements of the indicative content. A sustained and substantiated line of reasoning is evident and scientific conventions and vocabulary is used accurately throughout.</i></p> <p><b>3-4 marks</b> Good knowledge of practical steps; basic understanding of data processing; some idea of link between graph and Arrhenius equation <i>The candidate constructs a coherent account including many of the key elements of the indicative content. Some reasoning is evident in the linking of key points and use of scientific conventions and vocabulary is generally sound.</i></p> <p><b>1-2 marks</b> Some knowledge of practical steps; some idea of data processing <i>The candidate attempts to link at least two relevant points from the indicative material. Coherence is limited by omission and/or inclusion of irrelevant materials. There is some evidence of appropriate use of scientific conventions and vocabulary.</i></p> <p><b>0 marks</b> <i>The candidate does not make any attempt or give an answer worthy of credit.</i></p>							
				<b>Question 4 total</b>	<b>1</b>	<b>8</b>	<b>12</b>	<b>21</b>	<b>12</b>	<b>6</b>	

**COMPONENT 3: CHEMISTRY IN PRACTICE****SUMMARY OF MARKS ALLOCATED TO ASSESSMENT OBJECTIVES**

<b>Question</b>	<b>AO1</b>	<b>AO2</b>	<b>AO3</b>	<b>Total</b>	<b>Maths</b>	<b>Prac</b>
<b>1</b>	<b>8</b>	<b>8</b>	<b>1</b>	<b>17</b>	<b>0</b>	<b>8</b>
<b>2</b>	<b>1</b>	<b>8</b>	<b>3</b>	<b>12</b>	<b>5</b>	<b>5</b>
<b>3</b>	<b>3</b>	<b>3</b>	<b>4</b>	<b>10</b>	<b>3</b>	<b>4</b>
<b>4</b>	<b>1</b>	<b>8</b>	<b>12</b>	<b>21</b>	<b>12</b>	<b>6</b>
<b>Totals</b>	<b>13</b>	<b>27</b>	<b>20</b>	<b>60</b>	<b>20</b>	<b>23</b>