



# **GCE MARKING SCHEME**


**CHEMISTRY  
AS/Advanced**

**JANUARY 2011**

CH1

Section A

1.



[1]

2.

(a)  $M_r = 172.24$

[1]

(b)  $\% = 20.9$


[1]

3.

D

[1]

4.



[3]

5.

(a)

C	H	O	
<u>40</u>	<u>6.7</u>	<u>53.3</u>	
12	1.01	16	
3.33	6.63	3.33	(1)
1	2	1	

Empirical Formula =  $\text{CH}_2\text{O}$

(1) [2]

(b)  $\frac{180}{30.02} = 6$

Molecular Formula =  $\text{C}_6\text{H}_{12}\text{O}_6$

[1]

Total [10]

## Section B

6. (a) (i) Average mass of one atom of the element (1) relative to  $1/12^{\text{th}}$  mass of one atom of carbon-12. (1) [2]
- (ii)  $A_r = \frac{(39 \times 93.26) + (40 \times 0.012) + (41 \times 6.73)}{100}$  (1)  
 $= 39.14$  (1) [2]
- (b) (i) (Gaseous potassium) atoms bombarded by electrons. [1]
- (ii) Deflected through a magnetic field. [1]
- (c) (i)  ${}^{40}_{19}\text{K} \longrightarrow {}^{40}_{20}\text{Ca} + {}^0_{-1}\beta$  (accept  ${}^0_{-1}\text{e}$ )  
 (1 mark for  ${}^{40}_{20}\text{Ca}$ , 1 mark for balanced equation) [2]
- (ii)  $3.75 \times 10^9$  years. [1]
- (d) (i) Energy required to remove one mole of electrons from 1 mole of atoms / an electron from each atom in 1 mole (1) in the gaseous state. (1) [2]  
 (Accept equation)
- (ii) I In K greater shielding of outer electron (1) outweighs larger nuclear charge (1) / Na has greater effective nuclear charge (1) / Na outer electron closer to nucleus (1). (Maximum 2 marks) [2]
- II Shielding effect on outer electron is less (1) / 2nd electron is removed from inner shell / closer to nucleus (1) / after 1st electron is removed effective nuclear charge is greater. (1) [2]  
 (Maximum 2 marks)

Total [15]

7. (a) Bubbles (of gas) / fizzing /  $\text{CaCO}_3$  disappears / apparatus gets warmer [1]
- (b) Gas syringe / burette / graduated tube/measuring cylinder [1]
- (c) (Use scales to) weigh aqueous product / sampling and titration / change in pH at set times [1]
- (d) (i) Moles HCl = 0.020 [1]
- (ii) Moles  $\text{CaCO}_3$  = 0.01 (1)  
Mass = 1.00 g (1) [2]
- (iii) Moles  $\text{CO}_2$  = 0.010 (1)  
Volume = 0.240  $\text{dm}^3$  (1) [2]
- (e) (i) Smooth curve passing through 150  $\text{cm}^3$  ending at 200  $\text{cm}^3$  [1]
- (ii) Curve less steep (1) ending at 100  $\text{cm}^3$  (1) [2]
- (iii) When the acid is less concentrated it has fewer (acid) particles (1) therefore there is less chance of (successful) collisions (between the acid and carbonate) / fewer collisions per unit time. (1) [2]
- (f) Diagram with two reasonable curves. (1 mark) Activation energy labelled (1) The fraction of molecules that have the required activation energy is much greater at a higher temperature. (1) [3]
- QWC Selection of a form and style of writing appropriate to purpose and to complexity of subject matter. [1]

Total [17]

8. (a) (i) Between 1800 and 1900 the global temperature was fairly constant as was the concentration of  $\text{CO}_2$  in the atmosphere. (1)  
 Since 1900 the global temperature has risen steadily as has the concentration of  $\text{CO}_2$  in the atmosphere. (1)
- As concentration of  $\text{CO}_2$  increases, global temperature increases. (1 mark only). [2]
- QWC Legibility of text; accuracy of spelling, punctuation and grammar, clarity of meaning [1]
- (ii) There is an uncertainty in the results / temperature dropped between 1900 and 1910 / between 1940 and 1950 / at some points. [1]
- (iii) Before 1900 the instruments were less accurate (1) and there were fewer records (1)  
 Temperatures are estimates. (1)  
 Any 2 from 3 [2]
- (iv) More burning of fossil fuels / more industries / more transportation / deforestation. (Any two) [2]
- (b) (i) Rate of forward reaction = rate of back reaction. [1]
- (ii) (Molecules can escape from the bottle) so concentration amount of  $\text{CO}_2(\text{g})$  falls / pressure falls (1) and position of equilibrium moves to the left (so concentration of  $\text{CO}_2(\text{aq})$  falls) / rate of molecules entering solution is less than rate leaving solution. (1) [2]
- QWC The information is organised clearly and coherently, using specialist vocabulary where appropriate [1]

Total [12]

9. (a) (i) Furthest line on left hand side. [1]
- (ii) The (electron) energy levels of a hydrogen atom become closer. [1]
- (b) (i) If a system at equilibrium is subject to a change the equilibrium tends to shift so as to minimize the effect of the change. [1]
- (ii) I Yield increases. (1)  
Forward reaction is endothermic. (1) [2]
- II Yield decreases. (1)  
More (gaseous) molecules on the right hand side. (1) [2]
- (iii) Atom economy =  $\frac{\text{mass hydrogen}}{\text{mass reactants}} \times 100$  (1)  
= 17.8% (1) [2]
- (c) Bonds broken = 3296 kJ                      Bonds formed = 3132 kJ (1)  
 $\Delta H = 3296 - 3132 = 164 \text{ kJ mol}^{-1}$  (1) [2]

Total [11]

10. (a) To ensure that the (initial) temperature is constant / temperature difference is required between initial and maximum temperature. [1]
- (b) (i) Best fit lines (1)  
 Temperature rise = 9.6°C (1) [2]  
 (Accept  $\pm 0.2^\circ\text{C}$ )
- (ii) Extrapolation gives the temperature that would have been reached if the reaction occurred instantly / to allow for heat loss during the experiment [1]
- (c) Heat =  $50 \times 4.18 \times 9.6$   
 = 2006 J [1]
- (d) (i) Moles Mg = 0.037 [1]
- (ii) Moles  $\text{CuSO}_4$  = 0.025 [1]
- (e)  $\Delta H = \frac{2006}{0.025}$  (1)  
 =  $-80.2 \text{ kJ mol}^{-1}$  (1) [2]
- (f) Burette / pipette [1]
- (g) Magnesium was in excess. [1]
- (h) Rate of reaction is quicker. Allow greater surface area if qualified. [1]
- (i)  $\frac{12.9}{93.1} \times 100 = 13.9\%$  [1]
- (j) Energy/Heat is lost to the environment. (1)  
 States how insulation could be improved e.g. place a lid on the polystyrene cup (1) [2]

Total [15]

Section B Total [70]