

# **GCE MARKING SCHEME**

CHEMISTRY AS/Advanced

**SUMMER 2015** 

#### **INTRODUCTION**

The marking schemes which follow were those used by WJEC for the Summer 2015 examination in GCE CHEMISTRY. They were finalised after detailed discussion at examiners' conferences by all the examiners involved in the assessment. The conferences were held shortly after the papers were taken so that reference could be made to the full range of candidates' responses, with photocopied scripts forming the basis of discussion. The aim of the conferences was to ensure that the marking schemes were interpreted and applied in the same way by all examiners.

It is hoped that this information will be of assistance to centres but it is recognised at the same time that, without the benefit of participation in the examiners' conferences, teachers may have different views on certain matters of detail or interpretation.

WJEC regrets that it cannot enter into any discussion or correspondence about these marking schemes.

	Page
CH1	1
CH2	7
CH4	13
CH5	20

## CH1

# **SECTION A**

1.	Ne	10p, 10n, 10e (1)	
	O <sup>2-</sup>	8p, 10n, 10e (1)	[2]
2.	(a)	<sup>222</sup> Rn	[1]
	(b)	Time taken for half of the atoms in a radioisotope to decay (or similar)	[1]
	(c)	Mass = $0.25 g (1)$	
		Moles = $1.11 \times 10^{-3}$ (1) do not accept $1 \times 10^{-3}$	[2]
3.	(a)	The mass of one mole of compound	[1]
	(b)	$\Delta H_f = -417 \text{ kJ mol}^{-1}$	[1]
4.	(a)	Measure the volume of $\mathrm{CO}_2$ produced / mass of $\mathrm{CO}_2$ lost at constant time intervals	[1]
	(b)	No effect since concentration of acid has not changed	[1]

**Total Section A [10]** 

## **SECTION B**

0

(ii) 
$$2NH_3 + 2O_2 \longrightarrow N_2O + 3H_2O$$
 [1]

(iii) Moles 
$$Ca(NO_3)_2 = 5.40 \times 10^{-3}$$
 (1) Moles gas =  $1.35 \times 10^{-2}$  (1) Volume gas =  $0.324$  dm<sup>3</sup> (1) [3]

(c) Moles 
$$Ca(NO_3)_2 = 0.0256$$
 (1)   
Moles  $H_2O = 0.102$  (1)   
 $x = 4$  (1) [3]

**Total** [11]

6.	(a)	(i)	Energy required to remove one mole of electrons from one mole of atoms / to form one mole of positive ions from one mole of atoms (	1)
			in the gaseous state (to form 1 mol of gaseous ions) (1) (Accept correct equation)	[2]
		(ii)	Cross between Na and Mg crosses	[1]
		(iii)	P only has unpaired electrons, S has a pair of electrons in 3p	
			orbital (1) Repulsion between the paired electrons makes it easier to remove one of the electrons (1)	[2]
	(b)	(i)	Effective nuclear charge is greater / electron being removed from a positive ion	[1]
		(ii)	Accept from 6000 to 9000	[1]
	(c)	energy Falling	are formed from electron being excited and jumping up to a higher level (1) back down to the n = 2 level (1) g energy / photon of light (1)	
		Lines b	pecome closer since the electron energy levels of a hydrogen atom e closer (1)	[4]
			Selection of a form and style of writing appropriate to purpose and to exity of subject matter	) [1]

Total [12]

7. (a) (i) Sample is bombarded by high energy electrons / electron gun used on sample (1)
Electron knocked out (to form ions) (1) [2]

(b) (i) 
$$(7.25 \times 6) + (92.75 \times 7)$$
 (1)  $100$  6.928 (1) (accept 6.93) [2]

(c) (i) 
$$M_r(NH_4)_2SO_4 = 132.18$$
 (1) Moles = 0.0156 (1) [2]

(ii) Moles LiOH = 
$$0.0312$$
 (1)

Concentration =  $0.0312 = 1.05 \text{ mol dm}^{-3}$  (1) [2]

(iii) Atom economy = 
$$\frac{34.06}{180.08} \times 100$$
 (1)  
=  $18.9\%$  (1) [2]

**Total [14]** 

## 8. (a) Benefits:

Stops fossil fuels from running out

Reduces CO<sub>2</sub> emissions / greenhouse emissions / global warming / effect of global warming

Reduces SO<sub>2</sub> emissions / acid rain

There will be an investment in new technology

#### Difficulties:

Dependence on fossil fuel/Unlikely to meet current demand

Renewable energy currently more expensive

Reliability of supply from renewables

Major development in energy efficiency technologies required

Opposition by vested interests

(Maximum 3 marks from list, but need examples of both) (3)

Consideration and discussion of benefits/difficulties (1) [4]


QWC Legibility of text; accuracy of spelling, punctuation and grammar, clarity of meaning [1]

- (b) (i) I As temperature increases yield decreases

  As pressure increases yield decreases [1]
  - II As temperature is increased, equilibrium moves to the left (1)
    Therefore forward reaction is exothermic (1)
    As pressure is increased, equilibrium moves to the left (1)
    Therefore more gas moles in products (1)

    QWC The information is organised clearly and coherently, using specialist vocabulary where appropriate

    [1]
  - (ii) If temperature is too low, then reaction is too slow (1)
    If temperature is too high, yield is too low (1)
    Compromise temperature acceptable rate and yield (1)
    (Accept any two points)
    [2]
  - (iii) Heterogenous catalyst [1]
  - (iv) Lower temperatures could be used (1)
    Less energy consumption / increased yield (1)
    Equilibrium could be reached more quickly (1)
    (Accept any two points) [2]



(vi) 
$$\Delta H = E_f - E_b$$
 [1]

Total [19]

9.	(a)	Otherwise a temperature change would occur on adding the acid which h nothing to do with the reaction		
	(b)	(i) Best fit lines (1)		
		Temperature rise = 6.4 °C (1) (Take value from candidate's best fit lines)	[2]	
		(ii) Volume of acid = 26.0 cm <sup>3</sup>	[1]	
		[If no best fit lines award 0 in (i) and accept 25 cm <sup>3</sup> in (ii)]		
	(c)	Moles acid = 0.02425 (1)		
		Conc acid = $\frac{0.02425}{0.026}$ = 0.933 mol dm <sup>-3</sup> (1)	[2]	
	(d)	Heat = 51 × 4.18 x 6.4		
		= 1364 J	[1]	
	(e)	$\Delta H = -\frac{1364}{0.02425}$ (1)		
		$= -56.2 \text{ kJ mol}^{-1}$ (1)	[2]	
	(f)	Pipette / burette	[1]	
	(g)	No further reaction occurs (1)		
		The excess acid cools the solution (1)	[2]	
	(h)	Heat / energy is lost to the environment (1)		
		Insulation is improved e.g. lid on the polystyrene cup (1)	[2]	
		Т	otal [14]	
		Section B T	otal [70]	