Q1. There is a link between the properties of the oxides of the Period 3 elements and their structure and bonding. The table below shows the melting points of the oxides of some Period 3 elements.

	Na₂O	SiO <sub>2</sub>	P <sub>4</sub> O <sub>10</sub>
T <sub>m</sub> /K	1548	1883	573

- (c) Phosphorus(V) oxide has a lower melting point than sodium oxide.
  - (i) State the structure of and bonding in phosphorus(V) oxide.

(2)

		(ii)	Explain why the melting point of phosphorus(V) oxide is low.	
				(1)
	(d)	water In eacti	ch case, predict the pH of the solution formed and write an equation fo ion.	
			ith P₄O₁₀	
		-	tion	
			ith Na <sub>2</sub> O	
		Equa	tion	(4)
	(e)		e an equation for the reaction between Na $_2$ O and P $_4$ O $_0$ the general type of reaction illustrated by this example.	
		Equa	tion	
		Reac	tion type	(2)
				(Total 16 marks)
Q2.		(a) s	State and explain the trend in electronegativities across Period 3 from fur.	sodium

Explain why the oxides of the Period 3 elements sodium and phosphorus have different melting points. In your answer you should discuss the structure of and bonding in these oxides, and the link between electronegativity and the type of bonding.  A chemical company has a waste tank of volume 25 000 dm². The tank is full of phosphoric acid (H,PO,) solution formed by adding some unwanted phosphorus(V) oxide to water in the tank.  A 25.0 cm² sample of this solution required 21.2 cm² of 0.500 mol dm² sodium hydroxide solution for complete reaction.  Calculate the mass, in kg, of phosphorus(V) oxide that must have been added to the water in the waste tank.	
A chemical company has a waste tank of volume 25 000 dm³. The tank is full of phosphoric acid (H,PO,) solution formed by adding some unwanted phosphorus(V) oxide to water in the tank.  A 25.0 cm³ sample of this solution required 21.2 cm³ of 0.500 mol dm³ sodium hydroxide solution for complete reaction.  Calculate the mass, in kg, of phosphorus(V) oxide that must have been added to the water in the waste tank.	
A chemical company has a waste tank of volume 25 000 dm³. The tank is full of phosphoric acid (H.PO.) solution formed by adding some unwanted phosphorus(V) oxide to water in the tank.  A 25.0 cm³ sample of this solution required 21.2 cm³ of 0.500 mol dm³ sodium hydroxide solution for complete reaction.  Calculate the mass, in kg, of phosphorus(V) oxide that must have been added to the water in the waste tank.	
phosphoric acid (H₃PO₄) solution formed by adding some unwanted phosphorus(V) oxide to water in the tank.  A 25.0 cm³ sample of this solution required 21.2 cm³ of 0.500 mol dm⁻³ sodium hydroxide solution for complete reaction.  Calculate the mass, in kg, of phosphorus(V) oxide that must have been added to the water in the waste tank.	different melting points. In your answer you should discuss the structure of and bonding in these oxides, and the link between electronegativity and the type of
phosphoric acid (H₃PO₄) solution formed by adding some unwanted phosphorus(V) oxide to water in the tank.  A 25.0 cm³ sample of this solution required 21.2 cm³ of 0.500 mol dm⁻³ sodium hydroxide solution for complete reaction.  Calculate the mass, in kg, of phosphorus(V) oxide that must have been added to the water in the waste tank.	
phosphoric acid (H₃PO₄) solution formed by adding some unwanted phosphorus(V) oxide to water in the tank.  A 25.0 cm³ sample of this solution required 21.2 cm³ of 0.500 mol dm⁻³ sodium hydroxide solution for complete reaction.  Calculate the mass, in kg, of phosphorus(V) oxide that must have been added to the water in the waste tank.	
phosphoric acid (H₃PO₄) solution formed by adding some unwanted phosphorus(V) oxide to water in the tank.  A 25.0 cm³ sample of this solution required 21.2 cm³ of 0.500 mol dm⁻³ sodium hydroxide solution for complete reaction.  Calculate the mass, in kg, of phosphorus(V) oxide that must have been added to the water in the waste tank.	
phosphoric acid (H₃PO₄) solution formed by adding some unwanted phosphorus(V) oxide to water in the tank.  A 25.0 cm³ sample of this solution required 21.2 cm³ of 0.500 mol dm⁻³ sodium hydroxide solution for complete reaction.  Calculate the mass, in kg, of phosphorus(V) oxide that must have been added to the water in the waste tank.	
phosphoric acid (H₃PO₄) solution formed by adding some unwanted phosphorus(V) oxide to water in the tank.  A 25.0 cm³ sample of this solution required 21.2 cm³ of 0.500 mol dm⁻³ sodium hydroxide solution for complete reaction.  Calculate the mass, in kg, of phosphorus(V) oxide that must have been added to the water in the waste tank.	
phosphoric acid (H₃PO₄) solution formed by adding some unwanted phosphorus(V) oxide to water in the tank.  A 25.0 cm³ sample of this solution required 21.2 cm³ of 0.500 mol dm⁻³ sodium hydroxide solution for complete reaction.  Calculate the mass, in kg, of phosphorus(V) oxide that must have been added to the water in the waste tank.	
phosphoric acid (H₃PO₄) solution formed by adding some unwanted phosphorus(V) oxide to water in the tank.  A 25.0 cm³ sample of this solution required 21.2 cm³ of 0.500 mol dm⁻³ sodium hydroxide solution for complete reaction.  Calculate the mass, in kg, of phosphorus(V) oxide that must have been added to the water in the waste tank.	
phosphoric acid (H₃PO₄) solution formed by adding some unwanted phosphorus(V) oxide to water in the tank.  A 25.0 cm³ sample of this solution required 21.2 cm³ of 0.500 mol dm⁻³ sodium hydroxide solution for complete reaction.  Calculate the mass, in kg, of phosphorus(V) oxide that must have been added to the water in the waste tank.	
hydroxide solution for complete reaction.  Calculate the mass, in kg, of phosphorus(V) oxide that must have been added to the water in the waste tank.	phosphoric acid (H <sub>3</sub> PO <sub>4</sub> ) solution formed by adding some unwanted phosphorus(V)
water in the waste tank.	

				(5)
			(To	tal 15 marks)
Q3.		Consi	ider the following oxides.	
		Na₂C	$O$ , $MgO$ , $Al_2O_3$ , $SiO_2$ , $P_4O_{10}$ , $SO_3$	
	(a)	lder	ntify one of the oxides from the above which	
		(i)	can form a solution with a pH less than 3	
		(ii)	can form a solution with a pH greater than 12	(2)
				(2)
	(b)	Writ	te an equation for the reaction between	
		(i)	MgO and HNO₃	
		(ii)	SiO₂ and NaOH	
		(iii)	Na₂O and H₃PO₄	
				(3)
		_		
	(c)		lain, in terms of their type of structure and bonding, why $P_4O_{10}$ can be vapo entle heat but $SiO_2$ cannot.	rised

Page 5

••••							
							(Total 9 mai
. (a)	The melt	ina points of	f some of th	ne oxides for	rmed bv Per	riod 3 elemen	its are
` '		dom order b		10 0/1100 101	illiou by 1 of	iod o diomion	ito di o
Oxide	Α	В	С	D	E		
T <sub>m</sub> /°C	2852	-73	1610	1275	300	1	
(i)	structure Explain	es. your answe	r.		hich have s	imple molecu	ular 
	Explana	ntion					
(ii)						ow which of t would make.	
(ii)	in the ta	ble is sodiu	m oxide. Sta	ate the obse	ervation you		
(ii)	in the ta	ble is sodiu	m oxide. Sta	ate the obse	ervation you	would make.	

(b)	The base calcium oxide can be used to remove sulfur dioxide from flue-gases produced when fossil fuels are burnt in coal-fired power stations. Calcium oxide is produced when calcium carbonate, is decomposed by heat.						
	(i)	Write an equation for the action of heat on calcium carbonate.					
	(ii)	Identify the product formed when sulfur dioxide reacts with calcium oxide.					
	/iii\	Despite the additional cost, energtors of power stations are encouraged to					
	(iii)	Despite the additional cost, operators of power stations are encouraged to remove the sulfur dioxide from flue-gases. Explain why this may not be environmentally beneficial.					
		(4) (Total 10 marks)					