Q1. (a)	Some metal ions are toxic to humans. A substance that can be used to treat such poisoning contains the ion EDTA ⁴⁻ . EDTA ⁴⁻ forms very stable complexes with metal ions. These complexes are not toxic.			
	(i)	Write an equation for the reaction of EDTA ⁴⁻ with aqueous copper(II) ions, $[Cu(H_2O)_6]^{2^*}$.		
			(1)	
	(ii)	A solution containing EDTA ⁴⁻ can also be used in a titration to determine the concentration of metal ions in solution. A river was polluted with copper(II) ions. When a 25.0 cm³ sample of the river water was titrated with a 0.0150 mol dm⁻³ solution of EDTA⁴-, 6.45 cm³ were required for complete reaction. Calculate the concentration, in mol dm⁻³, of copper(II) ions in the river water. Show your working.		
			(2)	
(b)	wate	e determination of the concentration of copper(II) ions in a single sample of river er gives an unreliable value for the copper(II) ion pollution in the river. e one reason why this value is unreliable.		
			(1)	
(c)	Give	ver complexes can be used to identify a particular organic functional group. e one example of a silver complex that can be used in this way and state the anic functional group it identifies.		
	Silv	er complex		
	Org	anic functional group(Total 6 m	(2) arks)	

Q2.		Transition elements form complex ions with a range of colours and shapes.	
	(a)	By considering its electron arrangement, state how an element can be classified as a transition element.	
			(1)
	(b)	Explain the meaning of the term <i>complex ion</i> .	
			(0)
			(2)
	(c)	In terms of electrons, explain why an aqueous solution of cobalt(II) sulfate has a red colour.	
			(3)
	(d)	The ligand EDTA⁴ is shown below.	
	~ (%)	$C - CH_2$ $H_2C - C$ $C - CH_2$ $H_2C - C$ $C - CH_2$ $H_2C - C$	

Draw circles around the atoms of **two** different elements that link to a

transition metal ion by a co-ordinate bond when EDTA⁴ behaves as a ligand.

(i)


Write an equation for the reaction between EDTA⁴ and a [Co(H₂O)₀]² ion. Use (ii) the abbreviation EDTA[←] in your equation.

(1)

(iii) Explain why the complex ion, formed as a product of the reaction in part (d) (ii), is more stable than the $[Co(H_2O)_6]^{2+}$ ion.

(2)

(e) The diagram below shows part of the structure of haemoglobin.




Haemoglobin contains an iron(II) ion bonded to five nitrogen atoms and one other ligand. The fifth nitrogen atom and the additional ligand are not shown in this diagram.

In this diagram, bonds between nitrogen and iron are shown as N→Fe and as (i) N—Fe.

State the meaning of each of these symbols.

		Meaning of →	
		Meaning of —	(2)
	(ii)	State the function of haemoglobin in the blood.	(4)
			(1)
	(iii)	With reference to haemoglobin, explain why carbon monoxide is tox	ic.
			(2)
			(Total 16 marks)
00	T		. former d
Q3.	by transitiona lineara tetral	sition metals form complex ions. Using actual examples of complex ions on metal ions, give the formula of r complex ion, nedral complex ion and	s formed
	an octa	ahedral complex ion formed by using a bidentate ligand.	(Total 4 marks)

Q4. This question is about some reactions of cobalt compounds.



(e)	Give the formula of the cobalt-containing complex in \mathbf{Y} and explain why the complex is more stable than the cobalt-containing complex in \mathbf{W} .	nis
	Formula	
	Explanation	
		 (3)
		(-7
(5)		
(f)	Identify the cobalt-containing complex in solution Z and explain why its co-ordination number is different from that in the pink solution of CoCl ₂	
	Complex	
	Explanation	
		(2)
		(Total 13 marks)