

**Q1.**This question is about copper chemistry.

(a) Aqueous copper(II) ions  $[\text{Cu}(\text{H}_2\text{O})_6]^{2+}(\text{aq})$  are blue.

(i) With reference to electrons, explain why aqueous copper(II) ions are blue.

.....  
.....  
.....  
.....  
.....  
.....  
.....

*(Extra space)* .....  
.....  
.....

**(3)**

(ii) By reference to aqueous copper(II) ions, state the meaning of each of the **three** terms in the equation  $\Delta E = h\nu$ .

.....  
.....  
.....  
.....  
.....  
.....  
.....

*(Extra space)* .....  
.....  
.....

**(3)**

(iii) Write an equation for the reaction, in aqueous solution, between  $[\text{Cu}(\text{H}_2\text{O})_6]^{2+}$  and an excess of chloride ions.

State the shape of the complex produced and explain why the shape differs from that of the  $[\text{Cu}(\text{H}_2\text{O})_6]^{2+}$  ion.

.....  
.....  
.....  
.....  
.....  
.....  
.....  
(Extra space) .....  
.....  
.....

(3)

- (b) Draw the structure of the ethanedioate ion ( $\text{C}_2\text{O}_4^{2-}$ ). Explain how this ion is able to act as a ligand.

.....  
.....  
.....  
.....  
.....

(2)

- (c) When a dilute aqueous solution containing ethanedioate ions is added to a solution containing aqueous copper(II) ions, a substitution reaction occurs. In this reaction four water molecules are replaced and a new complex is formed.

- (i) Write an ionic equation for the reaction. Give the co-ordination number of the complex formed and name its shape.

.....  
.....  
.....  
.....  
.....  
.....

(4)

- (ii) In the complex formed, the two water molecules are opposite each other. Draw a diagram to show how the ethanedioate ions are bonded to a copper ion and give a value for one of the O–Cu–O bond angles. You are **not** required to show the water molecules.

(2)


(Total 17 marks)

**Q2.** Complexes containing transition elements have a wide variety of uses including acting as dyestuffs like *Prussian Blue*.

*Cisplatin* is a platinum-based chemotherapy drug used to treat various types of cancers. It was the first member of a class of anti-cancer drugs that react with DNA in tumour cells.

*Cisplatin* is prepared from  $K_2PtCl_4$  according to the following scheme.

**All the reactions shown are reversible.**



(a) Name the type of reaction occurring in all four steps of the scheme.

.....

(1)

(b) Explain why an excess of potassium iodide is used in Reaction 1.

.....  
 .....  
 .....

(2)

(c) (i) Write an equation for Reaction 1.

.....  
 .....

(1)

(ii) Calculate the percentage atom economy for the formation of  $\text{K}_2\text{PtI}_4$  in Reaction 1. Show your working.

.....  
 .....  
 .....

.....

(2)

(d) In Reaction 3, silver nitrate solution is added to improve the yield of product.

(i) Write the **simplest ionic** equation for the reaction of iodide ions with silver nitrate.

.....

(1)

(ii) Suggest why addition of silver nitrate improves the yield of product from Reaction 3.

.....

.....

(1)

(e) Suggest two reasons, other than poor practical technique, why the overall yield of *cisplatin* in this synthesis may be low.

Reason 1 .....

.....

Reason 2 .....

.....

(2)

(f) The *cisplatin* formed in Reaction 4 is impure. Outline how the impure solid is purified by recrystallisation.

.....

.....

.....

.....

.....

(3)

(g) Platinum compounds are highly toxic.

(i) State why *cisplatin* is used in cancer treatment despite its toxicity.

.....  
.....

(1)

(ii) Suggest a suitable precaution that should be taken by medical staff when using *cisplatin*.

.....

(1)

(Total 15 marks)

**Q3.(a)** Some metal ions are toxic to humans. A substance that can be used to treat such poisoning contains the ion  $\text{EDTA}^{4-}$ .  
 $\text{EDTA}^{4-}$  forms very stable complexes with metal ions. These complexes are **not** toxic.

(i) Write an equation for the reaction of  $\text{EDTA}^{4-}$  with aqueous copper(II) ions,  $[\text{Cu}(\text{H}_2\text{O})_6]^{2+}$ .

.....

(1)

(ii) A solution containing  $\text{EDTA}^{4-}$  can also be used in a titration to determine the concentration of metal ions in solution.  
A river was polluted with copper(II) ions. When a  $25.0 \text{ cm}^3$  sample of the river water was titrated with a  $0.0150 \text{ mol dm}^{-3}$  solution of  $\text{EDTA}^{4-}$ ,  $6.45 \text{ cm}^3$  were required for complete reaction.  
Calculate the concentration, in  $\text{mol dm}^{-3}$ , of copper(II) ions in the river water.  
Show your working.

.....  
.....  
.....  
.....

(2)

- (b) The determination of the concentration of copper(II) ions in a single sample of river water gives an unreliable value for the copper(II) ion pollution in the river. Give one reason why this value is unreliable.

.....  
.....

(1)

- (c) Silver complexes can be used to identify a particular organic functional group. Give **one** example of a silver complex that can be used in this way and state the organic functional group it identifies.

Silver complex .....

Organic functional group .....

(2)

(Total 6 marks)