

Allow multiples, including fractions.



Allow correct equation which includes water of crystallisation.

1

(b) $M_r \text{FeSO}_4 \cdot 7\text{H}_2\text{O} = 277.9$

Allow if shown clearly in the calculation.

Allow 278

1

Moles = $6.95 / 277.9 = 2.5(0) \times 10^{-2}$

Do not penalise precision but must be to a minimum of two significant figures.

Allow correct calculation using incorrect M_r .

Correct answer without working scores this mark only.

1

(c) $3(.00) \times 10^{-2}$

1

(d) Theoretical mass = $2.50 \times 10^{-2} \times 179.8 = 4.50\text{g}$

as long as 2.50×10^{-2} is the smaller of parts (b) and (c) **(M1)**

Allow consequential answer from parts (b) and (c).

Allow theoretical mass = (smaller of parts (b) and (c)) \times 179.8

*If larger of parts (b) and (c) used, lose **M1** but can score **M2**.*

Allow answers based on moles of reactant and product.

1

Yield = $3.31 \times 100 / 4.50 = 73.6\%$ **(M2)**

Award this mark only if answer given to 3 significant figures.

Correct answer without working scores this mark only, provided answer given to 3 significant figures.

1

(e) Some left in solution / some lost during filtration

Do not allow 'incomplete reaction'.

Do not allow 'reaction is reversible'.

1

(f) MnO_4^- will oxidise the iron(II) ion and the ethanedioate ion

1

MnO_4^- does not oxidise the Cu^{2+} ion / larger volume needed for iron(II) ethanedioate

1

[9]

M2.(a) A ligand is an electron pair / lone pair donor

Allow uses lone / electron pair to form a co-ordinate bond

1

A bidentate ligand donates two electron pairs (to a transition metal ion) from different atoms / two atoms (on the same molecule / ion)

QoL

1

(b) CoCl_4^{2-} diagram


1

Tetrahedral shape

1

$109^\circ 28'$

1



*Four chlorines attached to Co with net 2- charge correct
Charge can be placed anywhere, eg on separate formula
Penalise excess charges*

Allow 109° to 109.5°

[Co(NH₃)₆]²⁺ diagram


1

Octahedral shape

1

90°

1



Six ammonia / NH₃ molecules attached to Co with 2+ charge correct

Allow 180° if shown clearly on diagram

CE= 0 if wrong complex but mark on if only charge is incorrect

- (c) In different complexes the d orbitals / d electrons (of the cobalt) will have different energies / d orbital splitting will be different

1


Light / energy is absorbed causing an electron to be excited

1

Different frequency / wavelength / colour of light will be absorbed / transmitted / reflected

1

- (d) 1 mol of H₂O₂ oxidises 2 mol of Co²⁺



1

M_r CoSO₄·7H₂O = 281

If M_r wrong, max 3 for M1, M4, M5

1

$$\text{Moles Co}^{2+} = 9.87 / 281 = 0.03512$$

1

$$\text{Moles H}_2\text{O}_2 = 0.03512 / 2 = 0.01756$$

M4 is method mark for (M3) / 2 (also scores M1)

1

$$\begin{aligned} \text{Volume H}_2\text{O}_2 &= (\text{moles} \times 1000) / \text{concentration} \\ &= 0.01756 \times 1000 / 5.00 \end{aligned}$$

$$= 3.51 \text{ cm}^3 / (3.51 \times 10^{-3} \text{ dm}^3)$$

Units essential for answer

M5 is method mark for (M4) x 1000 / 5

Allow 3.4 to 3.6 cm³

If no 2:1 ratio or ratio incorrect Max 3 for M2, M3 & M5

Note: Answer of 7 cm³ scores 3 for M2, M3, M5 (and any other wrong ratio max 3)

Answer of 16.8 cm³ scores 3 for M1, M4, M5 (and any other wrong M, max 3)

Answer of 33.5 cm³ scores 1 for M5 only (so wrong M, AND wrong ratio max 1)

1

[16]

M3.(a) Orange dichromate

Allow max 2 for three correct colours not identified to species but in correct order

1

Changes to purple / green / ruby / red-violet / violet Chromium(III)
(Note green complex can be $[\text{Cr}(\text{H}_2\text{O})_5\text{Cl}]^{2+}$ etc)



Do not allow green with another colour

1

That changes further to blue Chromium(II)


Allow max 1 for two correct colours not identified but in correct order

1



Ignore any further reduction of Cr²⁺

1



Ignore additional steps e.g. formation of CrO₄²⁻

1


(b) Green precipitate

1

(Dissolves to form a) green solution


Solution can be implied if 'dissolves' stated

1



Penalise Cr(OH)₃ once only

1



Allow [Cr(H₂O)₆]³⁺ + 6OH⁻ → [Cr(OH)₆]³⁻ + 6H₂O

Allow formation of [Cr(H₂O)₂(OH)₄]⁺ and [Cr(H₂O)(OH)₅]²⁻ in balanced equations

Ignore state symbols, mark independently

1

(c) (ligand) substitution / replacement / exchange

Allow nucleophilic substitution

1

The energy levels/gaps of the d electrons are different (for each complex)

Ignore any reference to emission of light

1

So a different wavelength/frequency/colour/energy of light is absorbed (when d electrons are excited)

OR light is absorbed and a different wavelength/frequency/colour/energy (of light) is transmitted/reflected


1

(d) $E_{O_2} (/ H_2O) > E_{Cr^{3+}} (/ Cr^{2+}) / e.m.f = 1.67 V$
Allow $E_{(cell)} = 1.67$

1

So Cr^{2+} ions are oxidised by oxygen/air

Allow any equation of the form:



1

With $[Cr(H_2O)_6]^{2+}$ get $CrCO_3$

If named must be chromium(II) carbonate

1

with $[Cr(H_2O)_6]^{3+}$ get $Cr(H_2O)_3(OH)_3 / Cr(OH)_3$

Allow 0 to 3 waters in the complex

1

and CO_2

Can score M3, M4, M5 in equations even if unbalanced

1

$Cr(III)$ differs from $Cr(II)$ because it is acidic / forms H^+ ions

1

because Cr^{3+} ion polarises water

Ignore charge/size ratio and mass/charge

1

[19]

- M4.(a)** Co-ordinate / dative / dative covalent / dative co-ordinate
Do not allow covalent alone 1
- (b) (lone) pair of electrons on oxygen/O
If co-ordination to O²⁻, CE=0 1
- forms co-ordinate bond with Fe / donates electron pair to Fe
'Pair of electrons on O donated to Fe' scores M1 and M2 1
- (c) 180° / 180 / 90
Allow any angle between 85 and 95
Do not allow 120 or any other incorrect angle
Ignore units eg °C 1
- (d) (i) 3 : 5 / 5 FeC₂O₄ reacts with 3 MnO₄⁻
Can be equation showing correct ratio 1
- (ii) **M1** Moles of MnO₄⁻ per titration = $22.35 \times 0.0193/1000 = 4.31 \times 10^{-4}$
 Method marks for each of the next steps (no arithmetic error allowed for M2):
Allow 4.3×10^{-4} (2 sig figs)
Allow other ratios as follows:
eg from given ratio of 7/3 1
- M2** moles of FeC₂O₄= ratio from (d)(i) used correctly $\times 4.31 \times 10^{-4}$
M2 = $7/3 \times 4.31 \times 10^{-4} = 1.006 \times 10^{-3}$ 1

M3 moles of FeC_2O_4 in $250 \text{ cm}^3 = \text{M2 ans} \times 10$

$$\mathbf{M3} = 1.006 \times 10^{-3} \times 10 = 1.006 \times 10^{-2}$$

1

M4 Mass of $\text{FeC}_2\text{O}_4 \cdot 2\text{H}_2\text{O} = \text{M3 ans} \times 179.8$

$$\mathbf{M4} = 1.006 \times 10^{-2} \times 179.8 = 1.81 \text{ g}$$

1

M5 % of $\text{FeC}_2\text{O}_4 \cdot 2\text{H}_2\text{O} = (\text{M4 ans}/1.381) \times 100$

$$\mathbf{M5} = 1.81 \times 100/1.381 = 131 \% \text{ (130 to 132)}$$

1

(OR for M4 max moles of $\text{FeC}_2\text{O}_4 \cdot 2\text{H}_2\text{O} = 1.381/179.8 (= 7.68 \times 10^{-3})$

for M5 % of $\text{FeC}_2\text{O}_4 \cdot 2\text{H}_2\text{O} = (\text{M3 ans}/\text{above M4ans}) \times 100$

eg using correct ratio 5/3:

$$\text{Moles of } \text{FeC}_2\text{O}_4 = 5/3 \times 4.31 \times 10^{-4} = 7.19 \times 10^{-4}$$

$$\text{Moles of } \text{FeC}_2\text{O}_4 \text{ in } 250 \text{ cm}^3 = 7.19 \times 10^{-4} \times 10 = 7.19 \times 10^{-3}$$

$$\text{Mass of } \text{FeC}_2\text{O}_4 \cdot 2\text{H}_2\text{O} = 7.19 \times 10^{-3} \times 179.8 = 1.29 \text{ g}$$

$$\% \text{ of } \text{FeC}_2\text{O}_4 \cdot 2\text{H}_2\text{O} = 1.29 \times 100/1.381 = 93.4 \text{ (allow 92.4 to 94.4)}$$

Note correct answer (92.4 to 94.4) scores 5 marks

Allow consequentially on candidate's ratio

$$\text{eg } \mathbf{M2} = 5/2 \times 4.31 \times 10^{-4} = 1.078 \times 10^{-3}$$

$$\mathbf{M3} = 1.0078 \times 10^{-3} \times 10 = 1.0078 \times 10^{-2}$$

$$\mathbf{M4} = 1.078 \times 10^{-2} \times 179.8 = 1.94 \text{ g}$$

$$\mathbf{M5} = 1.94 \times 100/1.381 = 140 \% \text{ (139 to 141)}$$

Other ratios give the following final % values

1:1 gives 56.1% (55.6 to 56.6)

5:1 gives 281% (278 to 284)

5:4 gives 70.2% (69.2 to 71.2)

[10]