

Mark schemes

Q1.

(a)

This question is marked using levels of response. Refer to the Mark Scheme Instructions for Examiners for guidance on how to mark this question.	
Level 3 5-6 marks	All stages are covered and the description of each stage is generally correct and virtually complete. Answer is communicated coherently and shows a logical progression from stage 1 to stage 2 and stage 3.
Level 2 3-4 marks	All stages are covered but the description of each stage may be incomplete or may contain inaccuracies OR two stages are covered and the explanations are generally correct and virtually complete. Answer is mainly coherent and shows progression from stage 1 to stage 2 and/or stage 3.
Level 1 1-2 marks	Two stages are covered but the description of each stage may be incomplete or may contain inaccuracies, OR only one stage is covered but the explanation is generally correct and virtually complete. Answer includes isolated statements and these are presented in a logical order.
Level 0	0 marks Insufficient correct chemistry to gain a mark.

Stage 1

1a Heterogeneous means in a different phase/state from reactants

1b Catalyst speeds up reaction and is left unchanged **OR** lowers the activation energy for the reaction

Stage 2

2a Hydrogen and nitrogen/reactants adsorb onto the surface/active sites of the iron

2b Bonds weaken/reaction takes place

2c Products desorb/leave from the surface (of the iron)

Stage 3

3a Large surface area (of iron) by using powder or small pellets or support medium/mesh

3b Catalyst poisoned / sulfur poisons or binds to the catalyst

3c Active sites blocked

Ignore references to temperature and pressure


6

(b) Two negative ions repel


1

So activation energy is high

1



1



Ignore any state symbols given

Allow multiples for both equations

Allow equations in either order

1

(c) (Zn ions) have only one oxidation state

Or

Zn²⁺ is the only ion

Allow doesn't have variable oxidation state

Allow cannot be oxidised to Zn³⁺

Ignore has a full d shell

1

(d) M1 Amount of Fe = $0.998 \div 55.8 = 0.0179$ mol

1

M2 Amount of HCl = 0.0300 mol

1

M3 HCl is the limiting reagent

1

M4 Amount of H₂ produced = 0.0150 mol

M4 = M2 ÷ 2

1

M5 T = 303 K P = 100 000 Pa

1

$$\text{M6 } V \left(= \frac{0.0150 \times 8.31 \times 303}{100\,000} \right) = 3.78 \times 10^{-4} \text{ (m}^3\text{)}$$

$$\text{M6 } V \left(= \frac{\text{M4} \times 8.31 \times 303}{100\,000} \right) \text{ (m}^3\text{)}$$

1

(e) FeCO₃ or iron(II) carbonate

1

Green

Allow white

1

(f) Fe(H₂O)₃(OH)₃


Ignore square brackets if added

1

- brown 1
- $2 [\text{Fe}(\text{H}_2\text{O})_6]^{3+} + 3 \text{CO}_3^{2-} \rightarrow 2 \text{Fe}(\text{H}_2\text{O})_3(\text{OH})_3 + 3 \text{H}_2\text{O} + 3 \text{CO}_2$
Accept multiples 1
- (g) M1 Fe^{3+} is smaller (than Fe^{2+}) **OR** Fe^{3+} has a greater charge
OR Fe^{3+} has a greater charge density **OR** Fe^{3+} has a greater charge to size ratio
Penalise $\text{Fe}(\text{H}_2\text{O})_6^{3+}$ ions once in M1 or M2 1
- M2 Fe^{3+} ions are more polarising **OR** Fe^{3+} ions polarise water molecules more 1
- M3 So more O-H bonds (in the water ligands) break **OR** more H^+ ions released **OR** weaken O-H bonds in ligands more (in the Fe^{3+} solution)
Do not allow Fe^{3+} releases 3H^+ ions 1
- [25]**

Q2.

- (a) **M1** **B** = $\text{Al}(\text{H}_2\text{O})_3(\text{OH})_3$
Ignore [] 1
- M2** bubbles/effervescence
M2 *Do not allow gas evolved* 1
- M3** $2 [\text{Al}(\text{H}_2\text{O})_6]^{3+} + 3 \text{CO}_3^{2-} \rightarrow 2 \text{Al}(\text{H}_2\text{O})_3(\text{OH})_3 + 3 \text{H}_2\text{O} + 3 \text{CO}_2$
M3 *Ignore absence of square brackets around Al complex*
M3 *Allow correct balanced equations with Na_2CO_3* 1
- (b) **M1** **C** = $[\text{Al}(\text{OH})_4]^-$ **OR** $[\text{Al}(\text{H}_2\text{O})_2(\text{OH})_4]^-$ **OR** $[\text{Al}(\text{OH})_6]^{3-}$ 1
- M2** Excess NaOH
M2 *Allow excess OH* 1
- M3** $[\text{Al}(\text{H}_2\text{O})_6]^{3+} + 4 \text{OH}^- \rightarrow [\text{Al}(\text{OH})_4]^- + 6 \text{H}_2\text{O}$
 OR
 $[\text{Al}(\text{H}_2\text{O})_6]^{3+} + 4 \text{OH}^- \rightarrow [\text{Al}(\text{H}_2\text{O})_2(\text{OH})_4]^- + 4 \text{H}_2\text{O}$
 OR



M3 Allow equations to form $\text{Al}(\text{H}_2\text{O})(\text{OH})_5^{2-}$


M3 Allow correct balanced equations with NaOH

1



Do not penalise absence of square brackets

1



Accept other equations

1


M2 Al^{3+} has a small size and high charge OR has a high charge density

1

M3 Weakens the OH bond (in water) releasing H^+ ions

M2 Allow the aluminium ion has a small size and high charge OR has a high charge density

1


[10]**Q3.**

1

Brown

M2: Allow red-brown


1




M3: Allow correct equations with Na_2CO_3

M3: Ignore State symbols

1



1




M2: Allow correct equations with HCl

1



Allow KI/potassium iodide

1



1

green

1



This question is marked using levels of response. Refer to the Mark Scheme Instructions for Examiners for guidance on how to mark this question.	
Level 3 5-6 marks	All stages are covered and the description of each stage is generally correct and virtually complete. Answer is communicated coherently and shows a logical progression from stage 1 to stage 2 and stage 3 Answer is illustrated using diagrams of at least 2 specific examples of pairs of cobalt or platinum complex isomers.
Level 2 3-4 marks	All stages are covered but the description of each stage may be incomplete or may contain inaccuracies OR two stages are covered and the explanations are generally correct and virtually complete. Answer is mainly coherent and shows progression from stage 1 to stage 2 and/or stage 3. Answer is illustrated using diagrams of at least 1 specific example of a pair of cobalt or platinum complex isomers.
Level 1 1-2 marks	Two stages are covered but the description of each stage may be incomplete or may contain inaccuracies, OR only one stage is covered but the explanation is generally correct and virtually complete. Answer includes isolated statements and these are presented in a logical order. Answer is illustrated using at least 1 appropriate diagram or formula.
0 marks	Insufficient correct chemistry to gain a mark.

Indicative Chemistry content**Stage 1: shapes of complexes**

1a octahedral or 6 co-ordinate diagram

1b tetrahedral or square planar or 4 co-ordinate diagram

Stage 2: cis/ trans isomerism (or E-Z or geometric)

2a cis/trans isomerism in either square planar and/or octahedral complexes

2b Diagrams showing cis and trans isomerism in a square planar complex2c Diagrams showing cis and trans isomerism in both isomers of octahedral complexes eg draw cis and trans $M(H_2O)_4(OH)_2$ or $[M(NH_3)_4(H_2O)_2]^{2+}$


Stage 3: optical isomerism

3a optical isomerism / non superimposable mirror images in octahedral complexes


3b occurs with a specific bidentate ligands eg. $C_2O_4^{2-}$ or $NH_2CH_2CH_2NH_2$

3c draw both optical isomers of eg $[M(NH_2CH_2CH_2NH_2)_3]^{2+}$

6


[14]**Q4.**

Allow



Or equation to form $Al(H_2O)_4(OH)_2^+$

1



allow equations to form $[Al(H_2O)_4(OH)_2]^+$

1


(c) white ppt/solid

M1 and M2 in either order

1

effervescence/bubbles/fizzing

1




accept multiples

only allow spectator ions in a balanced equation

1

(d) White ppt/solid

1




only allow spectator ions in a balanced equation


1

Colourless solution forms / ppt or solid dissolves

1



OR



only allow 6 or 4 co-ordination

Allow $[Al(OH)_6]^{3-}$ in a balanced equation

1

[9]


Q5.

- (a) **A** Silver bromide / AgBr 1
- B** Iron(II) carbonate / FeCO₃ 1
- C** Iron(II) sulphate / FeSO₄ 1
- D** Carbon dioxide / CO₂ 1
- Y** Iron(II) bromide / FeBr₂ 1
- (b) $\text{Ag}^+ + \text{Br}^- \rightarrow \text{AgBr}$
 Allow equation if state symbols missing but penalise if state symbols are incorrect 1
- (c) $2\text{H}^+ + \text{CO}_3^{2-} \rightarrow \text{H}_2\text{O} + \text{CO}_2$
 Allow $\text{FeCO}_3 + 2\text{H}^+ \rightarrow \text{Fe}^{2+} + \text{CO}_2 + \text{H}_2\text{O}$ 1

[7]

Q6.

- (a) $[\text{Fe}(\text{H}_2\text{O})_6]^{3+} + 4\text{Cl}^- \rightarrow \text{FeCl}_4^- + 6\text{H}_2\text{O}$ 1
- (b) Cl⁻ is a bigger ligand 1
- So only 4Cl⁻ can fit around the metal
 Allow fewer Cl⁻ can fit around the metal 1



- (c) M1 for structure of complex

M2 for correct charge

(d) Change in entropy is positive

1
1(e) $5\text{Fe}^{2+} + \text{MnO}_4^- + 8\text{H}^+ \rightarrow \text{Mn}^{2+} + 5\text{Fe}^{3+} + 4\text{H}_2\text{O}$ 1
1(f) Amount of manganate (VII) = 6.50×10^{-4} mol

1

Amount of iron(II) = 3.25×10^{-3} mol*ie M1 x 5*

1

Mass of iron = 0.181 g = 181 mg

Allow M2 x 55.8

1

Percentage Fe = $181/1980 \times 100 = \underline{9.14}(\%)$ 3 sf

1

(g) Colourless to pale pink

1

[12]**Q7.**

(a) An electron pair on the ligand

1

Is donated from the ligand to the central metal ion

1

(b) Blue precipitate


1

Dissolves to give a dark blue solution

1



1



1

(c) $[\text{Cu}(\text{NH}_3)_4(\text{H}_2\text{O})_2]^{2+} + 2\text{H}_2\text{NCH}_2\text{CH}_2\text{NH}_2 \longrightarrow$
 $[\text{Cu}(\text{H}_2\text{NCH}_2\text{CH}_2\text{NH}_2)_2(\text{H}_2\text{O})_2]^{2+} + 4\text{NH}_3$

1

(d) Cu–N bonds formed have similar enthalpy / energy to Cu–N bonds broken

1

And the same number of bonds broken and made

- (e) 3 particles form 5 particles / disorder increases because more particles are formed / entropy change is positive

1

1

Therefore, the free-energy change is negative

M2 can only be awarded if M1 is correct

1

[11]

Q8.

D

[1]

Q9.

B

[1]

Q10.

B

[1]

Q11.

- (a) $4\text{Fe(s)} + 6\text{H}_2\text{O(g)} + 3\text{O}_2\text{(g)} \rightarrow 4\text{Fe(OH)}_3\text{(s)}$

Accept fractions, multiples and equations showing formation of hydrated hydroxide.

1

Correct states as above

Lose this mark if any state is missing or incorrect.

1

- (b) Aluminium (metal) forms an oxide coat (on exposure to air)

1

This coat prevents / inhibits further reaction (by water or oxygen)

Ignore references to a 'less reactive coating' unless further qualified.

1

[4]