

F322: Alcohols

1. 2-Methylpropan-2-ol ✓

ALLOW methylpropan-2-ol

[1]

2. Has O–H (bonds)

OR has hydroxyl (groups) **OR** has hydroxy (groups) ✓

ALLOW marks from a diagram of hydrogen bonding

IGNORE reference to alcohol functional group

Forms hydrogen bonds with water (molecules) ✓

DO NOT ALLOW 'forms hydrogen bonds'

[2]

3. $\text{CH}_3\text{COOCH}_2\text{CH}_2\text{OOCCH}_3$

1 mark for each ester end of molecule ✓✓


ALLOW displayed formula OR skeletal formula

ALLOW sticks


$\text{CH}_3\text{COOCH}_2\text{CH}_2\text{OH}$ shows one of the two ester groups and scores one mark

[2]

4. (i)



DO NOT ALLOW



2

- (ii) E/Z ✓

ALLOW cis-trans


IGNORE geometric

1

(iii) $\text{CH}_3\text{CH}_2\text{CH}=\text{CH}_2$ **OR** but-1-ene ✓

*If but-1-ene given in part (i),
ALLOW but-2-ene OR $\text{CH}_3\text{CH}=\text{CHCH}_3$
i.e. ECF from (i)*

DO NOT ALLOW methylpropene:



1

[4]

5. From the evidence, candidates may have identified compound **F** as propanone, propanal or propanoic acid

If **F** is propanone or propanoic acid, then maximum score = 7; **but** if **F** is propanal then maximum score = 6

The mark scheme for **F** = propanone and propanal is shown below.

mass spec of E – Remember to check the spectrum

Quality of Written Communication – mass spec gives M^+ or molecular ion of 60 **OR** mass spec gives parent ion of 60 **OR** highest m/z (**ALLOW** m/e) value is 60 ✓

$m/z = 45$ indicates loss of CH_3

OR $m/z = 45$ indicates presence of CH_3CHOH

OR $\text{CH}_2\text{CH}_2\text{OH}$ **OR** $\text{C}_2\text{H}_5\text{O}$ ✓

IR of F – Remember to check the spectrum

IR shows no broad absorption between 2500 to 3300 cm^{-1} so no O—H bond

OR no broad absorption between 2500 to 3300 cm^{-1} so not a carboxylic acid ✓

IR shows absorption at 1700 cm^{-1} due to a C=O bond

OR absorption at 1700 cm^{-1} indicates a ketone **OR** aldehyde present

Identification and equation

F is CH_3COCH_3 **OR** propanone ✓

E is $\text{CH}_3\text{CHOHCH}_3$ **OR** propan-2-ol ✓

$\text{CH}_3\text{CHOHCH}_3 + [\text{O}] \rightarrow \text{CH}_3\text{COCH}_3 + \text{H}_2\text{O}$ ✓

If **F** has been incorrectly identified as propanal, mark identification and equation as ECF, so max = 2

ALLOW **E** is $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$ ✓

ALLOW: $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH} + [\text{O}] \rightarrow \text{CH}_3\text{CH}_2\text{CHO} + \text{H}_2\text{O}$ ✓

The mark scheme for F = propanoic acid is shown below.

mass spec of E– Remember to check the spectrum

QWC – mass spec gives M^+ or molecular ion of 60

OR mass spec gives parent ion of 60

OR highest m/z (OR m/e) value is 60 ✓

$m/z = 45$ indicates loss of CH_3

OR $m/z = 45$ indicates presence of CH_3CHOH

OR CH_2CH_2OH OR C_2H_5O ✓

IR of F– Remember to check the spectrum

IR shows (broad) absorption somewhere between 3500 and 2500 cm^{-1} suggests carboxylic acid

OR O–H bond ✓

IR shows absorption at 1700 cm^{-1} due to C=O

OR absorption at 1700 cm^{-1} indicates a carboxylic acid ✓

Identification and equation

F is CH_3CH_2COOH OR propanoic acid ✓

E is $CH_3CH_2CH_2OH$ OR propan-1-ol ✓

$CH_3CH_2CH_2OH + 2[O] \rightarrow CH_3CH_2COOH + H_2O$ ✓

Extra guidance for marking of question

If E has **not** been identified OR if F has been identified as a **ketone or aldehyde**, use the **first** mark scheme

If F has been identified as a **carboxylic acid**, use the **second** mark scheme

Mass spec

These two marking points stand as **independent** marks whichever compounds have been identified.

The positive sign for fragment ions is not required. **IGNORE** negative charge.

The mass spec may well be on the actual spectrum.

IR mark

These stand as **independent** marks whichever compounds have been identified.

The IR analysis may well be on the actual spectrum.

Identification marks

If both structure and name are given they must **both** be correct but allow 'propanol' drawn with the correct structure because the position number of the –OH has been clearly identified

ALLOW ECF for identification of **F** e.g. if **E** is pentan-2-ol ✗ then an answer of pentan-2-one for **F** will be given a mark ✓ as ECF

ALLOW identification marks for **E** and **F** from equation

Equation mark

ALLOW ECF for any correct equation showing the oxidation of **any** alcohol to the appropriate product.

ALLOW molecular formulae in equations,

i.e. $C_3H_7OH + [O] \rightarrow C_2H_5CHO + H_2O$ ✓ ; $C_3H_8O + [O] \rightarrow C_3H_6O + H_2O$ ✓ ;

$C_3H_7OH + [O] \rightarrow C_2H_5COH + H_2O$ ✓

[7]

6. (a) **method 1:**
fermentation of sugars or carbohydrates **OR** reaction with yeast with sugar or carbohydrates ✓
 $C_6H_{12}O_6 \rightarrow 2C_2H_5OH + 2CO_2$ ✓

method 2:
hydration of ethene **OR** reaction of ethene with water **OR** reaction of steam with ethene ✓

$C_2H_4 + H_2O \rightarrow C_2H_5OH$ ✓

ALLOW sugar from equation

ALLOW C_2H_6O in equation

ALLOW correct multiples

IGNORE state symbols

ALLOW ethene from the equation

IGNORE mention of any catalyst



*ALLOW C_2H_6O in equation **OR** H_2O over the arrow*

ALLOW correct multiples

IGNORE state symbols

4

(b) (i) $(\text{CH}_3)_2\text{CO}$ **OR**




If name and formula given both need to be correct

ALLOW propanone **OR** acetone

IGNORE propone

NOT incorrect named compound



ALLOW O instead of [O]

ALLOW correct multiples

IGNORE state symbols

2

(ii) $\text{CH}_3\text{CH}_2\text{COOH}$ **OR** propanoic acid ✓

Any number or range of numbers between 1750–1640 (cm^{-1})
for C=O ✓

Any number or range of numbers between 2500–3300 (cm^{-1})
for O–H ✓

ALLOW C=O and O–H marks independent of compound
identified **i.e. stand alone marks**

ALLOW correct bonds shown by the appropriate absorption on
the IR spectrum

IGNORE reference to C–O bond

3

(c) (i) 2-methylpropan-2-ol ✓


ALLOW methylpropan-2-ol **OR** tertiarybutanol

1

(ii) ester ✓

1

(iii) $\text{CH}_3\text{CO}_2\text{C}(\text{CH}_3)_3$ **OR** $\text{CH}_3\text{COOC}(\text{CH}_3)_3$
OR



ester group shown ✓

rest of molecule ✓

ALLOW skeletal formula OR displayed formula

ALLOW ester linkage even if rest of structure is wrong

2

[13]

7. Availability of starting materials:

availability

sugar is renewable because it can be grown (1)

ethane is finite because it is obtained by processing of crude oil (1)

energy:

fermentation: energy is required for distillation/

hydration: energy is required to generate steam (1)

atom economy and waste products:

atom economy for fermentation < atom economy hydration (1)

In fermentation, CO_2 is produced in addition to ethanol/ethanol is

not the only product (1)

In hydration, ethanol is the only product/hydration is an addition reaction (1)

Atom economy of fermentation could be increased by finding a use CO_2 (1)



Atom economy linked to a chemical equation to show that hydration has 100% atom economy/fermentation has 51% atom economy (1) 7max

[7]

8. (a) (i) (volatile components) can escape/distil out (1)
ethanal is most volatile/bpt less than 60 °C/partial oxidation (1) 2
- (ii) (volatile components) cannot escape/ refluxed (1)
complete oxidation will be achieved/oxidised to the acid (1) 2
- (b) $C_2H_5OH + 2[O] \rightarrow CH_3COOH + H_2O$
 C_2H_5OH , $2[O]$ and CH_3COOH (1)
rest of equation (1) 2

[6]


9. (i) $C_6H_{12}O_6(aq) \rightarrow 2C_2H_5OH(l)$ or (aq) + $2CO_2(g)$ balanced equation 1
state symbols can be awarded only if equation shows $C_6H_{12}O_6$,
 C_2H_5OH and CO_2 1
- (ii) anaerobic, aqueous, temp range 25 – 40°C/warm to just above room temp 2
- (iii) no more bubbles/gas/ CO_2 1

[5]

10. $CH_3CH(OH)CH_3 + 4\frac{1}{2}O_2 \rightarrow 3CO_2 + 4H_2O / C_3H_8O$
(1 mark if correct formula for all four chemicals and 1 mark for
correct balancing)

[2]


11. (i) 2



- (ii) either (2-)methylpropan-1-ol or (2-)methylpropan-2-ol 1

[3]

12.




Minimum – must display/show C=C

[3]

13. (a) (i) H⁺ 1
 Cr₂O₇²⁻ 1
 (ii) Orange to green/black/blue 1
- (b) (i) contains a C=O/aldehyde, ketone, carboxylic acid and ester/
 carbonyl/carbonyl in an aldehyde 1
 (ii) does **not** contain a O–H/ (hydrogen bonded in a) carboxylic acid 1
 (iii) distillation (no mark) **because** distillation allows loss of volatile
 components /removes butanal from oxidising mixture 1
 prevents formation of RCOOH/ partial oxidation would be achieved 1
 or reverse argument for reflux not being used
 in that reflux prevents loss of volatile components
 hence complete oxidation would be achieved/RCOOH would be formed
 ✓

[7]



14. (a) (i) 1



- (ii) H₂SO₄/Al₂O₃/(hot) pumice/H₃PO₄ 1
 (H₂SO₄(aq) or dil H₂SO₄ loses the mark)

(iii)

1



(b) (i)

1




(ii)

2

from the diol allow

from the Cl-alcohol allow




[6]

15. (i) *low volatility*, = **high** boiling point/ not easy to vapourise/owtte 1
intermolecular bonds. = bonds/forces/attractions **between** molecules 1

(ii) type of intermolecular bond = hydrogen bond 1

dipoles on both O-H bonds 1

H-bond shown as a 'dashed bond' 1




(iii) (The boiling point of glycerol will be higher than ethanol because there are) 1
 more OH groups \therefore more H-bonds 1

[6]

16. (i) butan-2-ol by name or by formula ✓ 1

(ii)



curly arrow from the O of the OH⁻ to C^(δ+) ✓

curly arrow from C-Cl bond to Cl **and** correct dipoles ✓

correct products/ allow NaCl ✓

curly arrow from lone pair on :OH⁻ ✓

S_N1 route can still score all 4 marks:

curly arrow from C-Cl bond to Cl **and** correct dipoles ✓

curly arrow from the O of the OH⁻ to C⁺ ion ✓

correct products/ allow NaCl ✓


curly arrow from lone pair on :OH⁻ ✓

4

[5]

17. (i) H⁺ ✓ Cr₂O₇²⁻ 2

(ii)



(iii) carboxylic acid would have an absorption between $1680 - 1750 \text{ cm}^{-1}$ / 1700 cm^{-1} or $2500 - 3300 \text{ cm}^{-1}$.

1

[6]

18. (a) (i) H_2SO_4 – any mention of (aq) loses the mark

1

(ii) any correct formula/structure or name for benzoic acid




1

(b) (i) dichromate/ $\text{Cr}_2\text{O}_7^{2-}$ /permanganate

1

(ii)

1




[4]

19. $\text{C}_6\text{H}_{12}\text{O}_6 \rightarrow 2\text{C}_2\text{H}_5\text{OH} + 2\text{CO}_2$
($\text{C}_2\text{H}_5\text{OH}$ & CO_2 ✓)

[2]

20.




- dipoles 1
- hydrogen bond between O in one O-H
and H in the other O-H 1
- lone pair from O involved in the H-bond 1

[3]

21. (a) (i) (volatile components) can escape/distil out 1
- ethanal is most volatile/b pt less than 60°C/partial oxidation 1
- (ii) (volatile components) cannot escape/ refluxed 1
- complete oxidation will be achieved/oxidised to the acid 1
- (b) $C_2H_5OH + 2[O] \rightarrow CH_3COOH + H_2O$
($CH_3COOH + H_2O$ ✓) 2
- (c) spectrum C 1
- spectrum C only shows absorption at 1700 cm^{-1} for the C=O 1
- the other two spectra contain the OH group absorption at approx 3000 cm^{-1} 1

[9]

22. (a) (i) prop-2-en-1-ol $CH_2=CHCH_2OH$ must show the C=C double bond 1
- acrolein**




must clearly show the aldehyde group and the C=C

- (ii) alkene/C=C double bond 1


- (b) (i) acidified /H⁺ 1
dichromate/Cr₂O₇²⁻ 1
(ii) CH₂CHCH₂OH/ C₃H₆O/ C₃H₅OH + [O] → CH₂CHCHO/ C₃H₄O/
C₂H₃CHO + H₂O 1
not CH₂CHCOH

[6]

23. (i) CH₂CHCH₂OOCCHCH₂ / (C₆H₈O₂) 1
H₂O 1
(ii) 2






or



1 mark if the ester group, 1 mark for the rest of the molecule.
COO/CO₂ without displaying the ester, they can still get 1 mark.

[4]

24. (a) ✓✓✓




3

- (b) (i) orange to green/dark green/brown/black ✓ 1
(ii) C₄H₉OH/ C₄H₁₀O + 2[O] → C₃H₇COOH + H₂O ✓✓ 2
1 mark available for correct formula of the carboxylic acid
(iii) Identify isomer 2-methylpropan-1-ol by appropriate number/name/formula ✓ 1


- (c) (i) CH_2 has mass = 14, $14 \times 4 = 56$ ✓ 1
 $\therefore \text{C}_4\text{H}_8$ ✓ 1
(ii) $\text{C}_4\text{H}_9\text{OH} \rightarrow \text{C}_4\text{H}_8 + \text{H}_2\text{O}$ ✓ 1
(iii) Identify butan-2-ol by appropriate number/name/formula 1
- (d) (i) H_2SO_4 ✓ 1
(ii) 0.06 ✓ 1
(iii) 60% ✓ 1

[14]

25. (a) (i) alkene ✓ 1
alcohol/hydroxy/hydroxyl ✓ 1
- (b) (i) I = alkene & II = alcohol... both are needed ✓ 1
(ii) decolourised / colourless ✓ 1
(iii) ✓ 1



- (iv) X as shown below ✓ 1



- (c) (i) Ni/Pt/Rh/Pd ✓ 1
(ii) compound B is $\text{C}_{10}\text{H}_{22}\text{O}$ ✓ 1
(iii) $\text{C}_{10}\text{H}_{20}\text{O} + \text{H}_2 \rightarrow \text{C}_{10}\text{H}_{22}\text{O}$ ✓ 1

[9]

26. (a) $\text{C}_2\text{H}_5\text{OH} + 3\text{O}_2 \rightarrow 2\text{CO}_2 + 3\text{H}_2\text{O}$ ✓✓ 2

2CO₂ + 3H₂O gets 1 mark

- (b) **Fermentation** 1
- C₆H₁₂O₆ → 2C₂H₅OH + 2CO₂ ✓ 1
- Yeast /enzyme / temperature about 30 °C/ batch process ✓ 1
- Hydration** of ethene. ✓ 1
- C₂H₄ + H₂O → C₂H₅OH ✓ 1
- Temp > 100 °C/Press 370 – 100 atm / 6 –20 MPa/phosphoric acid catalyst/
continuous process ✓ 1
- Glucose is obtained from plants ✓ 1
- Ethene is obtained from crude oil/cracking/fossil fuel ✓ 1
- glucose is renewable/ethene isn't ✓ 1




1 mark available for *Quality of written communication*..... base the award of the mark on the ability to communicate the essential chemistry by correct use of at least two from:

fermentation/hydration/catalyst/renewable/sustainable/biofuel/
enzymes/finite/cracking ✓ 1

[12]

27. (a) (i) C₄H₁₀ ✓ 1
- (ii) C₂H₅O ✓ 1
- (iii) B and E ✓ 1
- (iv) A and F ✓ 1
- (b) (C₄H₉OH →) C₄H₈ + H₂O ✓ 1

(c) any unambiguous formula: ✓ 1



buta-1,3-diene ✓

name ecf to the structure only if structure above has formula C₄H₆

1

[7]

28. (a) (i) Alkene/C=C ✓ 1

Alcohol/ROH/hydroxy/hydroxyl/OH (not OH⁻ or hydroxide) ✓ 1

(ii) One of the C in both C=C is joined to two atoms or groups that are the same ✓ 1

(b) Observation decolourisation (of Br₂) ✓ 1

Molecular formula C₁₀H₁₈OBr₄ ✓✓ 2

C₁₀H₁₈OBr₂ gets 1 mark

(c) reagent CH₃COOH ✓ 1

catalyst H₂SO₄/H⁺/HCl (aq) *or dilute loses the mark* ✓ 1

(d) (i) C₁₀H₁₈O + 2[O] → C₁₀H₁₆O₂ + H₂O ✓✓ 2

1 mark for H₂O and 1 mark for 2[O]

(ii) The infra-red spectrum was of compound Y because absorption between 1680 – 1750 cm⁻¹ indicates a C=O ✓ 1

and the absence of a peak between 2500 – 3300 cm⁻¹ shows the absence of the OH hydrogen bonded in a carboxylic acid ✓ 1

[12]