M1.

Mark Range	The marking scheme for this part of the question includes an overall assessment for the Quality of Written Communication (QWC). There are no discrete marks for the assessment of QWC but the candidates' QWC in this answer will be one of the criteria used to assign a level and award the marks for this part of the question			
	Descriptor an answer will be expected to meet most of the criteria in the level descriptor			
4-5	 claims supported by an appropriate range of evidence 			
	 good use of information or ideas about chemistry, going beyond those given in the question 			
	 argument well structured with minimal repetition or irrelevant points 			
	 accurate and clear expression of ideas with only minor errors of grammar, punctuation and spelling 			
2-3	 claims partially supported by evidence 			
	 good use of information or ideas about chemistry given in the question but limited beyond this 			
	the argument shows some attempt at structure			
	 the ideas are expressed with reasonable clarity but with a few errors of grammar, punctuation and spelling 			
0-1	 valid points but not clearly linked to an argument structure 			
	 limited use of information or ideas about chemistry 			
	unstructured			
	errors in spelling, punctuation and grammar or lack of fluency			

(a) (i) M_r of $C_6H_5NH_2 = 93$ M_r of $CH_3COCI = 78.5$ total M_r of reagents = 264.5

% atom economy = $\frac{M_{r} \text{ of wanted product}}{\text{total M}_{r} \text{ of all reagents}} \times 100 \text{ QWC}$

1

1

$$= \frac{135}{264.5} \times 100 = 51.0 \%$$

1

(ii) expected yield =
$$\frac{10}{93} \times 0.5 \times 135 = 7.26 \text{ kg}$$

1

% yield =
$$\frac{5.38}{7.26} \times 100 = 74.1 \%$$

1

(iii) Although yield appears satisfactory (74%) % atom economy is only 51% QWC


1

nearly half of the material produced is waste and must be disposed of QWC

1

(b) (nucleophilic) addition-elimination


1



4

(c) $HNO_3 + 2H_2SO_4 \rightarrow NO_2^+ + H_3O^+ + 2HSO_4^-$

1



3

[16]

M2.D

[1]

M3.B

[1]

M4. (a) (nucleophilic) addition-elimination;

$$(CH_3CH_2 \xrightarrow{M_2} C)$$

$$M_1 C \longrightarrow CH_3CH_2 \xrightarrow{C} C$$

$$H C \xrightarrow{N+} H$$

$$(CH_3 \xrightarrow{N+} NH_2)$$

$$Allow C_2H_5 \xrightarrow{C} C$$

$$CH_3CH_2 \xrightarrow{N+} C$$

$$H C \xrightarrow{N+} H$$

(M3 for structure) (M4 for 3 arrows and lone pair) (M2 not allowed independent of M1, but allow M1 for correct attack on C+ if M2 show as independent first.) (+on C of C=O loses M2 but ignore δ + if correct) (Cl- removing Ft loses M4)


1

(If MS lost above for wrong C chain, do not penalise same error again here)

5

(b) $CH_3CH_2COCI + AICI_3 \rightarrow [CH_3CH_2CO]^+ + AICI_4^-;$

(penalise wrong alkyl group once at first error)
(position of + on electrophile can be on O or C or outside [])
(penalise wrong curly arrow in the equation or lone pair on



3

$$A|C|_4^- + H^+ \rightarrow A|C|_3^- + HC|_7^-$$

(or can be gained in mechanism);

1

(c) M1 CH₃CH₂COCI + H₂O → CH₃CH₂COOH + HCI 1 (penalise wrong alkyl group once at first error)

1

M2 M_r of CH₃CH₂COCI = 92.5 1 (if Mr wrong, penalise M2 only)

1

M3 moles of $CH_3CH_2COCI = 1.48/92.5 = 0.016 1$

1

M4 moles NaOH = $2 \times 0.016 = 0.0321$ (allow for $\times 2$ conseq to wrong no of moles)

1

1

M5 volume of NaOH = 0.032/0.42 = 0.0762 dm³ or 76.2 cm³ 1 (with correct units) (if ×2 missed in M4 lose M5 also)

[16]

M5. (a)
$$CH_3COCI + AICI_3 \rightarrow CH_3^{\stackrel{+}{C}}O + AICI_4^{\stackrel{-}{4}}$$
 (1) equation (1)

penalise wrong alkyl group once at first error position of + on electrophile can be on O or C or outside [] penalise wrong curly arrow in the equation or lone pair on AlCl₃ else ignore

2


1

3

1

Electrophilic substitution

NOT F/C acylation



horseshoe must not extend beyond C2 to C6 but can be smaller

+ not too close to C1


M3 arrow into hexagon unless Kekule allow M3 arrow independent of M2 structure

M1 arrow from within hexagon to C or to + on C

+ must be on C of RCO

(b) Nucleophilic addition

NOT reduction



M2 not allowed independent, but can allow M1 for attack of

H on C+ formed

	Do not anow ii	on oxides	1	[14]
	allow dilute an Do not allow in			
	(conc) H ₂ SO ₄ or (conc) H ₃ PO ₄			
(c)	dehydration or elimination	1	1	
	<u>1</u> –phenylethan(–1–)ol	or (1-hydroxyethyl)benzene	1	
			4	