

Q1.A green solution, **X**, is thought to contain $[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$ ions.

- (a) The presence of these ions can be confirmed by reacting separate samples of solution **X** with aqueous ammonia and with aqueous sodium carbonate.

Write equations for each of these reactions and describe what you would observe.

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

(4)

- (b) A 50.0 cm^3 sample of solution **X** was added to 50 cm^3 of dilute sulfuric acid and made up to 250 cm^3 of solution in a volumetric flask.

A 25.0 cm^3 sample of this solution from the volumetric flask was titrated with a $0.0205 \text{ mol dm}^{-3}$ solution of KMnO_4
At the end point of the reaction, the volume of KMnO_4 solution added was 18.70 cm^3 .

- (i) State the colour change that occurs at the end point of this titration and give a reason for the colour change.

.....
.....
.....
.....

(2)

- (ii) Write an equation for the reaction between iron(II) ions and manganate(VII)

ions.

Use this equation and the information given to calculate the concentration of iron(II) ions in the original solution **X**.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(5)
(Total 11 marks)

Q2. Which of these pieces of apparatus has the lowest percentage uncertainty in the measurement shown?

- A** Volume of 25 cm³ measured with a burette with an uncertainty of ± 0.1 cm³.
- B** Volume of 25 cm³ measured with a measuring cylinder with an uncertainty of ± 0.5 cm³.
- C** Mass of 0.150 g measured with a balance with an uncertainty of ± 0.001 g.
- D** Temperature change of 23.2 °C measured with a thermometer with an uncertainty of ± 0.1 °C.


(Total 1 mark)

Q3. N-phenylethanamide is used as an inhibitor in hydrogen peroxide decomposition and also in

the production of dyes.

N-phenylethanamide can be produced in a laboratory by the reaction between phenylammonium sulfate and an excess of ethanoic anhydride:

- (a) A student carried out this preparation using 1.15 g of phenylammonium sulfate ($M_r = 284.1$) and excess ethanoic anhydride.



- (i) Calculate the maximum theoretical yield of N-phenylethanamide that could be produced in the reaction. Record your answer to an appropriate precision.

Show your working.

.....
.....
.....
.....
.....
.....
.....

(3)

- (ii) In the preparation, the student produced 0.89 g of N-phenylethanamide. Calculate the percentage yield for the reaction.

.....
.....
.....
.....

(1)

- (b) The student purified the crude solid product, N-phenylethanamide, by recrystallisation.

- (i) Outline the method that the student should use for this recrystallisation.

.....
.....

.....
.....
.....
.....
.....
.....
.....

(4)

- (ii) Outline how you would carry out a simple laboratory process to show that the recrystallised product is a pure sample of N-phenylethanamide.

.....
.....
.....
.....
.....
.....
.....
.....

(3)

- (iii) Assume that the reaction goes to completion.

Suggest **two** practical reasons why the percentage yield for this reaction may **not** be 100%.

1

.....

2

.....

(2)

- (c) The reaction to form N-phenylethanamide would happen much more quickly if the student used ethanoyl chloride instead of ethanoic anhydride.

Explain why the student might prefer to use ethanoic anhydride, even though it has a slower rate of reaction.

.....
.....
.....

(2)
(Total 15 marks)

Q4. The maximum errors for the pipette and the burette are shown below. These errors take into account multiple measurements.

Pipette $\pm 0.05 \text{ cm}^3$
Burette $\pm 0.15 \text{ cm}^3$

Estimate the maximum percentage error in using each of these pieces of apparatus.

Use an average titre 24.25 cm^3 to calculate the percentage error in using the burette.

Show your working.

Pipette

.....

Burette

.....

(Total 2 marks)