

M1.B

[1]

M2.A

[1]

M3.(a) (i) $-\log[\text{H}^+]$

Penalise missing [] here and not elsewhere

1

(ii) $[\text{H}^+][\text{OH}^-]$

1

(b) (i) $[\text{H}^+] = 2.34 \times 10^{-7}$

1

pH = 6.63

Penalise fewer than 3 sig figs but allow more than 2 dp

1

(ii) $[\text{H}^+] = [\text{OH}^-]$

1

(iii) **M1** $[\text{H}^+] = K_w/[\text{OH}^-]$

if upside down or CE, allow M3 only for correct use of their $[\text{H}^+]$

1

M2 $(= 5.48 \times 10^{-14}/0.140) = 3.91 \times 10^{-13}$

1

M3 pH = 12.4(1)

not 12.40 (AE from 12.407)

1

Penalise fewer than 3 sig figs but allow more than 3 sfs

For values above 10, allow 3sfs - do not insist on 2 dp.

For values below 1, allow 2dp – do not insist on 3 sig figs

Not allow pH = 14 – pOH but can award M3 only for pH = 13.1(46)

Can award all three marks if $pK_w = 13.26$ is used

- (c) **M1** mol NaOH = mol OH⁻ = $(30 \times 10^{-3}) \times 0.20 = 6.0 \times 10^{-3}$
mark for answer 1
- M2** mol H₂SO₄ = $(25 \times 10^{-3}) \times 0.15 = 3.75 \times 10^{-3}$
mark for answer 1
- M3** mol H⁺ = $(25 \times 10^{-3}) \times 0.15 \times 2 = 7.5 \times 10^{-3}$
 OR XS mol H₂SO₄ = 0.75×10^{-3}
if factor of 2 missed or used wrongly, CE - lose M3 and next mark gained. In this case they must then use K_w to score any more.
see examples below 1
- M4** XS mol H⁺ = 1.5×10^{-3} 1
- M5** [H⁺] = $(1.5 \times 10^{-3}) \times (1000/55) = 0.0273$
if no use or wrong use of volume, lose M5 and M6 except if 1000 missed
AE -1 (pH = 4.56) 1
- M6** pH = 1.56
Penalise fewer than 3 sig figs but allow more than 3 sfs
For values above 10, allow 3sfs - do not insist on 2 dp.
For values below 1, allow 2dp – do not insist on 3 sig figs 1

[14]

M4.C

[1]

- M5.(a)** pH on the y-axis, volume of alkali on the x-axis
If axes unlabelled use data to decide that pH is on y-axis.

1

Uses sensible scales

*Lose this mark if plotted paths do not cover **half** of the paper.*

Lose this mark if the graph plot goes off the squared paper.

1

Labels the axes

Allow mark for axes labelled 'pH' and 'volume'.

1

Plots all of the points correctly

1

Line through the points is smooth and has the correct profile

Ignore 0–5 cm³ section of the graph.

Lose this mark if graph is kinked or not a single line.

1

Line ignores the point at 12 cm³

Lose this mark if point clearly not treated as an anomaly.

1

(b) (i) 24.4 cm³ ± 0.2

If no answer in (i) allow answer written on the graph.

*Allow this answer **only**.*

Do not penalise precision.

1

(ii) 12.2 cm³ ± 0.1

If no answer in (ii), allow answer written on the graph.

Allow answer to (i) divided by 2.

Do not penalise precision.

1

(iii) 3.9 ± 0.2

If no answer in (iii), allow answer written on the graph.

Consequential marking from (ii)

Lose this mark if answer not given to 1 dp.

1

- (c) $pK_a = -\log K_a$ or $K_a = 10^x$, where $x = -$ (answer to b(iii))

1

$$1.26 \times 10^{-4}$$

3.7 to 4.1 gives $K_a = 7.9 \times 10^{-5}$ to 2.0×10^{-4}

Consequential marking from b(i).

Correct answer without working scores 1 mark only.

Do not penalise precision.

1

- (d) Methanoic acid

Consequential marking from (c).

$pK_a = 3.7$ gives methanoic acid.

$pK_a = 4.1$ gives ethanoic acid.

No lucky guesses – candidates must apply answer from (c).

Do not allow answers based on data given in (f).

1

- (e) Error in using pipette is 0.2% **and**
Error in using burette is $0.15 \times 100 /$ (answer to b(i))

Using 24.4 for burette gives 0.6%

Do not penalise precision.

Allow if errors are given without working.

Lose mark if the burette error is not calculated on b(i).

*If the error being calculated is **not** stated, allow if the calculations are in the same order as in the question (pipette, burette).*

1

- (f) Difference is $1.6 \times 10^{-4} - 1.26 \times 10^{-4} = 0.34 \times 10^{-4}$
Allow consequential answer from (c).
Do not penalise precision.

$$0.34 \times 100 / 1.6 \text{ is a } 21\% \text{ error}$$

Correct final answer without working scores 1 mark.

Using 1.9×10^{-4} gives 0.3×10^{-4} and 18.8%.

1

- (g) Calibrate meter **or** thermostat the mixture **or** maintain constant temperature
Do not allow 'repeat experiment'.

1

- (h) Mixture is a buffer

1

[16]

M6.C

[1]

- M7.** (a) (i) $-\log[H^+]$
or $\log 1/[H^+]$
penalise ()

1

- (ii) $[H^+] = 0.56$
mark for the answer; allow 2dp or more


1

$$[H_2SO_4] = \frac{1}{2} \times 0.56 = 0.28$$

1

- (b) (i) $CH_3COOH + NaOH \rightarrow CH_3COONa + H_2O$

OR



Allow CH_3CO_2H etc

1

- (ii) mol acid = $(25.0 \times 10^{-3}) \times 0.41 = 1.025 \times 10^{-2}$ or 1.03×10^{-2}

1

$$[NaOH] = 1.025 \times 10^{-2} / 22.6 \times 10^{-3} = 0.45(4)$$

mark for answer
if not 0.454 look back for error

1

OR

$$[\text{NaOH}] = 1.03 \times 10^{-2} / 22.6 \times 10^{-3} = 0.456 \text{ or } 0.46$$

(iii) cresol purple

1

(iv) NaOH reacts with carbon dioxide (in the air)

1

(c) (i)
$$K_a = \frac{[\text{H}^+][\text{CH}_3\text{COO}^-]}{[\text{CH}_3\text{COOH}]}$$

allow molecular formulae or minor slip in formulae

penalise ()

allow H_3O^+

not allow HA etc

1

(ii)
$$K_a = \frac{[\text{H}^+]^2}{[\text{CH}_3\text{COOH}]}$$
 or with numbers

1

allow HA etc here

This can be scored in part (c)(i) but doesn't score there.

$$[\text{H}^+] = (\sqrt{1.74 \times 10^{-5} \times 0.410}) = \sqrt{7.13 \times 10^{-6}} = 2.67 \times 10^{-3}$$

1

mark for 2.67×10^{-3} or 2.7×10^{-3} either gives 2.57

pH = 2.57 can give three ticks here for (c)(ii)
penalise decimal places < 2 >

1

pH mark conseq on their $[\text{H}^+]$

so 5.15 gets 2 marks where square root not taken

(iii) **M1** mol $\text{OH}^- = (10.0 \times 10^{-3}) \times 0.10 = 1.0 \times 10^{-3}$

*If no subtraction or other wrong chemistry the max score is 3
for M1, M2 and M4*

1

$$\mathbf{M2} \text{ orig mol HA} = (25.0 \times 10^{-3}) \times 0.41 = 0.01025$$

1

$$\text{or } 1.025 \times 10^{-2} \text{ or } 1.03 \times 10^{-2}$$

$$\mathbf{M3} \text{ mol HA in buffer} = \text{orig mol HA} - \text{mol OH}^-$$

1

$$= 0.00925 \text{ or } 0.0093$$

*If A⁻ is wrong, max 3 for M1, M2 and M3 or use of
pH = pKa - log [HA]/[A⁻]*

$$\mathbf{M4} \text{ mol A}^- \text{ in buffer} = \text{mol OH}^- = 1.0 \times 10^{-3}$$

*Mark is for insertion of correct numbers in correct expression
for [H⁺]*

1

$$\mathbf{M5} \text{ [H}^+] = \left(\frac{K_a \times [\text{CH}_3\text{COOH}]}{[\text{CH}_3\text{COO}^-]} \right)$$

1

$$\frac{(1.74 \times 10^{-5})(0.00925)}{0.0010} \text{ or } \frac{(1.74 \times 10^{-5})(0.00930)}{0.0010}$$

$$(\text{= } 1.61 \times 10^{-4} \text{ or } 1.62 \times 10^{-4})$$

$$\mathbf{M6} \text{ pH} = 3.79 \text{ can give six ticks for 3.79}$$

if [HA]/[A⁻] upside down lose M5 & M6

If wrong method e.g. [H⁺]/[HA] max 3 for M1, M2 and M3

Some may calculate concentrations

*[HA] = 0.264 and [A⁻] = 0.0286 and rounding this to 0.029
gives pH = 3.80 (which is OK)*

NB Unlike (c)(ii), this pH mark is NOT awarded conseq to their [H⁺] unless following AE

*BEWARE: using 0.01025 wrongly instead of 0.00925 gives
pH = 3.75*

(this gets 3 for M1, M2 & M4)

1

[18]

