

M1. (a) moles HA = $\frac{25}{10^3} \times 0.150 = 3.75 \times 10^{-3}$ (1)

$$\therefore \text{vol NaOH} = \frac{3.75 \times 10^{-3}}{0.20} = 1.875 \times 10^{-2} \text{ dm}^3 \text{ (1)}$$

or 18.75 cm³

2

(b) (i) pH = $-\log_{10} [\text{H}^+]$ (1)

(ii) Value above 7 but below 11 (1)

(iii) phenol red / thymol blue / phenolphthalein / thymolphthalein
i.e. indicator with $7 < pK_n < 11$

3

(c) (i) Only slightly dissociated/ionised (1)
NOT "not fully dissociated / ionised"

(ii) $K_a = \frac{[\text{H}^+][\text{A}^-]}{[\text{HA}]}$ (1)

NOT $\frac{[\text{H}^+]^2}{[\text{HA}]}$

(iii) For weak acid alone:

$$K_a = \frac{[\text{H}^+]^2}{[\text{HA}]} \text{ (1)}$$

$$\therefore [\text{H}^+] = \sqrt{(2.75 \times 10^{-6}) \times 0.15}$$

$$= 2.03 \times 10^{-3} \text{ (1)}$$

$$\therefore \text{pH} = 2.69 \text{ (1)}$$

*pH should be given to 2 decimal places
penalise answer to 1 d.p. once in question*

5

(d) moles OH⁻ added = 1.875×10^{-3} = moles A⁻ = moles HA left **(1)**

or $[A^-] = [HA]$

∴ $K_a = [H^+]$ or $pH = pK_a$ **(1)**

∴ $pH = 4.56$ **(1)**

3

[13]

M2. (a) before any KOH added: $K_a = \frac{[H^+][A^-]}{[HA]}$ or $\frac{[H^+][CH_3COO^-]}{[CH_3COOH]}$ **(1)**

∴ $K_a = \frac{[H^+]^2}{[CH_3COOH]}$ **(1)**

∴ $[H^+] = \sqrt{1.74 \times 10^{-5} \times 0.160} = 1.67 \times 10^{-3}$ **(1)**

∴ $pH = 2.78$ **(1)**

4

(b) at 8 cm³ KOH:

Moles KOH added = $(8 \times 10^{-3}) \times 0.210 = 1.68 \times 10^{-3}$ **(1)**

∴ moles of CH₃COO⁻ formed = 1.68×10^{-3} **(1)**

Original moles of CH₃COOH = $(25 \times 10^{-3}) \times 0.160 = 4.0 \times 10^{-3}$ **(1)**

∴ moles of CH₃COOH left = $(4.0 \times 10^{-3}) - (1.68 \times 10^{-3})$
 $= 2.32 \times 10^{-3}$ **(1)**

$[H^+] = K_a \times \frac{[CH_3COOH]}{[CH_3COO^-]}$ **(1)**

$= 1.74 \times 10^{-5} \times \frac{2.32 \times 10^{-3} / V}{1.68 \times 10^{-3} / V} = 2.40 \times 10^{-5}$ **(1)**

∴ $pH = 4.62$ **(1)**

It forget subtraction : max 5

If K_a expression not used max 5

- (c) at 40 cm³ of KOH:
 Total moles of KOH = (40 × 10⁻³) × 0.21 = 8.4 × 10⁻³ (1)
 ∴ excess moles of KOH = (8.4 × 10⁻³) - (4.0 × 10⁻³)
 = 4.4 × 10⁻³ (1)
 in total volume = 40 + 25 = 65 cm³ (1)

$$\therefore [\text{OH}^-] = 4.4 \times 10^{-3} \times \frac{1000}{65} = 0.0677 \text{ (1)}$$

$$\therefore [\text{H}^+] = \frac{10^{-14}}{0.0677}$$

OR pOH = 1.17

$$= 1.477 \times 10^{-13} \text{ (1)}$$

$$\therefore \text{pH} = 12.83 \text{ (1)}$$

If volume missed : max 4

If moles of acid wrong but method includes subtraction : max 5

If no subtraction : max 4

M3. (a) (i)
$$K_a = \frac{[\text{H}^+][\text{CH}_3\text{COO}^-]}{[\text{CH}_3\text{COOH}]} \text{ (1)}$$

or H₃O⁺

(ii) (1)
$$K_a = \frac{[\text{H}^+]^2}{[\text{CH}_3\text{COOH}]} \text{ (1)}$$

(2)
$$[\text{H}^+] = \sqrt{1.74 \times 10^{-5} \times 0.220} = 1.96 \times 10^{-3} \text{ (1)}$$

(3)
$$\text{pH} = -\log_{10}[\text{H}^+] \text{ (1)}$$

can score independently

- (4) pH = 2.71 (1)
 2 d.p. essential
 If forget $\sqrt{\quad}$ can score (1) and (3) for pH = 5.42

5

(b) (i) moles acid = $\frac{25}{1000} \times 0.220$ (1) = 5.50×10^{-3}
 $= \frac{x}{10^3} \times 0.150$
 $\therefore x = 25 \times \frac{0.220}{0.150}$ or $5.50 \times 10^{-3} \times \frac{1000}{0.150}$
 $= 36.7$ (or 37) cm³ (or 36.6) (1)

NOT 36 NOR 37.0 units must match

- (ii) *Indicator:* thymol blue (1)
Explanation: weak acid – strong base (1)
 equivalent at pH > 7 (1)
 or high pH

5

- (c) (1) mol NaOH added = $\frac{2.0}{40.0} = 0.050$ (1)
 If wrong M_r: CE \therefore lose marks (1) and (2) then mark on
 consequentially \rightarrow max 4
 (2) mol CH₃COOH left = $0.220 - 0.050 = 0.170$ (1)
 (3) mol CH₃COO⁻ formed = 0.050 (1)

(4) $[H^+] = K_a \frac{[\text{acid}]}{[\text{salt}]}$ OR $\text{pH} = \text{p}K_a + \log \frac{[A^-]}{[HA]}$ etc (1)
 If expression wrong no marks for 4 / 5 / 6

can score (1) to (4) in (5)

$$(5) \quad [H^+] = 1.74 \times 10^{-5} \times \frac{(0.170)}{(0.05)} \quad \text{OR} \quad \text{pH} = 4.76 + \log \left(\frac{0.05}{0.17} \right) \quad (1)$$

$$(6) \quad \text{pH} = 4.23 \quad (1)$$

Correct answer gets (1)(1)(1)(1)(1)

Mark (5) is for use of correct values of (acid moles) and (salt moles)

if one wrong allow pH conseq

if both wrong, no further marks

e.g. if candidate forgets substitution in (2)

he loses (2) and (5) but can score (1) (3) (4) (6) = max 4

for pH = 4.12 if $\frac{[\text{acid}]}{[\text{salt}]}$ upside down; answer 5.29 scores 3 for (1) (2) (3)

6

[16]