


**Q1.** The sketch below shows the change in pH when a  $0.200 \text{ mol dm}^{-3}$  solution of sodium hydroxide is added from a burette to  $25.0 \text{ cm}^3$  of a  $0.150 \text{ mol dm}^{-3}$  solution of the weak acid HA at  $25^\circ\text{C}$ .



(a) The volume of sodium hydroxide solution added at the equivalence point is  $x \text{ cm}^3$ . Calculate the value of  $x$ .

.....  
 .....  
 .....

(2)

(b) (i) Define the term pH.

.....

(ii) The pH at the equivalence point is  $c$ . Suggest a value for  $c$ .

.....

(iii) Identify a suitable indicator for detecting the equivalence point of the titration.

.....

(3)

(c) The value of  $K_a$  for the weak acid HA at 25 °C is  $2.75 \times 10^{-5} \text{ mol dm}^{-3}$ .

(i) Explain the term *weak* as applied to the acid HA.

.....

(ii) Write an expression for  $K_a$  for the acid HA.

.....

(iii) Calculate the pH of the  $0.150 \text{ mol dm}^{-3}$  solution of acid HA before any sodium hydroxide is added, i.e. the pH at point *a*.

.....  
.....  
.....  
.....  
.....

(5)

(d) Calculate the pH of the solution formed when  $\frac{x}{2} \text{ cm}^3$  of the  $0.200 \text{ mol dm}^{-3}$  solution of sodium hydroxide are added to  $25.0 \text{ cm}^3$  of the  $0.150 \text{ mol dm}^{-3}$  solution of HA, i.e. the pH at point *b*.

.....  
.....  
.....

(3)

(Total 13 marks)

**Q2.** A  $0.210 \text{ mol dm}^{-3}$  solution of potassium hydroxide was added from a burette to  $25.0 \text{ cm}^3$  of a  $0.160 \text{ mol dm}^{-3}$  solution of ethanoic acid in a conical flask. Given that the value of the acid dissociation constant,  $K_a$ , for ethanoic acid is  $1.74 \times 10^{-5} \text{ mol dm}^{-3}$ , calculate the pH at  $25 \text{ }^\circ\text{C}$  of the solution in the conical flask at the following three points:

before any potassium hydroxide had been added;

after  $8.0 \text{ cm}^3$  of potassium hydroxide solution had been added;

after  $40.0 \text{ cm}^3$  of potassium hydroxide solution had been added.

**(Total 16 marks)**

**Q3.** The value of the acid dissociation constant,  $K_a$ , for ethanoic acid is  $1.74 \times 10^{-5} \text{ mol dm}^{-3}$  at  $298 \text{ K}$ .

(a) (i) Write an expression for  $K_a$  for ethanoic acid.

.....

(ii) Calculate the pH at  $298 \text{ K}$  of a  $0.220 \text{ mol dm}^{-3}$  solution of ethanoic acid.

.....  
.....  
.....  
.....  
.....  
.....

**(5)**

(b) A sample of the  $0.220 \text{ mol dm}^{-3}$  solution of ethanoic acid was titrated against sodium hydroxide solution.

(i) Calculate the volume of a  $0.150 \text{ mol dm}^{-3}$  solution of sodium hydroxide required to neutralise  $25.0 \text{ cm}^3$  of the ethanoic acid solution.

.....  
.....  
.....  
.....

(ii) From the list below, select the best indicator for this titration and explain your choice.

<b>Name of indicator</b>	<b>pH range</b>
bromophenol blue	3.0 – 4.6
methyl red	4.2 – 6.3
bromothymol blue	6.0 – 7.6
thymol blue	8.0 – 9.6

*Indicator* .....

*Explanation* .....

.....  
.....

(5)

(c) A buffer solution is formed when  $2.00 \text{ g}$  of sodium hydroxide are added to  $1.00 \text{ dm}^3$  of a  $0.220 \text{ mol dm}^{-3}$  solution of ethanoic acid.

Calculate the pH at  $298 \text{ K}$  of this buffer solution.

.....  
.....  
.....  
.....  
.....  
.....  
.....

(6)  
(Total 16 marks)