


- 1 The Ostwald Process is a method for making nitric acid. The equation for the first stage of this process is



- (a) The equilibrium yield of nitrogen monoxide, NO, is **increased** by

(1)

- A increasing both the pressure and the temperature.
- B decreasing both the pressure and the temperature.
- C decreasing the pressure and increasing the temperature.
- D increasing the pressure and decreasing the temperature.

- (b) For this stage of the process, the catalyst is an alloy of platinum and rhodium. A pressure of between 4 and 10 atm and a temperature of 1150 K are used. Unreacted reactants are recycled.


Which one of the following changes will affect the value of the equilibrium constant,  $K_p$ ?

(1)

- A Changing the composition of the platinum-rhodium catalyst.
- B Increasing the pressure above 10 atm.
- C Decreasing the temperature below 1150 K.
- D Not recycling unreacted reactants.

**(Total for Question = 2 marks)**


2 Iodine is soluble in both water and hexane. If iodine is added to a mixture of the two solvents, then the following equilibrium is set up.




The equilibrium constant, known as the partition coefficient, is 85.

The density of hexane is  $0.66 \text{ g cm}^{-3}$ . The density of water is  $1.00 \text{ g cm}^{-3}$ .


Which of the following diagrams is correct for this system at equilibrium?




A



B




C



D

(Total for Question = 1 mark)


- 3 What are the units of the equilibrium constant ( $K_c$ ) for the hypothetical reaction below?



- A  $\text{mol}^2 \text{dm}^{-9}$
- B  $\text{mol}^{-2} \text{dm}^9$
- C  $\text{mol}^2 \text{dm}^{-6}$
- D  $\text{mol}^{-2} \text{dm}^6$

**(Total for Question = 1 mark)**

- 4 This question is about the reversible reaction below.



- (a) A chemist investigating this reaction started with 10 moles of  $\text{NO}_2$  and allowed the system to reach equilibrium. If 3 moles of  $\text{N}_2\text{O}_4$  are formed, the number of moles of  $\text{NO}_2$  at equilibrium is

(1)

- A 8.5
- B 7
- C 6
- D 4


- (b) Under different conditions, 40% of the moles present at equilibrium is  $\text{N}_2\text{O}_4$ . If the total pressure of the system is 2.0 atm, the numerical value of the equilibrium constant,  $K_p$  is

(1)

- A 0.56
- B 0.67
- C 1.5
- D 1.8

**(Total for Question = 2 marks)**

5 Ammonium chloride decomposes on heating:



The equilibrium constant,  $K_p$ , for this reaction equals

**A**  $P_{\text{NH}_3} \times P_{\text{HCl}}$


**B**  $\frac{1}{P_{\text{NH}_3} \times P_{\text{HCl}}}$

**C**  $\frac{P_{\text{NH}_3} \times P_{\text{HCl}}}{P_{\text{NH}_4\text{Cl}}}$

**D**  $\frac{P_{\text{NH}_4\text{Cl}}}{P_{\text{NH}_3} \times P_{\text{HCl}}}$

**(Total for Question = 1 mark)**

6 Consider the equilibrium below.



(a) An increase in pressure by a factor of 2 will

**(1)**

- A** quadruple  $K_p$ .
- B** double  $K_p$ .
- C** have no effect on  $K_p$ .
- D** halve  $K_p$ .


(b) The units of  $K_p$  are

**(1)**

- A**  $\text{atm}^{-2}$
- B**  $\text{atm}^{-1}$
- C**  $\text{atm}$
- D**  $\text{atm}^2$

**(Total for Question 2 marks)**

7 Methanol is produced in the equilibrium reaction



Addition of more hydrogen to the equilibrium mixture at constant temperature

- A increases the equilibrium yield of methanol.
- B decreases the equilibrium yield of methanol.
- C increases the value of  $K_p$ .
- D decreases the value of  $K_p$ .

**(Total for Question 1 mark)**

8 The equation for the equilibrium between  $\text{NO}_2(\text{g})$  and  $\text{N}_2\text{O}_4(\text{g})$  can be written in two ways.



**or**




Which expression is correct?

- A  $K_c = K'_c$
- B  $K_c = (K'_c)^2$
- C  $K_c = 2(K'_c)$
- D  $K_c = \frac{1}{2}K'_c$

**(Total for Question 1 mark)**

9 4.0 mol of methanoic acid are reacted with 6.0 mol of ethanol.




The equilibrium mixture contains 3.0 mol of  $\text{HCOOC}_2\text{H}_5$ .

The equilibrium constant,  $K_c$ , for the reaction is

- A 0.33
- B 1.0
- C 3.0
- D 4.0

(Total for Question 1 mark)

10 This question is about the equilibrium reaction




Which statement is **not** correct?

- A The units of  $K_p$  are  $\text{atm}^{-2}$ .
- B  $K_p$  increases as temperature is decreased.
- C  $K_p$  increases when the pressure increases.
- D  $K_p$  increases when the total entropy change,  $\Delta S_{\text{total}}$ , increases.

(Total for Question = 1 mark)

11 The equation for the synthesis of methanol is



At equilibrium, when the temperature is 340 K, the total pressure is 20 atm. The moles of each component present at equilibrium are shown in the table below.

Formula	Equilibrium moles / mol	Mole fraction
CO	0.15	0.23
H <sub>2</sub>	0.32	
CH <sub>3</sub> OH	0.18	0.28

(a) The mole fraction of hydrogen in the equilibrium mixture is

(1)

- A 0.23
- B 0.46
- C 0.49
- D 0.92

(b) The numerical value for the equilibrium partial pressure of the carbon monoxide, in atmospheres, is

(1)

- A 3.0
- B 4.6
- C 5.0
- D 9.2


(c) Units for the equilibrium constant,  $K_p$ , for this reaction are

(1)

- A no units
- B atm
- C atm<sup>-1</sup>
- D atm<sup>-2</sup>

**(Total for Question = 3 marks)**


12 What are the units of  $K_c$  for the following equilibrium?



- A atm
- B  $\text{atm}^{-1}$
- C  $\text{dm}^3 \text{mol}^{-1}$
- D  $\text{mol dm}^{-3}$

(Total for Question 1 mark)

13 Consider the equilibrium




Which of the following is true when the total pressure of the system is increased at constant temperature?

		Value of $K_p$	Mole fraction of $\text{PCl}_5(\text{g})$
<input type="checkbox"/>	<b>A</b>	decreases	decreases
<input type="checkbox"/>	<b>B</b>	unaltered	increases
<input type="checkbox"/>	<b>C</b>	decreases	increases
<input type="checkbox"/>	<b>D</b>	unaltered	unaltered

(Total for Question = 1 mark)



14 Iron and steam at high temperature react in a closed vessel to give an equilibrium mixture



Which of the following is the correct expression for  $K_p$ ?

A  $K_p = \frac{P_{\text{H}_2}}{P_{\text{H}_2\text{O}}}$

B  $K_p = \frac{P_{\text{Fe}_3\text{O}_4} P_{\text{H}_2}^4}{P_{\text{Fe}}^3 P_{\text{H}_2\text{O}}^4}$

C  $K_p = \frac{P_{\text{H}_2}^4}{P_{\text{H}_2\text{O}}^4}$

D  $K_p = P_{\text{H}_2}^4$

(Total for Question = 1 mark)