


1 Alcohols **A**, **B**, **C** and **D** are shown below.



(a) Compound **A** is ethanol, a very useful alcohol.

Identify the two main methods used in the industrial production of ethanol.
Write an equation for each method.

method 1

.....

equation

method 2

.....

equation [4]

(b) A student heated each alcohol, **A–D**, with acidified potassium dichromate(VI) as the oxidising agent. With alcohols **A**, **B** and **C**, the colour turned from orange to green.

(i) Identify the organic product and write a balanced equation for the reaction of alcohol **B** with acidified potassium dichromate(VI).


Use [O] to represent the oxidising agent, acidified potassium dichromate(VI).

organic product:

balanced equation:

(ii) The organic product obtained from **C** was analysed by infrared (IR) spectroscopy.

The IR spectrum of the product is shown below.



Use your *Data Sheet* to identify the organic product. Explain your reasoning.

organic product:

reasoning

.....

..... [3]

(c) The student heated alcohol **D** with ethanoic acid in the presence of an acid catalyst. An organic product **E** was formed with a fruity smell.

(i) Name alcohol **D**.

..... [1]

(ii) Name the functional group in the organic product **E**.

..... [1]

(iii) Draw the structure of the organic product **E**.

[2]

[Total: 13]

2 In this question, you are asked to suggest structures for several organic compounds2

(a) Compounds **F**, **G** and **H** are **unbranched** alkenes that are isomers, each with a relative molecular mass of 70.0.

Compounds **F** and **G** are *E/Z* stereoisomers.

Compound **H** is a structural isomer of compounds **F** and **G**.

- Explain what is meant by the terms *structural isomer* and *stereoisomer*.
- Explain why some alkenes have *E/Z* isomerism.
- Analyse this information to suggest possible structures for compounds **F**, **G** and **H**.



In your answer you should make clear how each structure fits with the information given above.


.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

3 Ethanoic acid, CH₃COOH, is used to make esters.

Some information about two of the processes used to make ethanoic acid is given below.

Process 1

This is a one-step process that involves the reaction of methanol with carbon monoxide.



The conditions used are 180°C and 30 atmospheres pressure. A rhodium/iodine catalyst is used.

The percentage yield for this process is 99%.

Process 2

This involves the oxidation of naphtha, a fraction obtained from crude oil.

Liquid naphtha is oxidised using air at a temperature of 180°C and 50 atmospheres pressure. No catalyst is needed.

A large variety of other products are also formed in this oxidation.

(a) Suggest **three** advantages of making ethanoic acid using **Process 1** rather than **Process 2**.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


[3]

(b) The other products formed in **Process 2** are carboxylic acids, aldehydes and ketones.

A research chemist investigates some of these other products of **Process 2**.

(i) The research chemist isolates product, **J**.

The infrared spectrum of **J** is shown below.



The chemist also finds that 0.172 g of a pure sample of **J** contains 2.00×10^{-3} mol of **J**.

Suggest, with reasons, **one** possible structure for **J**.



In your answer you should link the evidence with your explanation.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) The chemist isolates another product, the carboxylic acid, **K**.

K has the molecular formula $C_4H_8O_2$.

Suggest a possible structure and name for **K**.

structure

name [2]

(c) Ethanoic acid is used in the manufacture of the ester, propyl ethanoate.

Describe how ethanoic acid is converted into propyl ethanoate.
Include an equation in your answer.

.....
.....
.....
.....
.....
.....
..... [4]

[Total: 14]

4 Chlorofluorocarbons, CFCs, were once used as propellants in aerosols. CFCs contribute to ozone depletion in the upper atmosphere.

(a) A CFC has the formula CF_2Cl_2 .

State the three-dimensional shape of a CF_2Cl_2 molecule and the F-C-Cl bond angle.

shape

bond angle [2]

(b) Two reasons that CF_2Cl_2 was used as an aerosol propellant are that it has low reactivity and will not hydrolyse in water.

(i) State **one** other reason why CF_2Cl_2 was developed for use as an aerosol.

.....
..... [1]

(ii) Suggest why CF_2Cl_2 does **not** hydrolyse in water.

.....
.....
..... [1]

(c) Explain, with the aid of equations, how the presence of CFCs in the upper atmosphere leads to ozone depletion.

.....
.....
.....
.....
..... [3]

(d) Why are scientists concerned about ozone depletion?

.....
.....
..... [1]

(e) International agreements have reduced the use of CFCs. However the concentration of atmospheric CFCs has hardly changed.

Suggest **two** reasons why.

.....

.....

.....

.....

.....

[2]

[Total: 10]

5 This question is about the six alcohols below.

butan-2-ol
2-methylpentan-3-ol
propan-1-ol

ethane-1,2-diol
2-methylpropan-2-ol
propan-2-ol

(a) Which alcohol is an example of a tertiary alcohol?

..... [1]

(b) Draw the skeletal formula for 2-methylpentan-3-ol.

[1]

(c) Butan-2-ol and 2-methylpropan-2-ol are structural isomers.

(i) What is meant by the term *structural isomer*?

.....
.....
..... [1]

(ii) Draw another structural isomer of these two alcohols.

[1]

(d) Ethane-1,2-diol can be dissolved in water to act as an anti-freeze in car radiators.

Explain why ethane-1,2-diol is very soluble in water.

.....
.....
.....

..... [2]
PhysicsAndMathsTutor.com

- (e) Ethane-1,2-diol is heated under reflux with ethanoic acid and a small amount of H_2SO_4 catalyst. Compound **A** is formed with molecular formula $\text{C}_6\text{H}_{10}\text{O}_4$.

Draw the structure of compound **A**.

[2]

- (f) Butan-2-ol is heated with H_2SO_4 catalyst.

- A mixture of **three** alkenes forms, **B**, **C** and **D**.
- The alkenes **B** and **C** are stereoisomers.

- (i) Draw the structures of the two stereoisomers **B** and **C**.

[2]

- (ii) What type of stereoisomerism is shown by **B** and **C**?

..... [1]

- (iii) Draw the structure of the other alkene, **D**, that is formed in this reaction.

[1]


(g) Alcohol **E** is one of the following alcohols.

butan-2-ol
2-methylpentan-3-ol
propan-1-ol

ethane-1,2-diol
2-methylpropan-2-ol
propan-2-ol

A student oxidises alcohol **E** by heating under reflux with excess acidified potassium dichromate(VI). An organic product **F** is isolated.

The mass spectrum of the alcohol **E** is shown below.



The infrared spectrum of the organic product **F** is shown below.

