

Question			Answer	Marks									
1	(a)		<table border="1"> <tr> <td>E°</td> <td>redox system</td> </tr> <tr> <td>Most negative</td> <td>E</td> </tr> <tr> <td></td> <td>C</td> </tr> <tr> <td>Least negative</td> <td>D</td> </tr> </table> <p style="text-align: center;">✓</p>	E°	redox system	Most negative	E		C	Least negative	D	1	ALL 3 correct for 1 mark
E°	redox system												
Most negative	E												
	C												
Least negative	D												
	(b)	(i)	pH = 0 ✓	1	Guidance								
	(b)	(ii)	<p>H redox system is more negative (e.g. has a more -ve E OR less +ve E OR is -ve electrode) OR H redox system releases electrons (May be in equation, e.g. $H_2 \rightarrow 2H^+ + 2e^-$) ✓</p> <p>Equilibrium shifts to increase $[H^+]$ OR H^+ OR standard hydrogen equation shifts to increase $[H^+]$ OR H^+ ✓</p>	2	<p>ALLOW ORA, ie Ag redox system (D) has more positive E / less negative E</p> <p>ALLOW equilibrium sign</p> <p>IGNORE H is more reactive ORA</p> <p>IGNORE direction of equilibrium shift</p>								
	(b)	(iii)	$H_2 + 2Ag^+ \rightarrow 2Ag + 2H^+$ ✓	1	<p>ALLOW multiples e.g. $\frac{1}{2}H_2 + Ag^+ \rightarrow Ag + H^+$</p> <p>State symbols NOT required ALLOW equilibrium sign</p>								
	(c)	(i)	$\text{AND } \begin{array}{ccccccc} & - & & & & & \\ & \text{Base}_2 & \text{H}_2\text{O} & \rightleftharpoons & \text{HCN} & \text{OH}^- & \\ & \text{Acid } 1 & & & \text{Acid } 2_+ & \text{Base } 1 & \checkmark \\ & \text{CN} & & & & & \end{array}$	1	<p>State symbols NOT required ALLOW CNH and HO^- (i.e. any order)</p> <p>ALLOW 1 and 2 labels the other way around. ALLOW 'just acid' and 'base' labels throughout if linked by lines so that it is clear what the acid-base pairs are.</p>								

Question		Answer	Marks	Guidance
	(c) (ii)	H ⁺ reacts with CN ⁻ OR HCN forms OR equation: H ⁺ + CN ⁻ → HCN (ALLOW ⇌) OR CN ⁻ accepts a proton/H ⁺ OR equilibrium shifts right AND CN ⁻ is removed ✓	1	ALLOW Acid reacts with/removes OH ⁻ ions (to form HCN) ALLOW CNH (i.e. any order) IGNORE other equilibrium comments
	(d) (i)	Fuel reacts with oxygen/oxidant to give electrical energy/voltage ✓	1	ALLOW named fuel. e.g. hydrogen/H ₂ ; ethanol; methanol, etc ALLOW fuel cell requires constant supply of fuel AND oxygen/an oxidant OR fuel cell operates continuously as long as a fuel AND oxygen/an oxidant are added IGNORE 'reactants' 'products' and comments about pollution and efficiency
	(d) (ii)	ethanol is a liquid OR is less volatile OR ethanol is easier to store/transport/stored more safely OR hydrogen is explosive/more flammable OR ethanol has more public/political acceptance ✓	1	Assume that 'it' refers to ethanol ALLOW ORA throughout IGNORE ethanol has a higher boiling point IGNORE H ₂ is a gas IGNORE 'produces no CO ₂ ' OR less pollution IGNORE comments about efficiency IGNORE comments about biomass and renewable
	(d) (iii)	C ₂ H ₅ OH + 3O ₂ → 2CO ₂ + 3H ₂ O ✓	1	Correct species AND balancing needed ALLOW multiples ALLOW C ₂ H ₆ O for formula of ethanol IGNORE state symbols
	(d) (iv)	O ₂ + 4H ⁺ + 4e ⁻ → 2H ₂ O ✓	1	Correct species AND balancing needed ALLOW multiples, e.g. 3O ₂ + 12H ⁺ + 12e ⁻ → 6H ₂ O 2 + 2H ⁺ + 2e ⁻ → H ₂ O ALLOW e (ie no +/2 sign) ALLOW O ₂ + 2H ₂ O + 4e ⁻ → 4OH ⁻ OR 3O ₂ + 6H ₂ O + 12e ⁻ → 12OH ⁻ IGNORE state symbols

Question		Answer	Marks	Guidance
	(d) (v)	oxidation: C from -2 to $+4$ '+' sign not required ✓ reduction: O from 0 to -2 ✓	2	ALLOW $2-$ and $4+$ ALLOW $C^{2-} \rightarrow C^{4+}$ ALLOW 0 and $2-$ ALLOW $O^0 \rightarrow O^{2-}$ ALLOW 1 mark if correct oxidation numbers shown for BOTH C and O but wrong way around <i>(ie C on reduction line and O on oxidation line)</i> IGNORE O_2 reduced IGNORE any reference to electron transfer <i>(not in question)</i>
		Total	13	

Question		Answer	Marks	Guidance
2	(a)	<p>Equations can be in either order</p> <p>$\text{Na}_2\text{O} + \text{H}_2\text{O} \rightarrow 2\text{NaOH} \checkmark$</p> <p>$\text{NaFeO}_2 + 2\text{H}_2\text{O} \rightarrow \text{Fe}(\text{OH})_3 + \text{NaOH} \checkmark$</p>	2	<p>ALLOW multiples throughout IGNORE state symbols</p> <p>ALLOW $\text{Na}_2\text{O} + \text{H}_2\text{O} \rightarrow 2\text{Na}^+ + 2\text{OH}^-$</p> <p>DO NOT ALLOW equations with uncanceled species. e.g. $\text{Na}_2\text{O} + 2\text{H}_2\text{O} \rightarrow 2\text{NaOH} + \text{H}_2\text{O}$</p> <p>ALLOW $2\text{NaFeO}_2 + \text{H}_2\text{O} \rightarrow \text{Fe}_2\text{O}_3 + 2\text{NaOH}$ OR $2 + \text{H}_2\text{O} \rightarrow \text{Fe}_2\text{O}_3 + 2\text{Na}^+ + 2\text{OH}^- \checkmark$</p>

2NaFeO

Question	Answer	Marks	Guidance
(b)	<p>FIRST, CHECK THE ANSWER ON ANSWER LINE IF answer = 33.7%, award 6 marks. IF there is an alternative answer, check to see if there is any ECF credit possible using working below</p> <p>-----</p> <p>amount $\text{S}_2\text{O}_3^{2-}$ used = $0.1000 \times \frac{25.50}{1000}$ = 2.550×10^{-3} (mol) ✓</p> <p>amount I_2 = $2.550 \times 10^{-3} \div 2$ = 1.275×10^{-3} (mol) ✓</p> <p>amount CrO_4^{2-} $\frac{2}{3} \times 1.275 \times 10^{-3}$ OR $1.275 \times 10^{-3} \div 1.5$ = $8.5(00) \times 10^{-4}$ (mol) ✓</p> <p>amount CrO_4^{2-} in original 1000 cm^3 = $40 \times 8.5(00) \times 10^{-4}$ = $3.4(00) \times 10^{-2}$ mol ✓</p> <p>Mass of Cr/Cr^{3+} in ore = $52.0 \times 3.4(00) \times 10^{-2}$ g = 1.768 g ✓</p> <p>percentage Cr in ore = $\frac{1.768}{5.25} \times 100$ = 33.7% ✓</p> <p>MUST be to one decimal place (in the question)</p>	<p>6</p>	<p>FULL ANNOTATIONS MUST BE USED</p> <p>IF a step is omitted but subsequent step subsumes previous, then award mark for any missed step Working: at least 3 SF throughout until final % mark BUT ignore trailing zeroes, ie for 0.490 allow 0.49</p> <p>-----</p> <p>ECF answer above $\div 2$</p> <p>ECF answer above $\div 1.5$</p> <p>ECF answer above $\times 40$</p> <p>ECF answer above $\times 52.0$ IMPORTANT: The last two marks are ONLY available by using 52.0 for Cr</p> <p>-----</p> <p>Common ECFs:</p> <p>0.8% $\times 40$ missing 5 marks (scaling error)</p> <p>0.84% $\times 40$ missing 4 marks (scaling error and 2 DP)</p> <p>33.68% 5 marks (2 DP)</p> <p>16.8% 5 marks (divide Cr somewhere by 2)</p> <p>144.9%; 72.5% 4 marks (Final 2 marks unavailable) Use of $M(\text{Fe}(\text{CrO}_2)_2) = 223.8$ instead of $M(\text{Cr})$.</p>

Question		Answer	Marks	Guidance
	(c)	<p><i>Overall:</i></p> $\text{CrO}_4^{2-} + 3\text{I}^- + 4\text{H}_2\text{O} \rightarrow \text{Cr}^{3+} + 1\frac{1}{2}\text{I}_2 + 8\text{OH}^- \checkmark$ <p>CrO</p> <p><i>Half equations:</i></p> $\text{CrO}_4^{2-} + 4\text{H}_2\text{O} + 3\text{e}^- \rightarrow \text{Cr}^{3+} + 8\text{OH}^- \checkmark$ <p>CrO</p> $2\text{I}^- \rightarrow \text{I}_2 + 2\text{e}^- \checkmark$	3	<p>ALLOW multiples and equilibrium signs throughout IGNORE state symbols throughout</p> <p>e.g. $2\text{CrO}_4^{2-} + 6\text{I}^- + 8\text{H}_2\text{O} \rightarrow 2\text{Cr}^{3+} + 3\text{I}_2 + 16\text{OH}^-$</p> <p>ALLOW equation using H^+. i.e.</p> <p>OR $\text{CrO}_4^{2-} + 3\text{I}^- + 8\text{H}^+ \rightarrow \text{Cr}^{3+} + 1\frac{1}{2}\text{I}_2 + 4\text{H}_2\text{O}$ $2\text{CrO}_4^{2-} + 6\text{I}^- + 16\text{H}^+ \rightarrow 2\text{Cr}^{3+} + 3\text{I}_2 + 8\text{H}_2\text{O}$ + </p> <p>ALLOW CrO_4^{2-} half equation using H^+. i.e.</p> $\text{CrO}_4^{2-} + 8\text{H}^+ + 3\text{e}^- \rightarrow \text{Cr}^{3+} + 4\text{H}_2\text{O}$ <p>CrO</p>
Total			11	

Question		Answer	Marks	Guidance	
3	(a)	<p>Definition The e.m.f. (of a half-cell) compared with/connected to a (standard) hydrogen half-cell/(standard) hydrogen electrode ✓</p> <p>Standard conditions <i>Units essential</i> Temperature of 298 K / 25°C AND (solution) concentrations of 1 mol dm⁻³ AND pressure of 100 kPa OR 10⁵ Pa OR 1 bar ✓</p>	2	<p>As alternative for e.m.f., ALLOW voltage OR potential difference OR p.d. OR electrode potential OR reduction potential OR redox potential ALLOW /(standard) hydrogen cell IGNORE S.H.E. (as abbreviation for standard hydrogen electrode)</p> <p>ALLOW 1M DO NOT ALLOW 1 mol ALLOW 1 atmosphere/1 atm OR 101 kPa OR 101325 Pa</p>	
	(b)	(i)	$2\text{Ag}^+(\text{aq}) + \text{Cu}(\text{s}) \rightarrow 2\text{Ag}(\text{s}) + \text{Cu}^{2+}(\text{aq}) \checkmark$	1	<p>State symbols not required ALLOW \rightleftharpoons provided that reactants on LHS</p>
	(b)	(ii)	<p>Assume Cu²⁺ Cu OR Cu half cell unless otherwise stated.</p> <p>[Cu²⁺] decreases OR < 1 mol dm⁻³ AND Equilibrium (shown in table) shifts to left ✓</p> <p>more electrons are released by Cu ✓</p> <p>The cell has a bigger difference in <i>E</i> ✓</p>	3	<p>FULL ANNOTATIONS MUST BE USED</p> <hr/> <p>ALLOW [Cu²⁺] less than standard concentration/1 mol dm⁻³ DO NOT ALLOW water reacts with Cu²⁺ OR Cu</p> <p>ALLOW <i>E</i> (for Cu²⁺ Cu) is less positive / more negative /decreases IGNORE standard electrode potential (<i>Cell no longer standard</i>) IGNORE <i>E</i>^o decreases CARE DO NOT ALLOW statements about silver <i>E</i> changing (CON)</p> <p>IGNORE just 'cell potential increases' (in the question) <i>The final mark is more subtle and is a consequence of the less positive E value of the copper half cell</i></p>

	(c)	(i)	no/less CO ₂ OR H ₂ O is only product OR greater efficiency ✓	1	IGNORE less pollution IGNORE less carbon emissions IGNORE less fossil fuels used IGNORE no/less greenhouse gas OR no global warming (H ₂ O vapour is a greenhouse gas)
	(c)	(ii)	liquefied/as a liquid AND under pressure/pressurised ✓	1	IGNORE adsorption or absorption IGNORE low temperature DO NOT ALLOW liquidise <i>processes are described in the question</i>
	(d)	(i)	$E = -2.31$ (V) ✓	1	– sign AND 2.31 required for the mark
	(d)	(ii)	$4\text{Al(s)} + 4\text{OH}^{\text{-}}(\text{aq}) + 3\text{O}_2(\text{g}) + 6\text{H}_2\text{O(l)} \rightarrow 4\text{Al(OH)}_4^{\text{-}}(\text{aq})$ species ✓ balance ✓	2	IGNORE state symbols ALLOW multiples ALLOW 1 mark for an equation in which OH ⁻ are balanced but have not been cancelled, e.g. $4\text{Al(s)} + 16\text{OH}^{\text{-}}(\text{aq}) + 3\text{O}_2(\text{g}) + 6\text{H}_2\text{O(l)} \rightarrow 4\text{Al(OH)}_4^{\text{-}}(\text{aq}) + 12\text{OH}^{\text{-}}(\text{aq})$ ALLOW 1 mark if charge on Al(OH) ₄ is omitted, i.e. $4\text{Al(s)} + 4\text{OH}^{\text{-}}(\text{aq}) + 3\text{O}_2(\text{g}) + 6\text{H}_2\text{O(l)} \rightarrow 4\text{Al(OH)}_4(\text{aq})$ ALLOW 1 mark for an 'correct equation' reversed, i.e. $4\text{Al(OH)}_4^{\text{-}}(\text{aq}) \rightarrow 4\text{Al(s)} + 4\text{OH}^{\text{-}}(\text{aq}) + 3\text{O}_2(\text{g}) + 6\text{H}_2\text{O(l)}$
			Total	11	

Question		Answer	Marks	Guidance
4	(a)	$\text{Fe}_2\text{O}_3 + 3\text{Cl}_2 + 10\text{OH}^- \rightarrow 2\text{FeO}_4^{2-} + 6\text{Cl}^- + 5\text{H}_2\text{O} \checkmark\checkmark$ <p>First mark for all 6 species Second mark for balancing</p>	2	<p>ALLOW multiples ALLOW oxidation half equation for two marks $\text{Fe}_2\text{O}_3 + 10\text{OH}^- \rightarrow 2\text{FeO}_4^{2-} + 5\text{H}_2\text{O} + 6\text{e}^-$ Correct species would obtain 1 mark – question: equation for oxidation</p> <p>ALLOW variants forming H⁺ for 1 mark, e.g: $\text{Fe}_2\text{O}_3 + 3\text{Cl}_2 + 5\text{OH}^- \rightarrow 2\text{FeO}_4^{2-} + 6\text{Cl}^- + 5\text{H}^+$ $\text{Fe}_2\text{O}_3 + 3\text{Cl}_2 + 5\text{OH}^- \rightarrow 2\text{FeO}_4^{2-} + 5\text{HCl} + \text{Cl}^-$</p>
	(b)	$\text{Ba}^{2+}(\text{aq}) + \text{FeO}_4^{2-}(\text{aq}) \rightarrow \text{BaFeO}_4(\text{s}) \checkmark$	1	<p>Balanced ionic equation AND state symbols required DO NOT ALLOW +2 or –2 for ionic charges</p>
	(c)	<p>Reason can ONLY be correct from correct reducing agent ----- reducing agent: I⁻ OR KI ✓</p> <p>I⁻ adds/donates/loses electrons AND to FeO₄²⁻ OR to BaFeO₄ OR to Fe(VI) or to Fe(+6) ✓ ALLOW Fe(6+) OR Fe⁶⁺</p>	2	<p>IGNORE H⁺ OR acidified ALLOW iodide/potassium iodide but DO NOT ALLOW iodine</p> <p>ALLOW I⁻ loses electrons AND to form I₂</p> <p>ALLOW Fe(6+) OR Fe⁶⁺</p>

(d)

FIRST, CHECK THE ANSWER ON ANSWER LINE
IF answer = 51.8%, award **4** marks.

$$n(\text{S}_2\text{O}_3^{2-}) \text{ used} = 0.1000 \times \frac{26.4}{1000} = 2.64 \times 10^{-3} \text{ (mol)} \checkmark$$

$$n(\text{FeO}_4^{2-}) = \frac{1}{2} \times \frac{2}{3} \times 2.64 \times 10^{-3} = 8.8(0) \times 10^{-4} \text{ (mol)} \checkmark$$

Mass BaFeO₄ in sample
 $= 8.8 \times 10^{-4} \times 257.1 \text{ g} = 0.226248 \text{ g} \checkmark$

$$\% \text{ purity} = \frac{0.226248}{0.437} \times 100 = 51.8\% \checkmark$$

MUST be to **one** decimal place (in the question)

As an alternative for the final two marks, **ALLOW**:

$$\text{Theoretical amount of BaFeO}_4 = \frac{0.437}{257.1} = 0.00170 \text{ (mol)} \checkmark$$

$$\% \text{ purity} = \frac{8.8 \times 10^{-4}}{1.70 \times 10^{-3}} \times 100 = 51.8\% \checkmark$$

FULL ANNOTATIONS MUST BE USED

For alternative answers, look first at common **ECFs** below.
Then check for **ECF** credit possible using working below
IF a step is omitted but subsequent step subsumes previous,
then award mark for any missed step

Working must be to at least 3 SF throughout until final % mark

BUT ignore trailing zeroes, ie for 0.880 allow 0.88

ECF answer above $\times \frac{1}{2} \times \frac{2}{3}$

This mark may be seen in 2 steps via I_2 but the mark is for both steps combined

ECF $257.1 \times$ answer above

ECF $\frac{\text{answer above}}{0.437} \times 100$

ALLOW 51.7% FROM 0.226 g BaFeO₄ (earlier rounding)

Common ECFs:

No $\times \frac{2}{3}$ for $n(\text{FeO}_4^{2-})$:

$$\% \text{ purity} = 77.7\%/77.6\% \quad 3 \text{ marks}$$

No $\div 2$ for $n(\text{FeO}_4^{2-})$:

$$\% \text{ purity} = 25.9\% \quad 3 \text{ marks}$$

24.6 used instead of 26.4:

$$\% \text{ purity} = 48.2\% \quad 3 \text{ marks}$$

4

	(e)	<p>gas: O₂ ✓</p> <p>precipitate: Fe(OH)₃ ✓</p> <p>equation: 2FeO₄²⁻ + 5H₂O → 1½O₂ + 2Fe(OH)₃ + 4OH⁻</p> <p>OR 2FeO₄²⁻ + H₂O + 4H⁺ → 1½O₂ + 2Fe(OH)₃ ✓</p>	<p>3</p>	<p>DO NOT ALLOW names IGNORE a balancing number shown before a formula</p> <p>ALLOW Fe(OH)₃(H₂O)₃</p> <p>ALLOW multiples ALLOW 2FeO₄²⁻ + 11H₂O → 1½O₂ + 2Fe(OH)₃(H₂O)₃ + 4OH⁻</p>
		Total	12	