

Question		Expected answers	Marks	Additional guidance
1	a	Complete circuit with electrodes to voltmeter AND salt bridge between solutions ✓ Fe ³⁺ /Fe ²⁺ half-cell with Pt electrode AND 1 mol dm ⁻³ /1 M Fe ²⁺ and 1 mol dm ⁻³ /1 M Fe ³⁺ ✓ Ni electrode in (1 mol dm ⁻³) Ni ²⁺ half-cell ✓	3	circuit shown must be complete, <i>i.e. must be capable of working</i> salt bridge must be labelled. electrodes AND salt bridge must dip into/touch both solutions ALLOW cells drawn either way around ALLOW Fe ³⁺ /Fe ²⁺ 1 mol dm ⁻³ /1 M /1 molar ALLOW BOTH solutions same concentration/equimolar DO NOT ALLOW 1 mol OR 1 dm ⁻³ IGNORE any temperature or pressure, even if wrong
	ii	1.02 V AND – sign ✓ 0.49 V AND + sign ✓	2	IGNORE any sign BEFORE cell potential ALLOW 1 mark for correct values AND signs BOTH the wrong way round: <i>i.e.</i> 1.02 V AND + sign AND 0.49 V AND – sign
	b	Cell A (based on 1 and 2) Ni + 2Fe ³⁺ → Ni ²⁺ + 2Fe ²⁺ ✓ Cell B (based on 1 and 3) 2Cr + 3Ni ²⁺ → 2Cr ³⁺ + 3Ni ✓ concentrations (of the ions in each cell) change OR concentrations are not standard ✓	3	In equations, ALLOW equilibrium sign, = instead of → Equations are required for the first two marking points ALLOW Ni → Ni ²⁺ + 2e ⁻ ALLOW Ni ²⁺ + 2e ⁻ → Ni ALLOW any statement that a concentration is changing IGNORE 'non-standard conditions'
	c	i	1	ALLOW MH → M + H ⁺ + e ⁻
		ii	1	DO NOT ALLOW adsorbed into the solid CON DO NOT ALLOW just 'as a liquid'
Total			10	

Question		Expected answers	Marks	Additional guidance
2	a	$\text{Fe}_2\text{O}_3 + 6\text{H}^+ \longrightarrow 2\text{Fe}^{3+} + 3\text{H}_2\text{O} \checkmark$	1	<p>ALLOW $\text{Fe}_2\text{O}_3 + 6\text{HCl} \longrightarrow 2\text{FeCl}_3 + 3\text{H}_2\text{O}$ OR $\text{Fe}_2\text{O}_3 + 6\text{HCl} \longrightarrow 2\text{Fe}^{3+} + 6\text{Cl}^- + 3\text{H}_2\text{O}$</p> <p>ALLOW correct multiples</p> <p>IGNORE state symbols</p> <p>DO NOT ALLOW Fe_2Cl_6 as a product</p>
	b	<p>$\text{Sn}^{2+} + 2\text{Fe}^{3+} \longrightarrow \text{Sn}^{4+} + 2\text{Fe}^{2+} \checkmark$</p> <p>$6\text{Fe}^{2+} + \text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ \longrightarrow$ $6\text{Fe}^{3+} + 2\text{Cr}^{3+} + 7\text{H}_2\text{O} \checkmark$</p>	2	<p>IGNORE state symbols</p> <p>ALLOW overall equations: $\text{SnCl}_2 + 2\text{FeCl}_3 \longrightarrow \text{SnCl}_4 + 2\text{FeCl}_2$</p> <p>$6\text{FeCl}_2 + \text{K}_2\text{Cr}_2\text{O}_7 + 14\text{HCl} \rightarrow$ $6\text{FeCl}_3 + 2\text{CrCl}_3 + 2\text{KCl} + 7\text{H}_2\text{O}$</p> <p>ALLOW correct multiples</p>

Question	Expected answers	Marks	Additional guidance
c	<p>FIRST, CHECK THE ANSWER ON ANSWER LINE IF answer = 54.6%, award 5 marks</p> <p>-----</p> <p>Amount Fe²⁺ in 250 cm³ solution – 3 marks</p> <p>amount Cr₂O₇²⁻ used = $0.0200 \times \frac{26.5}{1000}$ = 5.30×10^{-4} (mol) ✓</p> <p>amount Fe²⁺ = $6 \times 5.30 \times 10^{-4}$ = 3.18×10^{-3} mol ✓</p> <p>amount Fe²⁺ in original 250 cm³ = $10 \times 3.18 \times 10^{-3}$ = 3.18×10^{-2} (mol) ✓</p>		<p>ANNOTATIONS MUST BE USED IF there is an alternative answer, 1st check common errors below. Then see if there is any ECF credit possible using working below</p> <p>-----</p> <p>Working must be to at least 3 SF throughout BUT ignore trailing zeroes, <i>i.e.</i> for 0.490 allow 0.49</p> <p>ALLOW ECF from different Fe²⁺ ratio in equation from 8(b) BUT still ALLOW 6 : 1 even from different ratio in equation If no equation use actual 6 : 1 ratio DO NOT AWARD 'ratio mark' at all for use of 1 : 1 ratio – <i>makes problem easier</i></p> <p>ECF 10 × answer above</p>
	<p>% Fe in ore – 2 marks</p> <p>mass of Fe in ore = $55.8 \times 3.18 \times 10^{-2}$ g = 1.77444 g ✓</p>		<p>ECF 55.8 × answer above</p> <p>IF answer above has not been used AND × 55.8, DO NOT ALLOW this mark but do ALLOW final %</p> <p>IF answer above AND 55.8 are BOTH not used, then DO NOT ALLOW ANY further marks</p>
	<p>percentage Fe in ore = $\frac{1.77444}{3.25} \times 100$ = 54.6% ✓</p>	5	<p>ECF $\frac{\text{answer above}}{3.25} \times 100$</p> <p>ALLOW 54.5% (from 1.77 g) AND any answer with > 1 decimal place that rounds back to 54.5 OR 54.6</p>
			<p>COMMON ERRORS</p> <p>5.46 ✓✓✓✓ × 10 omitted 51.5 ✓✓✓✓ titre taken as 25.0 156.2 ✓✓✓✓ × 159.6 instead of 55.8 15.62 ✓✓✓ × 159.6 and × 10 omitted 45.5 ✓✓✓✓ 5 : 1 ratio 1.52 ✓✓✓✓ ÷ 6 instead of × 6</p>

Question	Expected answers	Marks	Additional guidance
d	E^\ominus for MnO_4^- is more positive/greater than Cl_2 OR E^\ominus for $\text{Cr}_2\text{O}_7^{2-}$ is less positive/smaller than Cl_2 ✓ MnO_4^- reacts with Cl^- OR HCl (forming Cl_2 gas) OR $\text{Cr}_2\text{O}_7^{2-}$ does not react with Cl^- ions ✓	2	ORA: E^\ominus for Cl_2 is less positive/smaller than MnO_4^- OR E^\ominus for Cl_2 is more positive/greater than $\text{Cr}_2\text{O}_7^{2-}$
Total		10	

Question			Answer	Mark	Guidance
3	(a)	(i)	<p>Complete circuit with electrodes to voltmeter AND salt bridge between solutions ✓</p> <p>Sn⁴⁺/Sn²⁺ half cell with Pt electrode AND both solutions labelled as 1 mol dm⁻³ / 1M</p> <p>H⁺/H₂ half cell with Pt electrode AND H⁺ solution labelled as 1 mol dm⁻³ / 1M ✓</p>	3	<p>ANNOTATE WITH TICKS AND CROSSES, etc circuit shown must be complete, <i>ie</i> must be capable of working salt bridge must be labelled and must dip into both solutions</p> <p>ALLOW concentration label of 'equimolar' or similar wording for Sn⁴⁺/Sn²⁺ half cell</p> <p>ALLOW any strong acid</p> <p>IF both half cells are correct with no concentrations, ALLOW 1 out of the 2 marks available for the 2 half cells</p> <p>IGNORE any stated temperature or pressure, even if wrong</p>
		(ii)	<p>2Cr + 3Sn⁴⁺ → 2Cr³⁺ + 3Sn²⁺ ✓ Cr + 3Cu⁺ → Cr³⁺ + 3Cu ✓ Sn²⁺ + 2Cu⁺ → Sn⁴⁺ + 2Cu ✓</p> <p>Conditions not standard OR concentrations not 1 mol dm⁻³ ✓</p> <p>High activation energy OR slow rate ✓</p>	5	<p>ANNOTATE WITH TICKS AND CROSSES, etc Correct species AND balancing needed for each mark ALLOW equations as shown with equilibrium sign ALLOW multiples but electrons must not be shown</p> <p>IF three equations have correct species but no balancing, AWARD 1 mark</p> <p>ALLOW not favoured kinetically</p>
	(b)	(i)	CH ₃ OH + 1½O ₂ → CO ₂ + 2H ₂ O ✓	1	<p>Correct species AND balancing needed ALLOW multiple, <i>ie</i> 2CH₃OH + 3O₂ → 2CO₂ + 4H₂O ALLOW CH₄O for formula of methanol</p>
		(ii)	CH ₃ OH + H ₂ O → 6H ⁺ + 6e ⁻ + CO ₂ ✓	1	
		(iii)	less CO ₂ OR less greenhouse gases ✓ greater efficiency ✓	2	<p>ALLOW no CO₂ OR no greenhouse gases ALLOW (very) efficient IGNORE less pollution OR 'renewable fuels'</p>
		(iv)	methanol is a liquid AND methanol is easier to store/transport ✓	1	<p>Both points required for mark Response MUST state that methanol is a liquid IGNORE methanol has a higher boiling point Assume that 'it' refers to methanol IGNORE safety issues, eg H₂ leakage, flammability, explosive</p>
Total				13	

Question			Answer	Marks	Guidance
4	(a)	(i)	<p> $\text{Fe}^+(\text{g}) + 2\text{I}(\text{g}) + \text{e}^- \checkmark$ $\text{Fe}(\text{g}) + 2\text{I}(\text{g}) \checkmark$ $\text{Fe}(\text{s}) + \text{I}_2(\text{s}) \checkmark$ $\text{Fe}^{2+}(\text{g}) + 2\text{I}^-(\text{g}) \checkmark$ </p> <p>Mark each marking point independently</p>	4	<p>Correct species AND state symbols required for each marks</p> <p>ALLOW e for e^-</p> <p>TAKE CARE: In top left box, e^- may be in centre of response and more difficult to see than at end.</p> <p>There is only ONE correct response for each line <i>From the gaps in the cycle, there is NO possibility of any ECF</i></p>

Question		Answer	Marks	Guidance
	(a) (ii)	<p>(The enthalpy change that accompanies) the formation of one mole of a(n ionic) compound from its gaseous ions (under standard conditions) ✓✓</p> <p>Award marks as follows. 1st mark: formation of compound from gaseous ions 2nd mark: one mole for compound only</p> <p>DO NOT ALLOW 2nd mark without 1st mark</p> <p>DO NOT ALLOW any marks for a definition for enthalpy change of formation BUT note the two concessions in guidance</p>	2	<p>IGNORE 'Energy needed' OR 'energy required'</p> <p>ALLOW one mole of compound is formed/made from its gaseous ions</p> <p>ALLOW as alternative for compound: lattice, crystal, substance, solid</p> <p>IGNORE: $\text{Fe}^{2+}(\text{g}) + 2\text{I}^{-}(\text{g}) \longrightarrow \text{FeI}_2(\text{s})$ (Part of cycle)</p> <p>ALLOW 1 mark for absence of 'gaseous' only, i.e. the formation of one mole of a(n ionic) compound from its ions (under standard conditions) ✓</p> <p>ALLOW 1 mark for ΔH_f definition with 'gaseous': the formation of one mole of a(n ionic) compound from its gaseous elements (under standard conditions) ✓</p>

Question		Answer	Marks	Guidance																						
(a)	(iii)	<p>FIRST, CHECK THE ANSWER ON ANSWER LINE IF answer = -2473 (kJ mol⁻¹) award 2 marks</p> <p>-----</p> $(-113) = 416 + (2 \times +107) + 759 + 1561 + (2 \times -295) + \Delta H_{LE}(\text{FeI}_2)$ <p>OR</p> $\Delta H_{LE}(\text{FeI}_2) =$ $-113 - (416 + (2 \times +107) + 759 + 1561 + (2 \times -295))$ <p>OR $-113 - 2360 \checkmark$</p> $= -2473 \checkmark \text{ (kJ mol}^{-1}\text{)}$	2	<p>IF there is an alternative answer, check to see if there is any ECF credit possible using working below. See list below for marking of answers from common errors</p> <p>-----</p> <p>ALLOW for 1 mark:</p> <table> <tr> <td>+2473</td> <td>wrong sign</td> </tr> <tr> <td>-2661</td> <td>107 and -295 used instead of 2 x 107 and 2 x -295</td> </tr> <tr> <td>-236</td> <td>+107 used instead of 2 x 107</td> </tr> <tr> <td>-276</td> <td>-295 used instead of 2 x -295</td> </tr> <tr> <td>-365</td> <td>wrong sign for 295</td> </tr> <tr> <td>-224</td> <td>wrong sign for 113</td> </tr> <tr> <td>-164</td> <td>wrong sign for 416</td> </tr> <tr> <td>-204</td> <td>wrong sign for 2 x 107</td> </tr> <tr> <td>-95</td> <td>wrong sign for 750</td> </tr> <tr> <td>+64</td> <td>wrong sign for 1561</td> </tr> <tr> <td>-365</td> <td>wrong sign for 2 x -295</td> </tr> </table> <p>Any other number: CHECK for ECF from 1st marking point for expressions with ONE error only e.g. one transcription error: e.g. +461 instead of +416</p>	+2473	wrong sign	-2661	107 and -295 used instead of 2 x 107 and 2 x -295	-236	+107 used instead of 2 x 107	-276	-295 used instead of 2 x -295	-365	wrong sign for 295	-224	wrong sign for 113	-164	wrong sign for 416	-204	wrong sign for 2 x 107	-95	wrong sign for 750	+64	wrong sign for 1561	-365	wrong sign for 2 x -295
+2473	wrong sign																									
-2661	107 and -295 used instead of 2 x 107 and 2 x -295																									
-236	+107 used instead of 2 x 107																									
-276	-295 used instead of 2 x -295																									
-365	wrong sign for 295																									
-224	wrong sign for 113																									
-164	wrong sign for 416																									
-204	wrong sign for 2 x 107																									
-95	wrong sign for 750																									
+64	wrong sign for 1561																									
-365	wrong sign for 2 x -295																									
(b)	(i)	$\text{Fe}^{2+}: 1s^2 2s^2 2p^6 3s^2 3p^6 3d^6 \checkmark$ $\text{Br}^-: 1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 \checkmark$	2	<p>ALLOW 4s before 3d, ie $1s^2 2s^2 2p^6 3s^2 3p^6 4s^2 3d^{10} 4p^6$ ALLOW $1s^2$ written after answer prompt (<i>ie</i> $1s^2$ twice) ALLOW upper case D, etc and subscripts, e.g.4S₂3D₁ ALLOW for Fe^{2+}4s⁰ DO NOT ALLOW [Ar] as shorthand for $1s^2 2s^2 2p^6 3s^2 3p^6$</p> <p>Look carefully at $1s^2 2s^2 2p^6 3s^2 3p^6$ – there may be a mistake</p>																						

Question		Answer	Marks	Guidance
(b)	(ii)	<p>With Cl₂ AND Br₂ AND I₂ products are Fe²⁺ (AND halide ion) FeCl₂ AND FeBr₂ AND FeI₂ ✓</p> <p>OR Evidence that two electrode potentials have been compared for at least ONE reaction, ✓ e.g. Fe -0.44 AND Cl₂ +1.36 e.g. Iron has more/most negative electrode potential</p> <p>With Cl₂ AND Br₂, products are Fe³⁺ (AND halide ion) FeCl₃ AND FeBr₃ ✓</p>	3	<p>FULL ANNOTATIONS NEEDED</p> <p>ALLOW products within equations (even if equations are not balanced) IF stated, IGNORE reactants</p> <p>ALLOW response in terms of positive 'cell reactions', e.g. Fe + Cl₂ → Fe²⁺ + 2Cl⁻ E = (+)1.80 V</p> <p>IGNORE comments about reducing and oxidising agents and electrons</p>
(c)		<p>BOTH EQUATIONS REQUIRE IONS PROVIDED IN QUESTION</p> <p>Reaction 1: 2 marks 1st mark for ALL CORRECT species e.g.: Fe²⁺ + NO₃⁻ + H⁺ → Fe³⁺ + NO + H₂O</p> <p>2nd mark for CORRECT balanced equation 3Fe²⁺ + NO₃⁻ + 4H⁺ → 3Fe³⁺ + NO + 2H₂O ✓✓</p> <p>-----</p> <p>Reaction 2: 1 mark ₂O)₆]²⁺ + NO → [Fe(H₂O)₅NO]²⁺ + H₂O ✓</p>	3	<p>ALLOW correct multiples throughout ALLOW equilibrium signs in all equations</p> <p>For 1st mark, IGNORE e⁻ present</p> <p>Check carefully for correct charges</p>
		[Fe(H	Total	16