


Rings, Polymers and Analysis

Arenes Mark Scheme /114

1. (a)




ALLOW $C_6H_6 + Br_2 \rightarrow C_6H_5Br + HBr$

DO NOT ALLOW multiple substitution

DO NOT ALLOW Br^+

1

(b) (i) White precipitate **OR** white solid **OR** white crystals ✓



DO NOT ALLOW colourless

DO NOT ALLOW white ppt and bubbles

DO NOT ALLOW

$Br_3C_6H_2OH$ **OR** 2,4,6-tribromophenol **OR** tribromophenol

2

(ii) 1,2-Dibromocyclohexane ✓

ALLOW 1,2dibromocyclohexane **OR** 1-2dibromocyclohexane

OR 12dibromocyclohexane **OR** cyclo-1,2-dibromohexane

DO NOT ALLOW dibromocyclohexane **OR** $C_6H_{10}Br_2$ **OR**

structures

1

- (iii) **MUST** spell delocalised/delocalized or localised/localized correctly once in the answer to obtain all 5 marks

benzene electrons or π -bonds are delocalised ✓

ALLOW diagram to show overlap of all 6 p-orbitals for delocalisation

DO NOT ALLOW benzene has delocalised structure or ring

phenol a lone or non-bonded pair of electrons on the oxygen or the OH group is (partially) delocalised into the ring ✓

ALLOW diagram to show movement of lone pair into ring for phenol

cyclohexene electrons are localised **OR** delocalised between two carbons ✓

ALLOW diagram or description of overlap of 2 adjacent p-orbitals for bonding in cyclohexene

DO NOT ALLOW cyclohexene has a C=C double bond

IGNORE slip if cyclohexene is written as cyclohexane but π -bonding correctly described

benzene has a lower **electron density** **OR** phenol has a higher electron density **OR** cyclohexene has a higher electron density ✓

DO NOT ALLOW charge density **OR** electronegativity instead of electron density


benzene cannot **polarise** or induce a dipole in Br₂ **OR** phenol can polarise the Br₂ **OR** cyclohexene can polarise Br₂ or the Br-Br bond ✓

ALLOW Br ^{δ^+} **OR** electrophile Br⁺ as alternate to polarise

5



[9]

2. (a) (i)




1

- (ii) Introduces a permanent dipole on Cl₂ / forms Cl⁺/



1

(iii)



correct dipole / Cl^+ (1)

curly arrow from benzene ring to Cl^+ / $Cl^{\delta+}$ (1)

intermediate (1)

curly arrow from H to regenerate benzene ring in intermediate (1)

H^+ as other product (1)

4

(iv) electrophilic substitution (1)

with electrophilic spelt correctly

1

(b) In benzene, π electrons are delocalised/spread out (1)

In alkenes, π electrons are concentrated between 2 carbons (1)

Electrophiles attracted more to greater electron density in alkenes (1)

3

[10]

3. Discussion of the π -bonding


p-orbitals overlap (1)

above and below the ring (1)

(to form) π -bonds / orbitals (1)

any of the first three marks are available from a labelled diagram

eg



(π -bonds / electrons) are delocalised (1)

4 marks

Other valid points – any two of:

- ring is planar /
- C-C bonds are equal length / have intermediate length/strength between C=C and C-C /
- σ -bonds are between C-C and/or C-H
- bond angles are 120°

6

MAX 2 out of 4 marks (1)(1)


Quality of written communication

two or more sentences with correct spelling, punctuation and grammar

1

[7]

4. (i)



3

(ii) dye / colouring / indicator (1)

1

(iii) phenylamine (1)
 $NaNO_2 / HNO_2$ (1) + HCl (1)
 $< 10^\circ C$ (1)
 add to alkaline phenol (1)

5

[9]

5. **bonding in benzene**
 overlap of p-orbitals / π bonds/electrons (or labelled) (1)



above and below the ring (or shown in a diagram) (1)
 electrons are delocalised (or labelled) (1)
 C–C bonds are: same length/strength / in between single and double / σ -bonded **AW** (1)

greater reactivity of phenol

(the ring is activated because ...)
lone pair from O is delocalised into the ring (1)
 so electron density (of the ring) is increased (1)
 so electrophiles are more attracted (to the ring) / dipole in electrophile more easily induced (1)
 (**NOT** just more easily “attacked” or “susceptible”)

Quality of written communication mark for at least two complete sentences in which the meaning is clear with correct spelling, punctuation and grammar (1)

8


[8]

6. (a) Correct structure of 3-nitrophenol or any multiple nitrated phenol (1) 1
- (b) M_r phenol (C_6H_6O) = 94.0 (1)
 M_r 4-nitrophenol ($C_6H_5NO_3$) = 139.0 (1)
 expected mass/moles of nitrophenol from 100 g =
 148 g/1.06 mol (or ecf from wrong M_r s) (1)
 at 27% yield gives 40 / 39.9 (g) (or ecf) (1) 4
last mark is for $0.27 \times$ expected mass to 2 or 3 sf

- (c) **conditions for nitration of benzene:**
 HNO₃ is concentrated (1)
 conc H₂SO₄ is present (1)
 heating or stated temp above 50°C (1) 3
- explanation for greater reactivity of phenol**
 lone pair from O atom is delocalised into the ring (1)
 greater (π) electron density around the ring (1)
 (the benzene ring in phenol) is activated (1)
- attracts electrophiles/⁺NO₂ more / makes it more
 susceptible to electrophiles **AW** (1) 4
- quality of Written Communication** mark for at least two legible
 sentences with correct spelling, punctuation and grammar 1

[13]

7.



4

[4]

8. **delocalised electrons**
 electrons are spread over more than two atoms **AW** (1)
- π-bond**
 formed by overlap of p-orbitals/ diagram to show (1) 2


[2]

9. (a) (i) **bromine as an electrophile**
 an electrophile accepts an electron pair (1)
NOT a lone pair

bromine is polarised/has + charge (centre)/dipole on Br-Br/Br⁺
 shown in diagram (1)

appropriate diagram showing a curly arrow from a double/
 π bond to the Br ^{δ^+} /Br⁺ (1)

eg



3

- (ii) **comparison of reactivity of cyclohexene and benzene**

benzene is (more) stable / more energy required (1)

benzene (π) electrons are delocalised (1)

benzene has lower electron/- charge density (1)

so bromine is less polarised / attracted to it /
 benzene is less susceptible to electrophiles (1)

ora for cyclohexene

4

quality of written communication mark for any **two** of the the terms:

delocalised/localised, π -electrons/bonds/system, electron density,
 dative covalent, activation/stabilisation energy, halogen carrier,
 heterolytic fission, addition/substitution, polarity used appropriately (1)

1

- (b) (i) iodobenzene because ...

Br is more electronegative than I (1) **ora**

so the I atom will be positive / δ^+ /the electrophile (1)

2

- (ii) $C_6H_6 + IBr \rightarrow C_6H_5I + HBr$ (1)

or ecf giving $C_6H_5Br + HI$

1

[11]

10. (a) (i) NaOH / Na (1)


1

- (ii) $C_6H_5OH + NaOH \rightarrow C_6H_5O^-Na^+ + H_2O$ /

$C_6H_5OH + Na \rightarrow C_6H_5O^-Na^+ + \frac{1}{2}H_2$ (1)

1

(b) (i)




(1)

allow a dipole on just one C=O bond

1

(ii)



(1)(1)

2

(iii) lone/electron pair from oxygen is delocalised into the ring /interacts with π -electrons (1)

increases π -electron density / negative charge
(around the ring) (1)

attracts electrophiles more (1)

3

(c) M_r salicylic acid = 138 (1)

moles (in 1:1 reaction) = $3500 \times 10^6 / 138 = 2.536 \times 10^7$ (1)

mass of phenol needed = $2.536 \times 10^7 \times 94 = 2384$ tonnes (1)

allowing for 45% yield = $2384 \times 100 / 45 = 5298 / 5300$ (tonnes) (1)

allow 5297.5–5300

allow *ecf* throughout

4

[12]

11. methylation stage (can come anywhere)

$\text{CH}_3\text{Cl} / \text{CH}_3\text{Br}$ (1)

$\text{AlCl}_3 / \text{FeBr}_3$ etc. (1)

equation – e.g. $\text{C}_6\text{H}_6 + \text{CH}_3\text{Cl} \rightarrow \text{C}_6\text{H}_5\text{CH}_3 + \text{HCl}$ (1)

intermediate name or unambiguous structure (1)

4 marks

*intermediates and equations will vary if methylation is done
after nitration or reduction*

nitration stage

(conc) H_2SO_4 (1)

(conc) HNO_3 (1)

equation – e.g.: $\text{C}_6\text{H}_5\text{CH}_3 + \text{HNO}_3 \rightarrow \text{C}_6\text{H}_4(\text{CH}_3)\text{NO}_2 + \text{H}_2\text{O}$ (1)

intermediate – name or unambiguous structure (1)

4 marks

reduction stage

tin/iron (1)

HCl (1)

equation – e.g.: $C_6H_4(CH_3)NO_2 + 6[H] \rightarrow C_6H_4(CH_3)NH_2 + 2H_2O$

or with H^+ also on left to give $C_6H_4(CH_3)NH_3^+$ (1)

3 marks

allow other suitable reducing agents:

Quality of Written Communication mark for a well organised answer with the three stages clearly distinguished and sequenced (1)

1 mark

12

[12]

12. (a) any two of ...
fibres / dyes / explosives / pharmaceuticals etc (1)(1) 2
allow any specific examples as long as they do involve aromatic nitro or amine groups – eg NOT nylon, fertiliser etc
- (b) temp 50-60° (1)
concentrated (acids) (1) 2
allow abbreviations for concentrated
- (c) $C_6H_6 + HNO_3 \rightarrow C_6H_5NO_2 + H_2O$
reactants (1) products (1) 2
allow a balanced equation for multiple nitration at any positions
- (d) (i) a pair of electrons ... (1)
... (electrons) move / transferred /
a (covalent) bond breaks/forms (1) 2
- (ii) it accepts a pair of electrons (from the benzene) (1) 1
NOT a 'lone' pair
- (iii) H^+ (on the ring) is replaced by NO_2^+ (1) 1
allow 'substitutes'
ignore + charges
- (iv) it is not used up / reformed at the end AW (1) 1

(e) π -bonding electrons are delocalised (1)

six π -electrons in benzene (1)

four π -electrons in the intermediate (1)

π -electrons are not over one carbon atom /

over **five** carbon atoms / p-orbitals in the intermediate (1)

this must be stated in words to compare benzene and the intermediate

π -electrons are over the **complete** ring / **all around** the ring

all six carbon atoms/ p-orbitals overlapping (1)

Quality of written communication

for at least two sentences/statements with legible text and correct spelling, punctuation and grammar (1)

6

[17]