

1. Two chemical tests are carried out on an aqueous solution of an aromatic organic compound **Y**.

The results of the tests are shown below.

<b>Test</b>	$\text{Br}_2(\text{aq})$	$\text{Na}_2\text{CO}_3(\text{aq})$
<b>Observation</b>	decolourised	effervescence

What is the minimum number of C atoms in **Y**?

- A** 6  
**B** 7  
**C** 8  
**D** 9

Your answer

[1]

2. Bromine is reacted separately with nitrobenzene and phenylamine.


Which organic products are likely to form?

	<b>Product from nitrobenzene</b>	<b>Product from phenylamine</b>
<b>A</b>	2-bromonitrobenzene	2-bromophenylamine
<b>B</b>	2-bromonitrobenzene	3-bromophenylamine
<b>C</b>	3-bromonitrobenzene	2-bromophenylamine
<b>D</b>	3-bromonitrobenzene	3-bromophenylamine

Your answer

[1]

3. The structure of a molecule that is used as a pain reliever is shown below.



Which statement about this molecule is **not** true?

- A It has the molecular formula  $C_8H_9NO_2$ .
- B It reacts with bases to form salts.
- C It has a ketone functional group.
- D It can be hydrolysed with aqueous acid.

Your answer

[1]


4. Which of the following support(s) the delocalised model for benzene rather than the Kekulé model?
- 1: Benzene is less reactive than cyclohexene
  - 2: A benzene molecule has a planar, hexagonal structure
  - 3: The enthalpy change of hydrogenation of benzene is more exothermic than predicted from the Kekulé structure
- A 1, 2 and 3
- B Only 1 and 2
- C Only 2 and 3
- D Only 1

Your answer

[1]

5. A student investigates reactions of aromatic compounds.

(a) The student first carries out the reaction shown below.



- (i) The student obtains a very low yield of compound E.  
The student obtains a much higher yield of a different organic product with molecular formula  $\text{C}_{14}\text{H}_{22}\text{O}_2$ .

Suggest an identity for the organic product  $\text{C}_{14}\text{H}_{22}\text{O}_2$  and draw its structure below.

[1]

- (ii) The student is told by a friend that the  $\text{FeCl}_3$  catalyst is not needed because quinol is more reactive than benzene.

Explain why the student's friend is correct.

You may draw a diagram to support your answer.

.....

.....


.....

.....

.....

..... [3]

- (b) 4-Nitrobenzoic acid is an important compound in chemical synthesis. The flowchart below shows a synthesis involving 4-nitrobenzoic acid.



- (i) State suitable reactant(s) and conditions for **step 1**.

[1]

- (ii) In **step 2**, the  $-\text{NO}_2$  group in compound **F** is reduced by tin and concentrated hydrochloric acid.


Write an equation for the reduction of compound **F**.

Show the structures of any organic compounds involved.

[2]

6. This question is about the chemistry of aromatic compounds.
- (a) Benzoic acid can be nitrated by concentrated nitric acid in the presence of concentrated sulfuric acid as a catalyst, as shown in **Equation 17.1**.

The organic product of this reaction is 3-nitrobenzoic acid.



- (i) Outline the mechanism for this nitration of benzoic acid.

Show how  $\text{H}_2\text{SO}_4$  behaves as a catalyst.



(b) A student investigates the relative ease of nitration of phenol, benzene, and benzoic acid.



The student finds that the conditions required for the nitration of each compound are different, as shown in **Table 17.1**.

Compound	phenol	benzene	benzoic acid
Conditions required for nitration	Dilute HNO <sub>3</sub> 20 °C No catalyst	Concentrated HNO <sub>3</sub> 55 °C H <sub>2</sub> SO <sub>4</sub> catalyst	Concentrated HNO <sub>3</sub> 100 °C H <sub>2</sub> SO <sub>4</sub> catalyst

**Table 17.1**

(i) State the trend in the relative ease of nitration of phenol, benzene, and benzoic acid.

.....

.....

..... [1]

(ii) Apply your knowledge of the bonding in arenes to explain the trend in part (b)(i).

.....

.....

.....

.....

.....


.....

..... [3]



(c) A student synthesises 3-bromophenylamine, shown below, starting from nitrobenzene.

- (i) Complete the flowchart showing the structure of the intermediate and the **formulae** of the reagents for each stage.



[3]

- (ii) Another student attempts the same synthesis but carries out reduction **before** bromination. The student was surprised to find that two structural isomers of 3-bromophenylamine had been formed instead of the desired organic product.

Explain this result and suggest the structures of the two isomers that formed.

Explanation .....


.....

.....

Structures

[3]

7. Benzene reacts with an organic reagent in the presence of a halogen carrier to form phenylethanone.



Which organic reagent is required?

- A  $\text{CH}_3\text{CH}_2\text{OH}$
- B  $\text{CH}_3\text{CHO}$
- C  $\text{CH}_3\text{COCl}$
- D  $\text{CH}_3\text{COOH}$

Your answer

[1]

8. Which statement(s) support(s) the delocalised model for the structure of benzene?
- 1 All carbon–carbon bonds have the same length.
  - 2 The enthalpy change of hydrogenation of benzene is less exothermic than expected.
  - 3 Bromine reacts with benzene less readily than with cyclohexene.
- A** 1, 2 and 3
- B** Only 1 and 2
- C** Only 2 and 3
- D** Only 1

Your answer

**[1]**



(ii) Explain why phenol is nitrated more readily than benzene.

.....

.....

.....

.....

.....

.....


.....

.....

.....


..... [3]

(b) Methylbenzene reacts with sulfur trioxide,  $\text{SO}_3$ , to form **D**, shown below.







The electrophile in this reaction is  $\text{SO}_3$ .

Complete the mechanism for the formation of **D**.  
Show curly arrows and the structure of the intermediate.



10. Phenol reacts with bromine.

Which is the **least** likely organic product?

<b>A</b>	
<b>B</b>	
<b>C</b>	
<b>D</b>	

Your answer

[1]

11. Which chemical(s) can react with phenol?

- 1 Potassium hydroxide
- 2 Ethanoyl chloride
- 3 Nitric acid

- A** 1, 2 and 3
- B** Only 1 and 2
- C** Only 2 and 3
- D** Only 1

Your answer


[1]






- (b) Benzene can be used as the starting material for the synthesis of compounds **D** and **E**, shown below.

In the diagrams  $C_6H_5$  is a phenyl group.





**compound D**



**compound E**

Compounds **D** and **E** can be converted into polymers.

- (i) Draw **two** repeat units of these polymers.



[3]


- (ii) State the **type** of polymer formed from compounds **D** and **E**.

From compound **D** .....

From compound **E** .....

[1]

- (iii) In the synthesis of compounds **D** and **E**, benzene is first reacted with ethanoyl chloride,  $\text{CH}_3\text{COCl}$ , to form phenylethanone, shown below.



**phenylethanone**

The reaction takes place in the presence of aluminium chloride,  $\text{AlCl}_3$ , which acts as a catalyst.


In the mechanism for this reaction,

- ethanoyl chloride first reacts with aluminium chloride to form the  $\text{CH}_3\text{-C}^+=\text{O}$  cation
- the  $\text{CH}_3\text{-C}^+=\text{O}$  cation then behaves as an electrophile.

Complete the mechanism for the reaction.


Include equations to show the role of the  $\text{AlCl}_3$  catalyst, relevant curly arrows and the structure of the intermediate.

Formation of electrophile .....



Regeneration of catalyst .....

(iv) Complete the flowchart for the synthesis of compounds **D** and **E** from phenylethanone.



13. Benzoic acid,  $C_6H_5COOH$ , is added to some foods as a preservative.

A student prepares benzoic acid as outlined below.

**Step 1** The student mixes  $4.00\text{ cm}^3$  of phenylmethanol,  $C_6H_5CH_2OH$ , (density =  $1.04\text{ g cm}^{-3}$ ) with sodium carbonate and aqueous potassium manganate(VII), as an oxidising agent. The mixture is heated under reflux.

**Step 2** The resulting mixture is cooled and then acidified with concentrated  $HCl$ . Impure crystals of benzoic acid appear.

**Step 3** The student recrystallises the impure crystals to obtain  $1.59\text{ g}$  of pure benzoic acid.

(a) In **Step 1**, sodium carbonate,  $Na_2CO_3$ , makes the reaction mixture alkaline.

Write an ionic equation to show how carbonate ions form an alkaline solution in water.

..... [1]

(b) In **Step 2**, explain why the mixture must be acidified so that crystals of benzoic acid appear.

.....

.....

.....

..... [1]

(c) Write the overall equation for the preparation of benzoic acid from phenylmethanol.

Use [O] for the oxidising agent.

..... [1]


(d) Calculate the percentage yield of benzoic acid.

Give your answer to **3** significant figures.

percentage yield = ..... % [3]



14. Dettol<sup>®</sup> is a disinfectant containing the antiseptic chloroxylenol, shown below.



**chloroxylenol**

(a) Chloroxylenol is a weak Brønsted–Lowry acid.

(i) What is the systematic name of chloroxylenol?

..... [1]

(ii) Predict the number of peaks in a <sup>13</sup>C NMR spectrum of chloroxylenol.

..... [1]

(iii) Name the functional group responsible for the acidity of chloroxylenol and describe a simple test which would confirm the presence of this group.

Functional group .....

Test .....

.....

..... [2]

(iv) A student measures the pH of the contents in a bottle of Dettol<sup>®</sup> as 5.14.

The label on the bottle shows the percentage of chloroxylenol in Dettol<sup>®</sup> as 4.80% i.e. 100 cm<sup>3</sup> of Dettol<sup>®</sup> contains 4.80 g of chloroxylenol.

Assume the following:

- Chloroxylenol is the only acidic component in Dettol<sup>®</sup>.
- Chloroxylenol is a weak monobasic acid.
- The density of Dettol<sup>®</sup> is 1.00 g cm<sup>-3</sup>.

Write the equation, using molecular formulae, for the acid dissociation of chloroxylenol.

Calculate the acid dissociation constant,  $K_a$ , for chloroxylenol.

$$K_a = \dots\dots\dots \text{mol dm}^{-3} \text{ [5]}$$





(iii)  $\alpha$ -Terpineol contains two functional groups.

For each functional group, choose a reagent that reacts with that group **only**.  
Draw the structures for the organic products of the reactions.

Show structures for organic compounds.

Reagent(s) .....

Name of functional group that reacts .....

Structure of organic product

Reagent(s) .....

Name of functional group that reacts .....

Structure of organic product

[4]




15. Which one of the following reacts with ethanoic acid **and** with phenol?

- A Aqueous potassium hydroxide
- B Bromine
- C Calcium carbonate
- D Methanol and an acid catalyst

Your answer

[1]

16. Which compound(s) could be prepared by reacting benzene with an acyl chloride in the presence of a halogen carrier?




- A** 1, 2 and 3  
**B** Only 1 and 2  
**C** Only 2 and 3  
**D** Only 1

Your answer

[1]

17 Compounds **B** and **C**, shown below, are unsaturated hydrocarbons containing nine carbon atoms.



(a) Compound **B** reacts with chlorine at room temperature, but compound **C** requires the presence of a halogen carrier.

In both reactions, the organic compound reacts with chlorine in a 1:1 molar ratio.

(i) Draw the structures of the organic product of each reaction.

Organic product with <b>B</b>	Organic product with <b>C</b>

[2]

(ii) Explain the relative resistance to chlorination of compound **C** compared with compound **B**.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

[3]

(iii) Outline the mechanism for the reaction of compound **C** with chlorine.

Show the role of the halogen carrier.

[5]


(b) Compound **C** can be prepared by 'trimerisation' of propanone using concentrated sulfuric acid as a catalyst.

Suggest an equation for this reaction, using **molecular** formulae.

..... [3]

(c) An organic chemist is investigating compound **D** for possible use as a medicine.

The chemist proposes a synthesis of compound **D** from compound **C**.




(i) Predict the number of peaks in the  $^{13}\text{C}$  NMR spectra of compounds **C** and **D**.

	Compound <b>C</b>	Compound <b>D</b>
<b>Number of peaks</b>	.....	.....

[2]

(ii) The chemist develops a three-stage synthesis of compound **D** from compound **C**.


Complete the flowchart.  
Show structures for organic compounds.




compound **C**

reagent: .....

catalyst: .....




1. Sn + HCl  
2. Neutralise



compound **D**

reagent: .....



[5]