
(b) Describe	Chatelier's principle. e, and explain, what would happen to the position of the NO ₂ /N ₂ O ₄ um if the following changes are made. ne temperature is increased.	[2
equilibri	um if the following changes are made. ne temperature is increased.	[:
equilibri	um if the following changes are made. ne temperature is increased.	[
equilibri	um if the following changes are made. ne temperature is increased.	[
equilibri	um if the following changes are made. ne temperature is increased.	
(i) Th 		
••••		
		[:
(ii) Th	ne pressure is increased.	
		[2
		l ⁴
(iii) A	catalyst is added.	
		[2

Nitrogen dioxide, NO_2 , and dinitrogen tetroxide, N_2O_4 , take part in the following

1.

2. The diagram below shows the energy distribution of molecules at a particular temperature. E_a represents the activation energy of the reaction.



(i) On the diagram, draw a second curve to represent the energy distribution of the same number of molecules at a higher temperature.

[2]

(ii)	Using your completed diagram,	explain how an	increase in t	emperature	causes
	the rate of reaction to increase				

[2]

[Total 4 marks]

- 3. A group of students investigated the effect of concentration on the rate of a reaction. They used the reaction between magnesium carbonate and dilute hydrochloric acid and measured the rate at which the gas was collected.
 - (a) (i) An incomplete equation for this reaction is given below. Complete the equation by balancing it and inserting state symbols.


$$MgCO_3(....) + HCl(....)MgCl_2(....) + CO_2(....) + H_2O(....)$$

[1]

(ii) Write the ionic equation for this reaction.

[1]

(b) The students added dilute hydrochloric acid to some magnesium carbonate. The students collected the gas and measured the volume, at regular intervals, until after the reaction was complete. They then plotted a graph of their results.



Use collision theory to explain the changes in the rate of the reaction as it proceeds.

[3]

(c) The students repeated the experiment using a weak acid instead of hydrochloric acid.

Assume the concentration of both acids and all other conditions are the same.

(i) On the axes in (**b**), sketch the graph the students obtained from this experiment.

		(ii)	State and explain what effect changing the acid has on the rate of the reaction.	
				[0]
			[Total 9 m	[2] arks]
4.			used as flavourings. They are made by a reversible reaction between a acid and an alcohol.	
			carboxylic acid + alcohol ← ester + water	
	(a)	Give set u	two features of a reversible reaction, when a dynamic equilibrium has been up.	
		1		
		2		
				[2]
	(b)	The	production of esters is catalysed homogeneously by the presence of acids.	
		(i)	What is meant by a catalyst?	
				[1]

(11)	what is meant by nomogeneous?
	[1]
(iii)	Using the fact that acids are needed to catalyse this reaction, deduce the formula of the ion that acts as the catalyst.
	[1]
(iv)	Catalysts do not affect the position of an equilibrium. Explain why not.
	[1]
	[Total 6 marks]

5.	(i)	On the axes below, draw and label the Boltzmann distribution to show the energies of molecules in a gas at a fixed temperature.	
			-
			[2
	(ii)	Use the graph to explain the effect of a catalyst on the rate of a reaction.	
			 [2
			[Total 4 marks

	Ammonia, NH_3 , is made industrially by the Haber process. This is an equilibrium reaction.				
	$N_2(g) + 3H_2(g) \implies 2NH_3(g) \Delta H = -92 \text{ kJ mol}^{-1}$				
(a)	State the pressure and temperature that are used in the Haber process.				
	pressure				
	temperature				
		[2]			
(b)	Describe and explain why these conditions are a compromise between rate and equilibrium.				
		[9]			
	[Total 1	1 marks]			

6.


[2]

7.	Many industrial processes, used to manufacture important chemicals, involve
	equilibrium reactions. Chemists use their understanding of rates of reaction and of
	yields at equilibrium to find the most economic conditions for the reactions.

Chemists were investigating the production of a chemical, X_2Y , that could be formed from X_2 and Y_2 as shown in **equilibrium 3.1** below.

		$2X_2(g) + Y_2(g) \rightleftharpoons 2X_2Y(g)$ equilibrium 3.1	
(a)	State	le Chatelier's principle.	
			[2]
(b)	State	and explain the effect on equilibrium 3.1 of a decrease in pressure on:	
	(i)	the equilibrium position of the reaction,	
			[2]
	(ii)	the rate of the reaction.	

(c) The chemists measured the percentage conversion of X_2 at various temperatures. The results are shown in the graph below.



(i)	Use the graph to	predict the	percentage	conversion	at 350°	C.
-----	------------------	-------------	------------	------------	---------	----

	_	_	_
	•	٦,	,
answer =	•	~/	_
3113WCI -		/	U

(ii) The forward reaction in **equilibrium 3.1** is exothermic. Explain how the graph supports this statement.

•••••	 •••••	 •••••
•••••	 •••••	

[2]

[1]

(a)		e, and explain, the effect of using a catalyst on:	
	(i)	the rate of conversion of X_2 and Y_2 into X_2Y ,	
			[2]
			i-j
	(ii)	the percentage conversion at equilibrium of X_2 and Y_2 into X_2Y .	
			[2]
		[Total 13 ma	

- **8.** The Boltzmann distribution can be used to show the effect of a change in temperature on the rate of a reaction.
 - Draw a labelled Boltzmann distribution diagram.
 - Explain the essential features of your diagram.
 - Using your diagram, explain how an increase in temperature affects the rate of a reaction.

[Total 9 marks]

9.	Limestone contains calcium carbonate, CaCO ₃ .			
		Limestone is an important source of commercially important chemicals such as lime, calcium oxide, CaO.		
	(a)		lcium carbonate is heated strongly in an open container, it decomposes ording to the equation below.	
			$CaCO_3(s) \rightarrow CaO(s) + CO_2(g)$ $\Delta H = +180 \text{ kJ mol}^{-1}$	
		•	gest two reasons why it is necessary to heat the calcium carbonate strongly chieve decomposition.	
		1		
		2		[2]
	(b)	bein	ther sample of calcium carbonate was placed in a closed container before g heated strongly. This allows an equilibrium to be set up. This equilibrium is wn below. $CaCO_3(s) \rightleftharpoons CaO(s) + CO_2(g)$	
		(i)	What can you say about the rates of the forward and reverse reactions when the calcium carbonate starts to decompose?	
				[1]
		(ii)	What can you say about the rates of the forward and reverse reactions when the equilibrium has been established?	
				[1]

(iii)	A valve allowed some of the carbon dioxide to escape.				
	State and explain what happens to the composition of the mixture in the container.				
	[3]				
	[Total 7 marks]				

10. A chemical C is made by reacting chemical A with chemical B in a reversible reaction.A, B and C are all gases under the reaction conditions.

Research chemists wanted to know the optimum conditions to use in the manufacture of **C**. They carried out a series of reactions under different conditions of temperature and pressure. The percentage conversion of **A** at equilibrium is shown in the table below.

pressure / MPa	temperature / °C	% A converted
	350	8
10	450	12
	550	16
	350	11
20	450	21
	550	29
	350	18
40	450	
	550	49

(a)	Suggest the percentage of A that is converted at 450 °C and 40MPa

(b)	(i)	Use the data in the table above to state the effect of increasing pressure on the percentage of A converted.	
			[1]
	(ii)	What can be deduced, from this change, about the total number of moles of reactants A and B compared with the number of moles of product C in the equation for the reaction? Explain how you reached your conclusion.	
			[2]
			[-]
(c)		the data in the table to deduce whether the reaction between A and B is hermic or endothermic. Explain how you reached your conclusion.	
	•••••		[2]
(d)	It wa	as found necessary to use a catalyst in the production of C.	
	(i)	What is meant by a catalyst?	
			[2]

	(ii)	Suggest and explain two reasons why catalysts are used in industrial processes.	
		1	
		2	
			[2]
(e)		ditions were used that should have given a conversion of A of 39%. In the ufacture, using these conditions, it was found that only 20% conversion was eved.	
	Sugg	est why the conversion was much less than theory suggested.	
			[1]
(f)	Give	two economically important processes that use catalysts.	
	proce	ess 1	
	catal	yst used in process 1	
	proce	ess 2	
	catal	yst used in process 2	
		[Total 13 ma	[2] rks]
		[Total 15 ma]


	$CO(g) + 2H_2(g) \rightleftharpoons CH_3OH(g)$ $\Delta H = -129 \text{ kJ mol}^{-1}$	
(i)	Describe and explain how the composition of the equilibrium mixture is affected by increasing the temperature	
	increasing the pressure in the reaction.	
		[4]
		ניו
(ii)	Describe and explain the effect of increasing the pressure on the rate of reaction.	
		[2]
(iii)	The reaction is carried out by passing gaseous reactants over a transition metal catalyst.	
	Name this type of catalysis.	
		[1]

Methanol can be used as a fuel or as the feedstock for a variety of organic compounds. It is manufactured from carbon monoxide and hydrogen.

11.

	(IV)	Suggest and explain the effect of a catalyst on the equilibrium position.	
		[Total 9 mar	[2] ks]
12.	(a)	A group of students were considering the factors that affect the rate of a chemical reaction. One of the students wrote the following.	
		The rate of a chemical reaction increases as the temperature is increased because there are more collisions.	
		Discuss to what extent this statement is true and describe any other factors that should be considered when looking at the effect of temperature on rate of reaction.	
			[6]

- (b) The students drew a Boltzmann distribution for a gas at temperature T_1 .
 - (i) Label the axes on the students' diagram below.



(ii) Sketch, on the diagram, the distribution that would be obtained at a **lower** temperature, T_2 . Label this line T_2 .

[2] [Total 9 marks]

13. Part of the manufacture of ammonia involves the equilibrium below.

$$N_2(g) + 3H_2(g) \Longrightarrow 2NH_3(g)$$
 $\Delta H = -92 \text{ kJ mol}^{-1}$ State le Chatelier's principle.

[Total 2 marks]

14.	A mixture of N_2 and H_2 was made and left to reach equilibrium.		
	Expla	ain how the following changes would affect the time taken to reach equilibriu	m.
	(i)	use of a catalyst	
			[2]
	(ii)	a higher temperature	
			[2]
			i - .
	(iii)	a lower pressure	
			-
			[2]
			[Total 6 marks]

15.	A mixture of N_2 and H_2 was left until it had reached equilibrium as shown below.			
		$N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$	$\Delta H = -92 \text{ kJ mol}^{-1}$	
	At th	at stage, N_2 , H_2 and NH_3 were present in the ed	quilibrium mixture.	
	-	ain how the following changes would affect the a	amounts of N ₂ , H ₂ and NH ₃	
	(i)	use of a catalyst		
				[1]
	(ii)	a higher temperature		
				[2]
	(iii)	a lower pressure		
				[2]
			[Total 5 ma	

16.	This	question is concerned with equilibria that exist between oxides of nitrogen.	
	State	e le Chatelier's principle.	
		т	otal 2 marks]
17.	Nitro	ogen dioxide, NO ₂ , is a brown gas whilst dinitrogen tetroxide, N ₂ O ₄ , is a colourle	266
17.		The following equilibrium between these two gases was set up.	733
		$2NO_2(g) \rightleftharpoons N_2O_4(g) \qquad \Delta H = -58 \text{ kJ mol}^{-1}$	
		cribe, and explain, what you would see after the following changes have been le and the system allowed to reach equilibrium again.	
	(i)	The temperature is increased.	
			[3]
	(ii)	The pressure is increased.	
			[3]
		т	otal 6 marks]

18.	What is a catalyst?	
		 [Total 2 marks]
19.	State le Chatelier's principle.	
		 [Total 2 marks]
20.	State two characteristics of a dynamic equilibrium.	
_0.	1	
	2	 [Total 2 marks]
21.	The following equation represents an equilibrium reaction.	
	$\operatorname{Cr_2O_7}^{2-}(\operatorname{aq}) + \operatorname{H_2O}(\operatorname{I}) \Longrightarrow 2\operatorname{CrO_4}^{2-}(\operatorname{aq}) + 2\operatorname{H}^+(\operatorname{aq})$ orange yellow	
	Use le Chatelier's principle to describe and explain the colour change (if any) the take place when dilute $HCl(aq)$ is added to a solution containing $K_2CrO_4(aq)$.	at might
		 [Total 2 marks]

22.	The following	equation	represents	another e	quilibrium	reaction.

$$2NO_2(g) \rightleftharpoons N_2O_4(g) \Delta H^{\theta} = -58 \text{ kJ mol}^{-1}$$
 brown colourless

Use le Chatelier's principle to describe and explain the colour change (if any) that might take place when

(i)	a mixture of NO ₂ (g) and N ₂ O ₄ (g) is compressed at constant temperature,	
		[2]
(ii)	a mixture of $NO_2(g)$ and $N_2O_4(g)$ is heated at constant pressure.	
		[2]
	[То	otal 4 marks]

In this question, one mark is available for the quality of written communication. 23.

What effect does a catalyst have on the rate of a reaction, and how does it achieve this effect?

[4]

Quality of Written Communication [1]

[Total 5 marks]


24. Diamond and graphite are two allotropes of carbon. Allotropes are different forms of the same element. Diamonds are made industrially from the much cheaper graphite.

Under high pressure the following equilibrium exists.

C(graphite)
$$\rightleftharpoons$$
 C(diamond) $\Delta H^{\theta} = +1.8 \text{kJmol}^{-1}$

substance	Density / gem ⁻³
graphite	2.25
diamond	3.51


Complete the following enthalpy profile diagram for the conversion of graphite (a) into diamond, labelling ΔH° .



[Total 10 marks]

Suggest which allotrope of carbon is thermodynamically more stable. Give a reason for your choice.	
n which allotrope do the carbon atoms take up less space? Give a reason fo our choice.	r
State Le Chatelier's principle.	
Use Le Chatelier's principle and the information given at the start of the quest of deduce the likely conditions of pressure and temperature needed to turn graphite into diamond. Explain your answer.	tion
Dressure	
emperature	

25. The diagram below shows the energy distribution of reactant molecules at a particular temperature. E_a represents the activation energy of the reaction.



(i) On the diagram, draw a second curve to represent the energy distribution of the same number of molecules at a higher temperature.

[2]

(ii) Use your completed diagram to explain how an increase in temperature can cause an increase in the rate of a reaction.

•••••	 	

[2]

[Total 4 marks]

26. The rate of the reaction between hydrogen and oxygen depends on the pressure as well as the temperature.

$$2H_2(g) + O_2(g) \rightarrow 2H_2O(g)$$
 $\Delta H = -286 \text{ kJ mol}^{-1}$

(i)	Describe and explain the effect of increasing the pressure on the rate of this reaction.	
		[2]
		ا2.
(ii)	A sudden rapid increase in the rate of a reaction causes an explosion to occur.	
	Suggest why highly exothermic reactions such as this one are more likely to explode than other reactions.	
		[2]
	[Total 4 m	arks