


Mark Scheme - OA4 Organic Synthesis and Analysis

- 1 (a) Chromophore [1]
- (b) (i) Melting temperature **lower** than literature value / melting occurs over a temperature range [1]
- (ii) Identify percentage or amount of impurities (1)
Identify the number of compounds present or number of impurities (1) [2]
- (c) (i) Acidified potassium dichromate (1)
Heat and distil (1) do not accept 'reflux' [2]
- (ii) M_r of phenylmethanol = 108.08 M_r of benzenecarbaldehyde = 106.06 (1)
100% conversion would be $10.0 \div 108.08 \times 106.06 = 9.815\text{g}$ (1)
86% yield = $9.815 \times 86 \div 100 = 8.44\text{g}$ (1) [3]
- (iii) Two resonances in the range 5.8-7.0 ppm (1)
These are doublets (1)
One **singlet** at around 11.0 ppm (1)
All resonances have the same area (1) [4]
- Total [13]**

- 2 (a) $\text{CH}_3\text{CH}(\text{CH}_3)\text{CH}_2\text{Cl}$ (1) AlCl_3 / FeCl_3 (1) Room temperature / in the dark (1) [3]
- (b) (i) 2,4-DNP (1) Orange precipitate (1) [2]
- (ii) Tollen's reagent (1) Silver mirror with C, no reaction with B (1) [2]
- (c) Optical isomerism is where a molecule and its mirror image are different / non-superimposable (1)
Compound C has a chiral centre / 4 different groups attached to one carbon atom (1)



The two isomers rotate the plane of polarised light in opposite directions (1) [4]

QWC: organisation of information clearly and coherently; use of specialist vocabulary where appropriate (1) [1]

- (d) Dilute acid (1) heat (1) hydrolysis (1) [3]
- (e) Acidified potassium dichromate (VI) (1) / heat (1)

One step reactions are generally better as they have a better yield / there is waste in each stage (1)

Two step process may be cheaper / use more sustainable reagents / may give a better yield in this case / produce less harmful waste materials / potassium dichromate may react with other parts of the molecule as well / may be easier to separate product (1)

Do not credit same idea twice e.g. if 'better yield' gains first mark, a different point is required to gain second mark [4]

QWC: selection of a form and style of writing appropriate to purpose and to complexity of subject matter [1]

Total [20]

3 (a) (i) Sodium / potassium cyanide [1]



(iii) Sulfuric / hydrochloric acid [1]



(v) eg



(vi) LiAlH_4 / H_2 / sodium, ethanol [1]

(vii) The nitrogen atoms act as electron pair donors / proton acceptors [1]


(b) (i) Molecular formula is $\text{C}_4\text{H}_4\text{O}_2$ [1]

(ii) 3 [1]

(iii) $\text{C} = \text{C}$ / alkene [1]

(iv) Two of the (remaining) protons are in equivalent environments (and one is not) / there are CH and CH_2 present [1]

(v) Possibilities




Total [12]

- 4 (a) (i) 48.5 / 49 % [1]
- (ii) Find a use for the calcium sulfate [1]
- (b) Total volume of aqueous sodium hydroxide needed = $\frac{26.40 \times 250}{25.00} = 264.0 \text{ cm}^3$ (1)
- from the graph this is equivalent to 0.011 mole of the acid (1)
- $\therefore M_r$ of the acid = $\frac{\text{mass}}{\text{no. of moles}} = \frac{2.31}{0.011} = 210$ (1)
- $\text{C}_6\text{H}_8\text{O}_7 \cdot n \text{ H}_2\text{O} = 210$
- \uparrow
- 192 $\therefore n = 18$ (1)
- since M_r of water is 18 $n = 1$ (1)
- (c) The two 'ends' of the double bond have different groups bonded to the carbon atoms (of the double bond) / they have different structural formulae, so cannot be stereo / geometric isomers [1]
- (d) eg sodium ethanoate / ethanoic acid (1) methane (1) [2]
- (e) $\text{C}_5\text{H}_8\text{O}_5 \rightarrow \text{CH}_3\text{COCH}_3 + 2\text{CO}_2$ [1]
- (f)
- The structure shows a central carbon-carbon single bond. The left carbon is bonded to two hydrogen atoms (one with a wedge bond, one with a dash bond) and a hydroxyl group (OH) with a dash bond. The right carbon is bonded to two hydrogen atoms (one with a wedge bond, one with a dash bond) and a hydroxyl group (OH) with a wedge bond.
- [1]
- (g) (Fractional) distillation / (preparative) gas chromatography / HPLC [1]
- (h) (i) eg An optically active isomer that will rotate the plane of polarised light / an isomer with a chiral centre [1]
- (ii) An equimolar mixture of both enantiomers (that has no apparent effect on the plane of polarised light) [1]

Total [15]


5 (a) (i) A compound that can rotate the plane of polarised light. [1]

(ii)



[1]

(iii)




[1]

(iv) Reflux / heat with $\text{H}_2\text{O}/\text{H}^+$ [1]

(v) It contains an equal amount of the two enantiomers /
it is a racemic mixture (1)


The rotating effect of one form exactly cancels out the effect of the
other (1) [2]

(b) (i)



[1]

(ii)



[1]

(c) (i) 2-aminopropanoic acid [1]

(ii) Nitrous acid / nitric(III) acid / HNO_2 [1]

(iii) It exists as a zwitterion (1)

strong electrostatic attractions / ionic bonds between different
zwitterions (1) [2]

Total [12]

- 6 (a) (i) $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3 + \text{Cl}_2 \rightarrow \text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{Cl} + \text{HCl}$ [1]
 (ii) $\text{CH}_3\text{CH}_2\text{CH}_2\dot{\text{C}}\text{HCH}_3$ [1]

- (b) (Anhydrous) aluminium chloride / iron(III) chloride allow AlCl_3 / FeCl_3 [1]

- (c) (i) orange / red precipitate [1]

(ii)



(1) $-\text{COCH}_3$ groups in any positions


It must contain a $\text{C}=\text{O}$ group but it is not an aldehyde as it does not react with Tollens' reagent (1) [2]

- (d) (i) (Alkaline) potassium manganate(VII) (solution) allow KMnO_4 / MnO_4^- [1]

- (ii) Dilute acid allow HCl / H^+ [1]

- (iii) Lithium tetrahydridoaluminate(III) / lithium aluminium hydride
allow LiAlH_4 [1]


(iv)



- (e) Only the infrared spectrum of benzoic acid would have a peak at $1650\text{--}1750\text{ cm}^{-1}$ (1)
 This is due to the carbonyl group present in the benzoic acid (1) [2]

Total [12]

7 (a)



[1]



(b) Moles of calcium carbide = $500/64.1 = 7.80$ (1)

Moles of ethyne = 7.80

Volume of ethyne = $7.80 \times 24.0 = 187 \text{ (dm}^3\text{)}$ (1)

[2]

(c) If the process is endothermic left to right then it needs to absorb energy
– hence the high temperature / endothermic reactions need a high temperature [1]



(f) Any two for (1) each
energy costs / cost of **catalyst** / problems of separation of products /
time taken / availability of starting materials / percentage yield /
atom economy / relative health and safety [2]

(g) $\text{C}_6\text{H}_5-\text{C}\equiv\text{C}-\text{CH}_2-\text{CH}_3$ (1) C_7H_8 (1) [2]



(ii) I sulfuric acid / H_2SO_4 / phosphoric acid / H_3PO_4 / Al_2O_3 [1]

II 3-hydroxypropanoic acid does not show a $\text{C}=\text{C}$ absorption at $1620-1670 \text{ cm}^{-1}$ but this is present in propenoic acid [1]

III The $\text{CH}_3-\text{C}(=\text{O})-$ / $\text{CH}_3\text{CH}(\text{OH})$ group is absent [1]

Total [16]

- 8 (a) (i) 2 mol of ethanol gives 1 mol of ethoxyethane (1)


$$\text{Moles of ethanol} = \frac{69}{46} = 1.5$$

$$\therefore \text{Moles of ethoxyethane if theoretical yield} = 0.75$$

$$\therefore \text{Moles of ethoxyethane if 45\% yield} = 0.75 \times 0.45 = 0.34 \quad (1)$$

$$\text{Mass of ethoxyethane} = 0.34 \times 74 = 25\text{g} \quad (1) \text{ allow error carried forward} \quad [3]$$


- (ii) Ethene / C_2H_4 [1]



(1) for correct curly arrows (1) for correct δ^+ and δ^- [2]

- (iv) They need to have an N-H / O-H / F-H bond / a highly electronegative atom bonded to hydrogen [1]

- (b) (i) For example



[1]


Accept any polybrominated species
Do not accept a monobrominated species

- (ii) Bromine decolorised / orange to colourless / white solid [1]

- (c) Reagent Iron(III) chloride solution / FeCl_3 (1)


Observation Purple coloration / solution (1) [2]

- (d) (i) $\text{C}_{10}\text{H}_{12}\text{O}_1$ [1]



[1]

- (e) Displayed formula, for example



(1)


Functional group carboxylic acid (1) [2]

Total [15]

- 9 (a) (i) A [1]
- (ii) D [1]
- (iii) C [1]
- (iv) C [1]
- (b) (i) Nucleophilic substitution [1]
- (ii) The C–Cl bond in chlorobenzene is stronger than in 1-chlorobutane (1) due to delocalization of electron density from the ring with the bond (1)
- OR
- Delocalised electrons in chlorobenzene (1)
repel lone pair of electrons on nucleophile / ammonia (1) [2]
- (iii) $\text{C}_4\text{H}_9\text{NH}_2 + \text{CH}_3\text{COCl} \longrightarrow \text{C}_4\text{H}_9\text{NHCOCH}_3 + \text{HCl}$ [1]
- (iv) I Tin and concentrated hydrochloric acid (1)
- Add sodium hydroxide (after cooling) (1)
- Steam distillation to separate the product (1) [3]
- II $\text{C}_6\text{H}_5\text{NN}^+\text{Cl}^-$ [1]
- III Azo dye / azo compound [1]

Total [13]

- 10 (a) (i) (Fractional) distillation / (preparative) gas chromatography / HPLC / TLC column chromatography / solvent extraction [1]
- (ii) the fragmentation pattern would be different / valid examples given [1]
- (iii) I




- II Heated electrically / by a naked flame with a water bath (1)
 Add compound G to the ethanol until the hot ethanol will (just) not dissolve any more solute (1)
 Filter hot (1)
 Allow to cool (1)
 Filter (1)
 Dry in air / window sill / < 60 °C in an oven (1) [5]

Maximum 4 out of 5 total if second marking point not given
 Note 5 marks maximum here


QWC Information organised clearly and coherently, using specialist vocabulary where appropriate [1]

- (iv) I The amine is reacted with sodium nitrite / HCl(aq) or nitrous acid (1)
 at a temperature of < 10 °C (1) [2]

II



- (b) (i) Nucleophilic addition (1)



Accept a mechanism that shows HCN polarisation and nucleophilic addition as a concerted process

polarisation / charges shown (1) curly arrows on first structure (1)

regeneration of $\text{C}\equiv\text{N}^-$ or capture of H^+ and curly arrow (1)


[4]

- (ii) Chromophores (1)

The colour will be black (1) as the compound absorbs blue / other colours (1)

[3]


- (iii)



[1]

Total [20]

- 11 (a) (i) Both molecules have lone pairs on nitrogen (1)
 The lone pairs can form (coordinate) bonds with H^+ ions (1) [2]
- (ii) Lone pair on N in phenylamine is delocalised over benzene ring (1) therefore less able to accept H^+ (1) [2]
- (iii) I Arrow in first step (1)
 Cation structure in second step (1)
 Arrow in second step (1)




[3]

II (fractional) distillation / steam distillation [1]

III Sn and conc. HCl (1) followed by NaOH (1) [2]

(b) (i)




- (ii) Addition polymerisation makes one product only /
 Condensation produces one product plus a small molecule like water (1)

Addition polymerisation uses one starting material /
 Condensation polymerisation has two different starting materials (1)


Addition polymerisation involves monomer with one functional group /
 Condensation polymerisation involves monomer with two functional groups
 (1)

(max 2) [2]

(c) (i)



(ii)



(iii) Alanine has strong (electrostatic) forces between the zwitterions (1)

Butanoic acid has hydrogen bonding between molecules /
electrostatic forces in alanine are stronger than forces in butanoic acid
(1)

[2]

(iv) Soda lime (1) $\text{CH}_3\text{CH}_2\text{NH}_2$ (1)

[2]

Total [20]