


## Mark Scheme - PI4.2 Entropy and Feasibility of Reactions

- 1
- (a) (i) atomisation of magnesium / vaporisation of magnesium [1]
- (ii) increased ratio positive charge on nucleus: number of electrons [1]
- (iii) is positive because the (negative) electron is repelled by negative species [1]
- (iv) lattice enthalpy is  $-3835(\text{kJ mol}^{-1})$  numerical value (1) negative sign (1) [2]
- (b) (i) gases are more random/ have more disorder / move more freely and therefore have a higher entropy [1]
- (ii)  $\Delta S = 21.8 (\text{JK}^{-1}\text{mol}^{-1})$  [1]
- (iii)  $\Delta G = \Delta H - T\Delta S$  (1) ft from (ii)
- $\Delta G$  must be  $-ve$  if reaction to be spontaneous/ to calculate T make  $\Delta G = 0$  (1)
- $0 = 318000 - T 21.8$   $T = 14587/14600$  (K) (1) [3]
- (c) use of aqueous sodium hydroxide (1)
- white precipitate for all possible ions (1)
- excess aqueous sodium hydroxide – precipitate dissolves for  $\text{Pb}^{2+}$  and  $\text{Al}^{3+}$  (1)
- use of aqueous (potassium) iodide/ hydrochloric acid/ sulfuric acid / soluble chloride/ soluble sulfate (1)
- result – yellow ppt for  $\text{Pb}^{2+} + \text{I}^-$  and no ppt for  $\text{Al}^{3+}$  / white ppt for  $\text{Pb}^{2+} + \text{Cl}^-$  or  $\text{SO}_4^{2-}$
- and no ppt for  $\text{Al}^{3+}$  [result for both needed] (1) [5]
- QWC Organisation of information clear and coherent* (1)
- Use of specialist vocabulary* (1) QWC [2]
- (d) (i) diagram to show central Al, 4 Cl and 4 shared pairs of electrons, all Cl outer electrons, dative pair identifiable [1]
- (ii) chlorination of benzene (1) produces  $\text{Cl}^+$  as electrophile (1)
- OR gives ionic liquids (1) with low vapour pressure/ non-volatile/ do not evaporate
- in use (1)
- OR catalyst (1) in polymerisation of alkenes (1) [2]
- Total [20]

- 2 (a)  $2 \times (0) + 3 \times (-394) - (-826) - 3 \times \Delta H_f^\ominus(\text{CO}) = -23$  (1)
- $2 \times (\Delta H_f^\ominus(\text{Fe})) + 3 \times (\Delta H_f^\ominus(\text{CO}_2)) - (\Delta H_f^\ominus(\text{Fe}_2\text{O}_3)) - 3 \times \Delta H_f^\ominus(\text{CO}) = -23$  (1)
- $-1182 + 826 + 23 = 3 \times \Delta H_f^\ominus(\text{CO})$
- $-333 = 3 \times \Delta H_f^\ominus(\text{CO})$
- $-111 \text{ kJ mol}^{-1} = \Delta H_f^\ominus(\text{CO})$  (1) [3]
- (b) Gases have higher entropies than solids as the molecules have a greater degree of freedom / disorder [1]
- (c) (i)  $\Delta G = \Delta H - T \Delta S = -23 - (298 \times 9/1000)$  (1)
- $= -25.7 \text{ kJ mol}^{-1}$  (1) [2]
- (ii) A reaction is feasible when  $\Delta G$  is negative (1)
- No temperature exists where  $\Delta G$  is positive /  $\Delta G$  is negative at all temperatures (1) [2]
- (iii) Higher temperature used to increase rate of reaction [1]
- Total [9]**

- 3 (a)  $\text{CO} \rightarrow \text{C} +2$        $\text{CO}_2 \rightarrow +4$  (1)
- Increase of (positive) oxidation number = oxidation / reducing agents themselves  
are always oxidised are always oxidised (1)

OR



Oxidation number of iodine reduced, reduction occurring, CO reducing agent (1) [2]


- (b) +2 state becomes more stable down the group and +4 becomes less stable. [1]

- (c) (i) Add (a little) sodium hydroxide solution (1) to each solution.  
A white precipitate (1) of aluminium / lead(II) hydroxide (1) is seen.  
When more sodium hydroxide solution is added these precipitates (dissolve giving  
a colourless solution). (1) [4]

*QWC Legibility of text: accuracy of spelling, punctuation and grammar;  
clarity of meaning.* [1]

- (ii) Yellow precipitate (1)  $\text{Pb}^{2+} + 2 \text{I}^- \rightarrow \text{PbI}_2$  [2]

- (d) (i) The bonding of **aluminium** in the monomer has not completed the octet / suitable  
diagram / 6 electrons in its outer shell (1)  
When the dimer is formed this octet of bonded electrons is formed (1)



- (ii) (As a catalyst) in the chlorination of benzene / making ionic liquids [1]

- (iii) I The number of (gaseous) species is increasing, leading to less order [1]

II For the reaction to be just spontaneous  $\Delta G = 0$  (1)

$$\text{substituting} \quad 0 = 60\,000 - 88 T$$

$$T = 60\,000 / 88 = 682 \text{ K} / 409^\circ\text{C} \quad (1) \quad [2]$$

(e) 
$$K_c = \frac{[[Al(H_2O)_5(OH)]^{2+}(aq)][[H^+](aq)]}{[[Al(H_2O)_6]^{3+}(aq)]}$$

$$\therefore 1.26 \times 10^{-5} = [H^+]^2 / 0.10$$

$$\therefore [H^+]^2 = 1.26 \times 10^{-6} \quad [1]$$

$$\therefore [H^+] = \sqrt{1.26 \times 10^{-6}} = 1.12 \times 10^{-3} / 0.00112 \quad (1)$$

- error carried forward

$$pH = -\log_{10} [H^+] = -\log_{10} 1.12 \times 10^{-3} = 2.95 \quad (1) \quad [3]$$

Total [20]