

Mark Scheme - 1.6 The Periodic Table

1.

- (a) (i) Energy required to remove one mole of electrons from one mole of atoms / to form one mole of positive ions from one mole of atoms (1)
in the gaseous state (to form 1 mol of gaseous ions) (1)
(Accept correct equation) [2]
- (ii) Cross between Na and Mg crosses [1]
- (iii) P only has unpaired electrons, S has a pair of electrons in 3p orbital (1)
Repulsion between the paired electrons makes it easier to remove one of the electrons (1) [2]
- (b) (i) Effective nuclear charge is greater / electron being removed from a positive ion [1]
- (ii) Accept from 6000 to 9000 [1]
- (c) Lines are formed from electron being excited and jumping up to a higher energy level (1)
Falling back down to the $n = 2$ level (1)
Emitting energy / photon of light (1)
Lines become closer since the electron energy levels of a hydrogen atom become closer (1) [4]
- QWC Selection of a form and style of writing appropriate to purpose and to complexity of subject matter [1]

Total [12]

2.

- (a) same number of protons and electrons (1)
0, 1 and 2 neutrons (1) [2]
- (b) (i) 3 energy levels between $n = 2$ and $n = \infty$
becoming closer together
first gap must be $<$ that between $n = 1$ and $n = 2$ [1]
- (ii) any arrow pointing upwards (1)
from $n = 1$ to $n = \infty$ (1) [2]
- (c) (i) visible [1]
- (ii) (not correct because) Balmer series corresponds to energy transitions involving $n = 2$ (1)
for ionisation energy need Lyman series / energy transitions involving $n = 1$ (1) [2]
- (d) (i) $Q(g) \rightarrow Q^+(g) + e^-$ / accept any symbol [1]
- (ii) Group 6 [1]
- (iii) In T there is more shielding (1)
The outer electron is further from the nucleus (1)
The increase in shielding outweighs the increase in nuclear charge / there is less effective nuclear charge (1) [3]
Legibility of text; accuracy of spelling, punctuation and grammar; clarity of meaning QWC [1]
- Total [14]**

3.

- (a) Weighing bottle would not have been washed / difficult to dissolve solid in volumetric flask / final volume would not necessarily be 250 cm^3 [1]
- (b) Pipette [1]
- (c) To show the end point / when to stop adding acid / when it's neutralised [1]
- (d) So that a certain volume of acid can be added quickly before adding drop by drop / to save time before doing accurate titrations / to give a rough idea of the end point [1]
- (e) To obtain a more reliable value [1]
- (f) (i) Moles = $0.730/36.5 = 0.0200$ (1)
Concentration = $0.02/0.1 = 0.200\text{ mol dm}^{-3}$ (1) [2]
- (ii) Moles = $0.2 \times 0.0238 = 0.00476$ [1]
- (iii) 0.00476 [1]
- (iv) $0.00476 \times 10 = 0.0476$ [1]
- (v) $M_r = 1.14/0.0476 = 23.95$ [1]
- (vi) Lithium [1]

- mark consequentially throughout (f)

Total [12]

4.

nitrogen / phosphorus (or any other Group 5 element)

[1]

5.

- (a) Name of any commercially/ industrially important chlorine containing compound e.g. (sodium) chlorate(I) as bleach/ (sodium) chlorate(V) as weedkiller/ aluminium chloride as catalyst in halogenation

- do not accept CFCs

[1]

- (b) (i) $K_c = \frac{[HI]^2}{[H_2][I_2]}$ must be square brackets

[1]

- (ii) $K_c = \frac{0.11^2}{3.11^2} = 1.25 \times 10^{-3}$ follow through error (ft)

[1]

- (iii) K_c has no units ft

[1]

- (iv) when temperature increases K_c increases (1)

this means equilibrium has moved to RHS

/ increasing temperature favours endothermic reaction (1)

therefore ΔH for forward reaction is +ve (1)

(mark only awarded if marking point 2 given)

[3]

- (c) (i) +2

[1]

- (ii) co-ordinate/ dative (covalent)

[1]

- (iii) pink is $[\text{Co}(\text{H}_2\text{O})_6]^{2+}$ **and** blue is $[\text{CoCl}_4]^{2-}$ (1)

(ligand is) Cl^- (1)

(addition of HCl sends) equilibrium to RHS (1)

[3]

- (iv) $[\text{Co}(\text{H}_2\text{O})_6]^{2+}$ shown as octahedral [with attempt at 3D] (1)

$[\text{CoCl}_4]^{2-}$ shown as tetrahedral/ square planar (1)

[2]

Total [14]

6.

(a) (i) Number of moles of HCl = $\frac{80 \times 0.20}{1000}$ = 0.016 (1)

Number of moles of calcium needed = 0.008 (1)

Number of moles of calcium actually used = $\frac{0.40}{40}$ = ~ 0.010 (1)

(∴ calcium is present in excess)

[Calculation could be carried out in grams] [3]

(ii) gas bubbles / effervescence / some calcium 'dissolves' / colourless solution produced [1]

(b) Mass of E in solution at 0 °C = 0.13×2 = 0.26 g (1)

∴ Quantity precipitated = $1.50 - 0.26$ = 1.24 g (1) [2]

(c) (i) Brick red / orange-red [1]

(ii) Cream precipitate (accept off-white precipitate) [1]



(iv) Red / brown solution [1]

(v) Calcium bromide is an ionic compound (1)
and contains Ca^{2+} and Br^- ions (1)
Chlorine reacts with the bromide ions in a redox / displacement reaction (1)
Chlorine is a more powerful oxidising agent / has a greater affinity for electrons than bromine (1)
 $2\text{Br}^- + \text{Cl}_2 \rightarrow \text{Br}_2 + 2\text{Cl}^-$ (1) [5]

QWC: ensure that text is legible and that spelling, punctuation and grammar are accurate so that the meaning is clear [1]

Total [16]

7. (a) (i)

	<i>magnesium nitrate</i>	<i>barium chloride</i>	<i>sodium hydroxide</i>
<i>potassium carbonate</i>	white precipitate	white precipitate	no visible change
<i>sodium hydroxide</i>	WHITE PRECIPITATE	NO VISIBLE CHANGE	
<i>barium chloride</i>	NO VISIBLE CHANGE		

All three correct for 2 marks, two correct for 1 mark [2]

(ii) Name of precipitate: Magnesium carbonate (1)
Ionic equation: $\text{Mg}^{2+} + \text{CO}_3^{2-} \rightarrow \text{MgCO}_3$ (1) [2]

(b) (i) Sodium hydroxide solution would turn blue/purple
[Ignore references to potassium carbonate] [1]

(ii) Potassium carbonate would give a lilac flame
Sodium hydroxide would give a golden yellow flame
Barium chloride would give an apple green flame
(2 for all correct, 1 mark for 2 correct)
1 max if any reference to white flame for magnesium [2]

(iii) Barium chloride (1) White precipitate (1) [2]

(c) (i) Sodium **ions** surrounded by δ^- on oxygen atoms of water (1)
Bromide **ions** surrounded by δ^+ on hydrogen atoms of water (1)
Marks can be obtained from a labelled diagram – must show minimum of two oxygen/hydrogen atoms around sodium/bromide ions [2]

(ii) Observation with sodium bromide cream precipitate (1)
Observation with sodium iodide yellow precipitate (1) [2]

(iii) Reagent: (dilute) ammonia solution (1)
Observation with sodium bromide: precipitate dissolves in part
Observations with sodium iodide: precipitate does not change
both observations required for (1)
[If concentrated ammonia (1) used then sodium bromide will dissolve completely] [2]

(iv) $2\text{NaI} + \text{Br}_2 \rightarrow 2\text{NaBr} + \text{I}_2$ allow ionic equation [1]

Total [16]

8.

(a) (i) 12 [1]

(ii) 14 [1]

(iii) Percentage / abundance / ratio / proportion of each isotope [1]

(b) (i) 0.125 g [1]

(ii) e.g. Cobalt-60 (1) in radiotherapy (1) / Carbon-14 (1) in radio carbon dating (1) / Iodine-131 (1) as a tracer in thyroid glands (1) [2]

(c) (i) Atoms are hit by an electron beam / electrons fired from an electron gun (and lose electrons) [1]

(ii) To be able to accelerate the ions (to high speed) / so that they can be deflected by a magnetic field
- no credit for 'so that *atoms* can be deflected...'
[1]

(iii) They are deflected by a magnetic field / according to the m/z ratio [1]

(d)



1s

2s

2p

3s

3p



[1]

(e) (i) $\text{Mg}_3\text{N}_2 + 6\text{H}_2\text{O} \longrightarrow 3\text{Mg}(\text{OH})_2 + 2\text{NH}_3$ [1]

(ii) moles $\text{Mg}(\text{OH})_2 = 1.75/58.32 = 0.0300$ (1)

moles $\text{Mg}_3\text{N}_2 = 0.0100$ (1)

mass $\text{Mg}_3\text{N}_2 = 0.01 \times 100.9 = 1.01 \text{ g}$ (1) [3]

- must be 3 significant figures to gain third mark

Total [14]

9.

- (a) apparatus in which reaction can occur, e.g. flask/ test tube, and delivery/ rubber tube (1)

apparatus in which to measure volume of gas, e.g. over water with measuring cylinder/ gas syringe (1) [2]

- (b) (i) fewer moles of barium used / barium has a higher A_r [1]

- (ii) reaction faster/ more vigorous/ less cloudy solution formed with barium (1)

because ionisation energy of barium is less/ electrons lost more easily from barium/ barium is lower in the group/ barium hydroxide is more soluble (1) [2]

- (c) flame test (1) brick red for calcium and (apple) green for barium (1)

OR


add sulfuric acid/ sodium sulfate solution/ potassium sulfate solution (1)

white precipitate with Ba^{2+} , less precipitate/ no precipitate with Ca^{2+} (1) [2]

- (d) electrons correct – oxide ion clearly shows that 2 electrons originated from calcium atom (1)
charges correct (1) [2]

- (e) (i) add sulfuric acid/ sodium sulfate solution/ potassium sulfate solution (1)

filter (1)



- (ii) moles Ba = $2/137$ (1)

$$\text{mass } BaSO_4 = \frac{2 \times 233.1}{137} = 3.4 \text{ (g)} \quad (1) \quad [2]$$

Total [14]

10.

- (a) (i) Potassium bursts into flames sodium does not / potassium darts about surface **more** vigorously than sodium [1]
- (ii) 1st ionisation energy decreases as group is descended / as element has higher A_r (1)
- (Atom) becomes larger / outer electron further from nucleus / more shielding / less effective nuclear charge (1) [2]
- (iii) As group descended outer electron more easily lost [1]
- (b) (i) Electronegativity (difference between the atoms) (1)
- The bigger the difference the more likely is an ionic bond / ORA for covalent (1) [2]
- (ii) Ionic: high electron density centred round ions / shown on diagram (1)
- Covalent: high electron density between nuclei/atoms / shown on diagram (1)
- Intermediate: high electron density between nuclei/atoms but higher nearer one of them / ions with electron distortion of negative ion (1) [3]
- (c) (i) Calcium [1]
- (ii) Calcium chloride/ CaCl_2 – error carried forward (ecf) from (i) [1]
- (iii) White precipitate/ solid – ecf from (i) [1]
- (iv) $\text{Ca}^{2+} + 2\text{OH}^- \rightarrow \text{Ca}(\text{OH})_2$ (ignore state symbols) – ecf from (i) [1]
- Penalise incorrect metal once only in (c)

Total [13]