


## Mark Scheme - 3.5 Chemical Kinetics

1.

- (a) (i) tangent drawn at  $t = 40$  (1)  
rate calculated 0.017 to 0.027 (ignore units) (1) [2]
- (ii) as reaction proceeds less collisions (per unit time) occur [1]
- (b) (i) 1<sup>st</sup> order shown by:  
calculation of rates at at least 2 concentrations (1)  
statement rate  $\propto$  concentration (1)  
OR  
constant half-life (1)  
half-life is 24 minutes (1) [2]
- (ii) rate =  $k[\text{N}_2\text{O}_5]$  (1) [1]
- (iii)  $k = \text{rate (from (i))} / [\text{N}_2\text{O}_5]$  (from graph) (1)  
(mark correct numbers – no need to check evaluation)  
units =  $\text{minutes}^{-1}$  (1) ft from (ii) [2]
- (iv) (student A more likely to be correct) reaction is 1<sup>st</sup> order and 1  $[\text{N}_2\text{O}_5]$  involved in rate determining step [1]
- (c) correct curve starting at 100 kPa and becoming horizontal (1)  
horizontal at 250 kPa (1) [2]

**Total [11]**

2.



(b) (i)

	$[\text{NH}_4^+(\text{aq})]/\text{mol dm}^{-3}$	$[\text{NO}_2^-(\text{aq})]/\text{mol dm}^{-3}$	Initial rate/ $\text{mol dm}^{-3} \text{s}^{-1}$
<b>1</b>	0.200	0.010	$4.00 \times 10^{-7}$
<b>2</b>	<b>0.100</b>	0.010	$2.00 \times 10^{-7}$
<b>3</b>	0.200	<b>0.030</b>	$1.20 \times 10^{-6}$
<b>4</b>	0.100	0.020	<b><math>4.00 \times 10^{-7}</math></b>

(1 mark for each correct answer) [3]

(ii)  $k = \frac{4.00 \times 10^{-7}}{0.200 \times 0.010} = 2.0 \times 10^{-4}$  (1)

Units =  $\text{mol}^{-1} \text{dm}^3 \text{s}^{-1}$  (1) [2]

(iii) No change [1]

(iv) Increases

If temperature is increased rate increases (1)

and since concentrations do not change the rate constant must increase  
(or similar) (1) [2]

**Total [10]**

3.

- (a) Lead(II) iodide or  $\text{PbI}_2$  (1) Bright yellow (1) [2]
- (b)  $2\text{Cu}^{2+} + 4\text{I}^- \rightarrow 2\text{CuI} + \text{I}_2$  (1)
- The precipitate is copper(I) iodide (stated or clearly indicated by state symbols) (1)  
[2]
- (c) Bromine has a more positive  $E^\ominus$  than iodine so it is a stronger oxidising agent (1)
- Bromine is able to oxidise iodide (1)
- Bromine has a less positive  $E^\ominus$  than chlorine so it is a weaker oxidising agent (1)
- Bromine is not able to oxidise chloride (1)
- MAX 3
- OR Calculate EMF for each reaction (1 each) and state that positive EMF means reaction is feasible (1) [3]
- QWC Legibility of text, accuracy of spelling, punctuation and grammar, clarity of meaning* [1]
- (d) 1 mark for each two products or observations  
 $\text{KHSO}_4$  HI  $\text{H}_2\text{S}$   $\text{SO}_2$  S  $\text{I}_2$  [MAX 2 for products]
- Yellow solid rotten egg smell steamy fumes
- Black solid or brown solution or purple fumes
- MAX 3 [3]
- (e) (i) Measure time taken for a sudden colour change (1)  
Rate =  $1 \div \text{time}$  (1) [2]
- (ii) I. pH 1 has a concentration of  $\text{H}^+$  ten times higher than pH 2. [1]
- II. Order with respect to  $\text{H}_2\text{O}_2 = 1$  (1)  
Order with respect to  $\text{I}^- = 1$  (1)  
Order with respect to  $\text{H}^+ = 0$  (1) [MAX 2 for the stated orders]  
Rate =  $k[\text{H}_2\text{O}_2][\text{I}^-]$  (1) [3]
- III.  $k = 0.028$  (1)  $\text{mol}^{-1}\text{dm}^3 \text{s}^{-1}$  (1) [ecf from rate equation] [2]
- IV. Rate equation is unchanged and increasing temperature increases the value of the rate constant [1]

**Total [20]**

4.

- (a)  $1 \text{ dm}^3$  at  $20^\circ\text{C}$  contains  $52.9 \text{ g}$  and at  $0^\circ\text{C}$  it contains  $17.5 \text{ g}$  (1)  
 $\therefore$  amount crystallised =  $52.9 - 17.5 = 35.4 \text{ g}$  (1) [2]

- (b) (i) 2 mol of  $\text{K}_2\text{S}_2\text{O}_8$  give 1 mol of  $\text{O}_2$   
2 mol of  $\text{K}_2\text{S}_2\text{O}_8$  give  $29.0 \text{ dm}^3$  of  $\text{O}_2$  (1)  
 $\therefore$  0.1 mol of  $\text{K}_2\text{S}_2\text{O}_8$  gives  $29.0/20 = 1.45 \text{ dm}^3$  of oxygen (1) [2]

- (ii) Measure the volume of oxygen produced at specified time intervals /  
Measure the pH of the solution at specified time intervals [1]

- (c) (i) An (inert) electrode that is used to carry the charge / current / electron flow [1]

- (ii) A comment on the relative values (e.g. the persulfate system is the more positive of the two systems) (1)  
The more positive 'reagent' / persulfate ions acts as the oxidising agent, accepting electrons via the external circuit (1)  
- must have the first mark to get second [2]

- (d) (i) The experiments show that both the concentrations of iodide and persulfate have doubled (1) therefore the initial rate should increase four times  
 $4 \times 8.64 \times 10^{-6} = 3.46 \times 10^{-5}$  (1) [2]

- (ii) Rate =  $k [\text{S}_2\text{O}_8^{2-}] [\text{I}^-]$  (1)  
$$\therefore k = \frac{8.64 \times 10^{-6}}{0.0400 \times 0.0100}$$
$$= 0.0216 \text{ (1) dm}^3 \text{ mol}^{-1} \text{ s}^{-1} \text{ (1)}$$
 [3]

- (iii) In the rate equation one  $\text{S}_2\text{O}_8^{2-}$  ion reacts with one  $\text{I}^-$  ion.  
The rate-determining step therefore has to have 1 mole of each reacting, as (only) seen in step 1 [1]

Total [14]

5.

- (a) Plotting (2)  
Best fit line (1) [3]
- (b) (i) C (1)  
Curve steeper (1) [2]
- (ii) Concentration of acid is greatest [1]
- (c)  $44 \text{ cm}^3 (\pm 1 \text{ cm}^3)$  [1]
- (d) Moles Mg =  $0.101/24.3 = 0.00416$  (1)  
Moles HCl =  $2 \times 0.02 = 0.04$  (1) [2]
- (e) (i) Mg is not the limiting factor /  
Mg now in excess / HCl not in excess [1]
- (ii) Moles acid =  $0.5 \times 0.04 = 0.02$  (1)  
Volume  $\text{H}_2 = 0.01 \times 24 = 0.24 \text{ dm}^3$   
- correct unit needed (1) [2]
- (f) Lower the temperature of the acid (1)  
Reactants collide with less energy (1)  
Fewer molecules that have the required activation energy (1)[3]
- or Use pieces of magnesium (1) less surface area (1) less chance  
of successful collisions (1)
- QWC Selection of a form and style of writing appropriate to purpose  
and to complexity of subject matter. [1]

Total [16]

6.

- (a) Use weighing scales to weigh the metal oxide (1)  
Use measuring cylinder to pour hydrogen peroxide solution and water into a conical flask (1)  
Immerse flask in water bath at 35 °C (1)  
Add oxide to flask and connect flask to gas syringe (1)  
Measure volume of oxygen every minute for 10 minutes / at regular time intervals (1)
- (any 4 of above, credit possible from labelled diagram) [4]
- (b) Oxide **A** because reaction is faster [1]
- (c) (i) 18 cm<sup>3</sup> [1]  
(ii) 10 cm<sup>3</sup> [1]
- (d) Concentration of hydrogen peroxide has decreased (1)  
reaction rate decreases / fewer successful collisions (1) [2]
- (e) All the hydrogen peroxide has decomposed / the same quantity of hydrogen peroxide was used [1]
- (f) 25 cm<sup>3</sup> [1]
- (g) Reaction will take less time (1)  
Reactants collide with more (kinetic) energy (1)  
More molecules have the required activation energy (1) [3]
- QWC Selection of a form and style of writing appropriate to purpose and to complexity of subject matter* [1]
- Total [15]**

7.

(a) (i) He may have lost carbon dioxide through leaks, this would have given a lower volume than expected. (1)  
He used lower concentration of acid / diluted the acid with water and the rate of carbon dioxide evolution was slower than expected. (1) [2]

(ii) The concentration of acid is higher in the first half (1) the collision rate is higher (1) [2]

(iii) eg  $k = \frac{V}{T}$  (1)  $\therefore k = \frac{130}{298}$  / 0.436

$$\therefore V = 0.436 \times 323 = 141 \text{ (cm}^3\text{)} \quad (1)$$

$$\text{or } \frac{V_1}{V_2} = \frac{T_1}{T_2} \quad (1) \quad \therefore V_1 = \frac{323 \times 130}{298} = 141 \text{ (cm}^3\text{)} \quad (1) \quad [2]$$

(b) (i) 260 (cm<sup>3</sup>) [1]

(ii) 0.45 (g) (0.43–0.48) [1]

(c) The diagram shows two reasonable distribution curves with T<sub>2</sub> flatter and 'more to the right' than T<sub>1</sub>. (1)  
Activation energy correctly labelled, or mentioned in the writing (1)  
Fraction of molecules having the required activation energy is much greater at a higher temperature (thus increasing the frequency of successful collisions) (in words) (1) [3]

*The candidate has selected a form and style of writing that is appropriate to purpose and complexity of the subject matter QWC* [1]

(d) Place the mixture on a balance and measure the (loss in) mass (1)  
at appropriate time intervals (1)

OR BY OTHER SUITABLE METHOD

eg. sample at intervals / quench (1) titration (1) [2]

**Total [14]**

8.

- (a) to increase rate of reaction / to increase surface area [1]
- (b)  $\text{MgCO}_3 + 2\text{HCl} \rightarrow \text{MgCl}_2 + \text{CO}_2 + \text{H}_2\text{O}$  (ignore state symbols) [1]
- (c) rate starts fast and gradually slows (1)  
because concentration becomes less so fewer collisions (per unit time) /  
less frequent collisions / lower probability of collisions (1)  
at time = 17/18 min rate = 0 (1) [3]
- (d) all the solid would all have disappeared / if more carbonate is added further  
effervescence is seen [1]
- (e) (i) volume  $\text{CO}_2 = 200 \text{ cm}^3$  (1)  
moles  $\text{CO}_2 = 200 / 24000 = 0.008333 = \text{moles MgCO}_3$  (1)  
[minimum 2 sf] [2]
- (ii) mass  $\text{MgCO}_3 = 0.008333 \times 84.3 = 0.702 \text{ g}$  (1)  
 $\% \text{ MgCO}_3 = \frac{0.702}{0.889} \times 100 = 79.0\% / 79\%$  [2]
- (e) carbon dioxide is soluble in water / reacts with water (1)  
volume collected less therefore % / moles of  $\text{MgCO}_3$  less (1) [2]
- (f) use of 40.3 and 84.3 (1)  
atom economy =  $40.3 / 84.3 \times 100 = 47.8\%$  (1) [2]

**Total [14]**