# 

Please write clearly in	n block capitals.		
		l	· · · · · · · · · · · · · · · · · · ·
Centre number		Candidate number	
_			
Surname			
Forename(s)			
Candidate signature			

Morning

# GCSE CHEMISTRY

Foundation Tier Paper 2

Wednesday 12 June 2019

## Materials

For this paper you must have:


- a ruler
- a scientific calculator
- the periodic table (enclosed).

## Instructions

- Use black ink or black ball-point pen.
- Fill in the boxes at the top of this page.
- Answer **all** questions in the spaces provided. Do not write outside the box around each page or on blank pages.
- Do all rough work in this book. Cross through any work you do not want to be marked.
- In all calculations, show clearly how you work out your answer.

#### Information

- The maximum mark for this paper is 100.
- The marks for questions are shown in brackets.
- You are expected to use a calculator where appropriate.
- You are reminded of the need for good English and clear presentation in your answers.




8462/2F

Time allowed: 1 hour 45 minutes



IB/G/Jun19/E12








	Answer <b>all</b> questions in the spaces provided	I.	Do not wr outside th box
0 1	This question is about drinking water.		
	There are two main steps in producing drinking water from	n fresh water.	
0 1.1	Draw <b>one</b> line from each step to the reason for the step.	[2 marks]	
	Step	Reason for step	
		Desalination	
	Filtration	Improve taste	
		Increase pH	
	Sterilisation	Kill bacteria	
		Remove solids	
01.2	Which <b>two</b> substances are used to sterilise fresh water?	[2 marks]	
	Tick (✓) <b>two</b> boxes.		
	Ammonia		
	Chlorine		
	Hydrogen		
	Nitrogen		
	Ozone		

0 3

box





IB/G/Jun19/8462/2F

			Do not write outside the box
0 1.4	Scientists tested two water samples from the drinking water supply. The scientists tested one sample for aluminium ions and the other sample for sulfate		
	ions.		
	Draw <b>one</b> line from each ion to the compound needed to identify the ion. [2 marks]		
	lon	Compound needed to identify ion	
		Barium chloride	
	Aluminium ion	Copper sulfate	
		Silver nitrate	
	Sulfate ion	Sodium hydroxide	
		Sulfuric acid	
01.5	How could pure water be proc solids?	duced from drinking water that contained dissolved [1 mark]	
	Tick (✓) <b>one</b> box.		
	Chromatography		
	Cracking		
	Distillation		
	Sedimentation		9



Turn over ►

02	Some central heating boilers use methane as a fuel.	Do not w outside t box
	Carbon monoxide detectors are placed near central heating boilers.	
02.1	Which <b>three</b> properties of carbon monoxide make it necessary to use carbon monoxide detectors?	
	Choose answers from the box. [3 marks]	
	acidic alkaline colourless corrosive	
	insoluble odourless toxic	
	1	
	2	
	3	
02.2	Complete the sentence.	
	[1 mark] Methane produces carbon monoxide when burning in a limited supply of	
02.3	8 g of methane has a volume of 12 dm <sup>3</sup> at room temperature and pressure.	
	Calculate the mass of 36 dm <sup>3</sup> of methane. [2 marks]	
	Mass = g	




02.4	Most methane is obtained from natural gas, which is a fossil fuel.		Do not write outside the box
	Methane can also be produced renewably.		
	Which <b>two</b> are renewable sources of methane?	[2 marka]	
	Tick (✓) <b>two</b> boxes.	[2 marks]	
	Animal waste		
	Food in landfill		
	Nitrogen in the air		
	Non-biodegradable plastics		
	Scrap iron		
			8
	Turn over for the next question		



● 3       Hydrogen is a raw material in the Haber process.         Hydrogen is produced from methane.         The word equation for the reaction is:         methane + steam ⇒ carbon monoxide + hydrogen         ● 3.1       How can you tell that the reaction is reversible?         [1 mark]	Hydrogen is produced from methane. The word equation for the reaction is: methane + steam ≓ carbon monoxide + hydrogen	[1 mark]
The word equation for the reaction is:       methane + steam ≠ carbon monoxide + hydrogen         0 3 . 1       How can you tell that the reaction is reversible?         0 3 . 2       The forward reaction is endothermic.         Name the type of energy change in the reverse reaction.       [1 mark]         0 3 . 3       A nickel catalyst is used in this reaction.         Why is a catalyst used in this reaction.       [2 marks]         Tick (<) two boxes.	The word equation for the reaction is: methane + steam ≓ carbon monoxide + hydrogen	[1 mark]
methane + steam = carbon monoxide + hydrogen   0 3 .1   How can you tell that the reaction is reversible?   [1 mark]     0 3 .2   The forward reaction is endothermic.   Name the type of energy change in the reverse reaction.   [1 mark]     0 3 .3   A nickel catalyst is used in this reaction.   Why is a catalyst used in this reaction?   My is a catalyst used in this reaction?   Tick (<) two boxes.	methane + steam ⇒ carbon monoxide + hydrogen	[1 mark]
0 3.1       How can you tell that the reaction is reversible?       [1 mark]         0 3.2       The forward reaction is endothermic.       Name the type of energy change in the reverse reaction.       [1 mark]         0 3.2       A nickel catalyst is used in this reaction.       [1 mark]       [2 marks]         Tick (~) two boxes.       [2 marks]       [2 marks]         To increase the temperature       [1 mark]       [2 marks]         To produce less carbon monoxide       [1 mark]       [1 mark]         To use less energy       [1 mark]       [1 mark]		[1 mark]
Image:	<b>0 3 . 1</b> How can you tell that the reaction is reversible?	[1 mark]
Name the type of energy change in the reverse reaction.       [1 mark]         0 3 . 3       A nickel catalyst is used in this reaction.         Why is a catalyst used in this reaction?       [2 marks]         Tick (✓) two boxes.       [2 marks]         To increase the temperature       [1 mark]         To produce less carbon monoxide       [1 mark]         To use less energy       [1 mark]		
Name the type of energy change in the reverse reaction.       [1 mark]         0 3.3       A nickel catalyst is used in this reaction.         Why is a catalyst used in this reaction?       [2 marks]         Tick (✓) two boxes.       [2 marks]         To increase the temperature       [1 mark]         To produce less carbon monoxide       [1 mark]         To use less energy       [1 mark]		
[1 mark] 0 3.3 A nickel catalyst is used in this reaction. Why is a catalyst used in this reaction? Tick (<) two boxes. To increase the temperature To produce less carbon monoxide To reduce costs To use less energy	<b>0 3</b> . <b>2</b> The forward reaction is endothermic.	
Why is a catalyst used in this reaction?   Tick (✓) two boxes.   To increase the temperature   To produce less carbon monoxide   To reduce costs   To use less energy	Name the type of energy change in the reverse reaction.	[1 mark]
Why is a catalyst used in this reaction?   Tick (✓) two boxes.   To increase the temperature   To produce less carbon monoxide   To reduce costs   To use less energy		
Why is a catalyst used in this reaction?   Tick (✓) two boxes.   To increase the temperature   To produce less carbon monoxide   To reduce costs   To use less energy		
[2 marks]         Tick (✓) two boxes.         To increase the temperature         To produce less carbon monoxide         To reduce costs         To use less energy	<b>0 3</b> . <b>3</b> A nickel catalyst is used in this reaction.	
Tick (<) two boxes.  To increase the temperature  To produce less carbon monoxide  To reduce costs  To use less energy	Why is a catalyst used in this reaction?	[2 marks]
To produce less carbon monoxide	Tick ( $\checkmark$ ) <b>two</b> boxes.	[z marks]
To reduce costs To use less energy	To increase the temperature	
To use less energy	To produce less carbon monoxide	
	To reduce costs	
To use less methane	To use less energy	
	To use less methane	



box





box

Table 1 shows data about four fertilisers, A, B, C and D.


#### Table 1

Fertiliser	Percentage by mass of nitrogen (%)	Percentage by mass of phosphorus (%)	Percentage by mass of potassium (%)
Α	35.0	0.0	0.0
В	21.2	0.0	0.0
с	21.2	23.5	0.0
D	0.0	0.0	52.3



IB/G/Jun19/8462/2F

box








0 4	Titan is a moon of the plane	t Saturn.			
	<b>Table 2</b> shows the percentaatmosphere of the Earth.	ges of some gases in the	atmosphere of Titan and in th		
		Table 2			
	Gas	Percentage of gas in atmosphere (%)			
	Gas	Titan	Earth		
	Nitrogen	98	78		
	Oxygen	Zero	21		
	Methane	1.4	0.0002		
	Argon	0.14	0.9		
	Carbon dioxide	0.0001	0.04		
04.1	Which <b>two</b> gases are prese		on the Earth than on Titan? [1 ma		
		and			



IB/G/Jun19/8462/2F

[1 mark]






	<del>.</del>		Do not write outside the
0 4 . 4		f Saturn because of the greenhouse effect.	box
	How do greenhouse gases trap energy	from the sun? [1 mark]	
	Tick (✓) <b>one</b> box.		
	All wavelengths of radiation are reflected	ed back to the surface of Titan.	
	Long wavelength radiation is reflected b	back to the surface of Titan.	
	Short wavelength radiation is reflected	back to the surface of Titan.	
	As well as methane, the atmosphere of Methane is an alkane and propene is a	Titan contains small amounts of propene gas. n alkene.	
04.5	Bromine water is an orange solution us	ed to identify alkenes.	
	Draw one line from each gas to its effe		
		[2 marks]	
	Gas	Effect on bromine water	
		Forms a blue solution	
		Forms a blue solution	
	Methane	Forms a blue solution Forms a colourless solution	
	Methane	Forms a colourless solution	
	Methane		
	Methane	Forms a colourless solution	
		Forms a colourless solution	




04.6	Propene reacts with water (steam) to make propanol.	Do not write outside the box
	The ratio of the masses of propene and water that react is:	
	propene : water	
	7:3	
	Calculate the mass of propene that reacts with 21 g water. [2 marks]	
	Mass = g	
		9
	Turn over for the next question	
	rum over for the next question	
	Turn over ▶	•



Do not write outside the box







Do not write outside the box

	Choose answers from the bo	DX.	[2 marl		
	air	ammonia	copper		
	limesto	ne	sand		
	The materials used to make t	the soda-lime glass fibres ar	e sodium carbonate,		
		and			
) 5.5	Suggest <b>two</b> reasons why su	urfboards are coated.	[2 marl		
	1				
	2				
	Some surfboards are made from wood.				
	Some surfboards are made f	rom wood.			
	Some surfboards are made for <b>Table 3</b> contains information a wooden surfboard.		ldition polymer surfboard a		
	Table 3 contains information		ldition polymer surfboard a		
	Table 3 contains information	about the materials in an ac	Idition polymer surfboard a Wooden surfboard		
	Table 3 contains information	about the materials in an ac Table 3 Addition polymer			
	Table 3 contains information         a wooden surfboard.	about the materials in an ac Table 3 Addition polymer surfboard	Wooden surfboard		
	Table 3 contains information         a wooden surfboard.         Relative strength	about the materials in an ac Table 3 Addition polymer surfboard 14	Wooden surfboard 38		



box



IB/G/Jun19/8462/2F

14

19

		Do not write outside the
06	This question is about the corrosion of metals.	box
	The corrosion of iron is called rusting.	
0 6 . 1	Plan an investigation to show that both water and air are needed for iron to rust.	
	You should include the results you expect to obtain.	
	Use apparatus and materials from the list:	
	<ul> <li>test tubes</li> <li>stoppers</li> <li>iron nails</li> <li>tap water</li> <li>boiled water</li> <li>drying agent</li> </ul>	
	• oil. [6 marks]	
I		1



Do not write outside the

box

A student investigated how the mass of three iron nails, **A**, **B** and **C**, increased after rusting.

Table 4 shows the student's results.

Nail	Mass of nail before rusting in g	Mass of nail after rusting in g	Increase in mass of nail in g
Α	1.22	1.30	0.08
В	1.25	1.36	x
С	1.24	1.33	0.09

Table 4

06.2	Calculate <b>X</b> in <b>Table 4</b> .	[1 mark]
	X =	g
06.3	Calculate the mean increase in mass of the three iron nails, <b>A</b> , <b>B</b> and <b>C</b> .	
	Use Table 4 and your answer to Question 06.2	[1 mark]
	Mean increase in mass =	g



Turn over ►


8

● 7       Some students investigated the rate of decomposition of hydrogen peroxide. The equation for the reaction is: hydrogen peroxide → water + oxygen         ● 7.1       Complete the sentence. Choose an answer from the box. <b>a burning splint a glowing splint d a pring splint a glowing splint d a burning splint a glowing splint f the students s two other the section</b> This is the method					
hydrogen peroxide → water + oxygen          ① 7.1       Complete the sentence.         Choose an answer from the box.       [1 mark]	0 7	Some studer	nts investigated the rate of decor	mposition of hydrogen perox	ide.
0 7.1       Complete the sentence.         Choose an answer from the box.       [1 mark]         a burning splint       a glowing splint         damp litmus paper       limewater         The students tested the gas produced to show that it was oxygen.       The students used		The equation	for the reaction is:		
Choose an answer from the box.       [1 mark]         a burning splint       a glowing splint         damp litmus paper       limewater         The students tested the gas produced to show that it was oxygen.         The students used			hydrogen peroxide –	→ water + oxygen	
Choose an answer from the box.       [1 mark]         a burning splint       a glowing splint         damp litmus paper       limewater         The students tested the gas produced to show that it was oxygen.         The students used					
a burning splint       a glowing splint         damp litmus paper       limewater         The students tested the gas produced to show that it was oxygen.         The students used	0 7 . 1				
damp litmus paper       limewater         The students tested the gas produced to show that it was oxygen.       The students used		Choose an a	nswer from the box.		[1 mark]
The students tested the gas produced to show that it was oxygen. The students used			a burning splint	a glowing splint	]
<ul> <li>The students used</li> <li>Student A investigated the effect of the particle size of a manganese dioxide catalyst on the rate of the reaction.</li> <li>This is the method used.</li> <li>1. Measure 25 cm<sup>3</sup> hydrogen peroxide solution into a conical flask.</li> <li>2. Add some fine manganese dioxide powder to the conical flask.</li> <li>3. Measure the volume of oxygen produced every 30 seconds for 10 minutes.</li> <li>4. Repeat steps 1 to 3 two more times.</li> </ul>			damp litmus paper	limewater	
<ul> <li>Student A investigated the effect of the particle size of a manganese dioxide catalyst on the rate of the reaction.</li> <li>This is the method used.</li> <li>Measure 25 cm<sup>3</sup> hydrogen peroxide solution into a conical flask.</li> <li>Add some fine manganese dioxide powder to the conical flask.</li> <li>Measure the volume of oxygen produced every 30 seconds for 10 minutes.</li> <li>Repeat steps 1 to 3 two more times.</li> </ul>		The students	s tested the gas produced to sho	w that it was oxygen.	
on the rate of the reaction. This is the method used. 1. Measure 25 cm <sup>3</sup> hydrogen peroxide solution into a conical flask. 2. Add some fine manganese dioxide powder to the conical flask. 3. Measure the volume of oxygen produced every 30 seconds for 10 minutes. 4. Repeat steps 1 to 3 two more times.		The students	sused		
on the rate of the reaction. This is the method used. 1. Measure 25 cm <sup>3</sup> hydrogen peroxide solution into a conical flask. 2. Add some fine manganese dioxide powder to the conical flask. 3. Measure the volume of oxygen produced every 30 seconds for 10 minutes. 4. Repeat steps 1 to 3 two more times.					
on the rate of the reaction. This is the method used. 1. Measure 25 cm <sup>3</sup> hydrogen peroxide solution into a conical flask. 2. Add some fine manganese dioxide powder to the conical flask. 3. Measure the volume of oxygen produced every 30 seconds for 10 minutes. 4. Repeat steps 1 to 3 two more times.					
<ol> <li>Measure 25 cm<sup>3</sup> hydrogen peroxide solution into a conical flask.</li> <li>Add some fine manganese dioxide powder to the conical flask.</li> <li>Measure the volume of oxygen produced every 30 seconds for 10 minutes.</li> <li>Repeat steps 1 to 3 two more times.</li> </ol>				cle size of a manganese dio	kide catalyst
<ul><li>2. Add some fine manganese dioxide powder to the conical flask.</li><li>3. Measure the volume of oxygen produced every 30 seconds for 10 minutes.</li><li>4. Repeat steps 1 to 3 two more times.</li></ul>		This is the m	ethod used.		
<ul><li>3. Measure the volume of oxygen produced every 30 seconds for 10 minutes.</li><li>4. Repeat steps 1 to 3 two more times.</li></ul>		1. Measure 2	5 cm <sup>3</sup> hydrogen peroxide solutio	on into a conical flask.	
4. Repeat steps 1 to 3 two more times.		2. Add some	fine manganese dioxide powde	r to the conical flask.	
		3. Measure tl	ne volume of oxygen produced e	every 30 seconds for 10 min	utes.
5. Repeat steps 1 to 4 with coarse manganese dioxide lumps.		4. Repeat ste	eps 1 to 3 two more times.		
		5. Repeat ste	eps 1 to 4 with coarse manganes	se dioxide lumps.	
		·		·	



0 7 2	The method student <b>A</b> used did <b>not</b> give repeatable results.		Do not write outside the box
	How could student <b>A</b> make the results repeatable?		
	Tick (✓) <b>one</b> box.	[1 mark]	
	Student <b>A</b> should make measurements every 2 minutes.	]	
	Student <b>A</b> should measure the mass of manganese dioxide.		
	Student <b>A</b> should use 50 cm <sup>3</sup> hydrogen peroxide.	]	
	Student <b>A</b> should use a beaker instead of a conical flask.		
	Student <b>B</b> used a method which gave repeatable results.		
0 7.3	How could student <b>B</b> improve the accuracy of these results?	[1 mark]	
	Tick (✓) <b>one</b> box.	[]	
	Calculate a mean but do not include any anomalous results.	]	
	Calculate a mean but do not include the first set of results.		
	Record the results in a table and plot the results on a bar chart.	]	
	Record the results in a table and plot the results on a line graph.		







0 7.5	Fine manganese dioxide powder produces a higher rate of reaction than coarse manganese dioxide lumps.	Do not write outside the box
	Sketch on <b>Figure 5</b> the results you would expect for student <b>B</b> 's experiment with fine manganese dioxide powder.	
	[2 marks]	
07.6	Hydrogen peroxide molecules collide with manganese dioxide particles during the reaction.	
	Why does fine manganese dioxide powder produce a higher rate of reaction than coarse manganese dioxide lumps?	
	[1 mark] Tick (✓) <b>one</b> box.	
	Fine manganese dioxide powder has a larger surface area.	
	Fine manganese dioxide powder has larger particles.	
	Fine manganese dioxide powder produces less frequent collisions.	
	Turn over for the next question	10
	Turn over ►	

i urn over i




Table 5 gives information about some of the fractions.

Table 5

Fraction	Boiling point range in °C
Petroleum gases	Below 30
Petrol	40–110
Kerosene	180–260
Diesel oil	260–320
Heavy fuel oil	320–400
Bitumen	400–450



08.1	Suggest a suitable temperature for the furnace in Figure 6. [1 mark]	Do not writ outside the box
08.2	C Explain why diesel oil collects above heavy fuel oil but below kerosene in the fractionating column. Use Table 5. [2 marks]	
08.3	Suggest <b>two</b> reasons why bitumen is <b>not</b> used as a fuel. [2 marks] 1	
	2	
	Question 8 continues on the next page	




08.4	Petrol contains mainly alkanes.	Do not write outside the box
	Which of the following compounds is an alkane?	
	[1 mark] Tick (✓) one box.	
	C <sub>2</sub> H <sub>4</sub>	
	C <sub>4</sub> H <sub>8</sub>	
	C <sub>6</sub> H <sub>14</sub>	
	C <sub>8</sub> H <sub>16</sub>	
	Large hydrocarbon molecules in the diesel oil fraction are cracked to produce smaller hydrocarbon molecules.	
08.5	Describe the conditions needed to crack hydrocarbon molecules from the diesel oil fraction.	
	[2 marks]	



	Turn over ►	
	Turn over for the next question	11
0 8 . 7	$C_{15}H_{32} \rightarrow C_{12}H_{26} + \_\_\_$	
08.7	Complete the equation for the cracking of $C_{15}H_{32}$	
	[2 marks]	
0 8.6	Explain why large hydrocarbon molecules in the diesel oil fraction are cracked to produce smaller hydrocarbon molecules.	Do not write outside the box








			Do not write outside the box
09.2	The tablet also contains other substances. The substances in tablets are present in fixed amounts.		
	What name is given to mixtures like tablets?	[1 mark]	
09.3	The tablet has a mass of 1.20 g and contains 700 mg of lithium carbonate.		
	Calculate the percentage by mass of lithium carbonate in this tablet.	[3 marks]	
	Percentage by mass of lithium carbonate =	%	10



Turn over ►






1 0	This question is about rate of reaction.	Do not write outside the box
	A student investigated the rate of the reaction between magnesium and dilute hydrochloric acid.	
	The equation for the reaction is:	
	$Mg(s) + 2 HCI(aq) \rightarrow MgCI_2(aq) + H_2(g)$	
10.1	Which state symbol in the equation for the reaction does <b>not</b> represent one of the three states of matter?	
	[1 mark]	
	The student determined the rate of production of hydrogen gas.	
10.2	What <b>two</b> pieces of measuring apparatus could the student use to find the rate of production of hydrogen gas?	
	[2 marks]	
	1	
	2	
	Question 10 continues on the next page	



34


Do not write outside the box





10.4	Give <b>three</b> conclusions that can be drawn about the rate of reaction b	petween	Do not write outside the box
	magnesium and dilute hydrochloric acid in this investigation.		
	Use data from <b>Figure 8</b> and <b>Table 6</b> .	[3 marks]	
	1		
	2		
	3		
10.5	The student repeated the investigation using dilute hydrochloric acid temperature.	at a higher	
	All the other variables were kept the same.		
	Which <b>two</b> statements are correct?	[2 marks]	
	Tick (✓) <b>two</b> boxes.	[]	
	More bubbles were produced in the first 10 seconds.		
	The activation energy for the reaction was higher.		
	The magnesium was used up more quickly.		
	The reaction finished at the same time.		
	The total volume of gas collected was greater.		
	END OF QUESTIONS		11





Copyright © 2019 AQA and its licensors. All rights reserved.





IB/G/Jun19/8462/2F